

## EXPERIMENTAL PAPER

### Preliminary study on the impact of methanolic extract of *Elephantopus scaber* Linn. on hair growth promoting effect in rats

HIMANSHU BHUSAN SAHOO<sup>2</sup>, RAKESH SAGAR<sup>1\*</sup>, SUBRAT KUMAR BHATTAMISRA<sup>3</sup>,  
AMRITA BHAIJI<sup>2</sup>

<sup>1</sup> Department of Pharmacognosy and Phytochemistry  
Dr. H.S. Gour Central University  
Sagar, 470003  
Madhya Pradesh, India

<sup>2</sup> Department of Pharmacology and Toxicology  
Vedica College of Pharmacy  
Bhopal 462021  
Madhya Pradesh, India

<sup>3</sup> Department of Pharmacology  
Roland Institute of Pharmaceutical Sciences  
Berhampur 760010  
Orissa, India

\*corresponding author: e-mail: drrsagar@gmail.com

#### Summary

Methanolic extract of *Elephantopus scaber* leaves was investigated for its role in hair growth in experimental rats. The 2% and 5% solution of prepared formulation using methanolic extract was studied for hair growth on wistar rats for 30 days. Minoxidil (2% solution) was taken as a reference standard. The prepared hair oil formulation of methanolic extract significantly ( $p < 0.01$ ) potentiated the hair growth initiation and completion time with dose-dependent improvement in skin biopsy, hair length and hair weight. *In vitro* hair follicles development study showed the significant ( $p < 0.05$ ) improvement in the initiation of new hair follicles. Above mentioned results indicated that the methanolic extract of *E. scaber* rendered significant hair growth promotive effect.

**Key words:** alopecia, hair growth, *Elephantopus scaber*, minoxidil, MESL, methanolic extract

## INTRODUCTION

Alopecia is a dermatological disorder and distressing symptom [1]. It is common universal problem in primary healthcare practice throughout the world and has been estimated to affect both genders of all races between 0.2% and 2% of the world population [2]. It has been suggested that alopecia could have an indirect effect on psychological and social lifestyle along with self esteem among both genders [3]. The main problems associated with alopecia such as pigmentation problems (fading), dandruff and falling of hair. Regardless from metabolic and hereditary causes, alopecia has been observed as a major side effect of anticancer, immunosuppressant and many other drug treatments [4]. The root hair reflects the growth, water and nutrient uptake and site of infection by fungus which is the primary factor of alopecia [5]. Many people suffer from hair loss or hair thinning despite the development of several medical treatments. Therefore, it is important to develop a novel therapy that could prevent hair loss and enhance the hair growth [6]. The search for treatment results in few drugs of synthetic origin, but their side effects cannot be neglected [7].

*Elephantopus scaber* Linn. (*Asteraceae*) popularly known as Elephant's foot is a wild shrub distributed worldwide in all tropical regions. Phytochemically, the plant has been reported to contain sesquiterpenes lactones deoxyelephantopin, isodeoxyelephantopin and scabertopin [8]. It also contains epifriedelinol, lupeol, stigmaterol [9], elephantopin and triterpenes [10]. The pharmacological properties of the leaf extracts have been evaluated for diuretic [11], anti-inflammatory [12], hepatoprotective [13] and antimicrobial activity [14]. In traditional claims, roots were used as an antipyretic, cardiogenic. Decoction of roots and leaves is used as emollient and given in dysuria, diarrhoea, dysentery, tonic, anthelmintic [15]. It is also used to treat stomachic pain [16] and also possesses hepatoprotective activity [17]. When the extracts of leaves are applied topically, they are employed as to treat eczema and ulcers [18], an antipyretic for treatment of erysipelas, skin infections, and measles [19]. In Siddha system of medicine, the leaves of this plant are used as a thaali for hair in case of hairfall, however, there is no scientific report available on hair growth promoting activity of *Elephantopus scaber*. Therefore, the present study was focused on the scientific investigation of the hair growth potential of *Elephantopus scaber* Linn. leaves.

## MATERIALS AND METHODS

### Plant material

Leaves of *Elephantopus scaber* Linn. were collected from local area of Bhopal, Madhya Pradesh, India in August 2011. The species was identified and authenticated at the Department of Botany, Jiwaji University, Gwalior, Madhya Pradesh, India, where a plant specimen was deposited. The collected leaves were air-dried, reduced to coarse powder using electric blender and stored in an airtight container at room temperature (35°C).

## Preparation of extract

The dried coarse powder (200 g) of leaves was continuously extracted with methanol using Soxhlet apparatus for 48 h. After complete extraction, the methanolic extract was concentrated under reduced pressure at 40°C in a vacuum dryer to obtain dried extract. The percent yield of the extract was found to be 8.74% (w/w).

## Preparation of herbal hair formulation

Accurately weighed 2 g and 5 g of crude extract were taken and mixed in 100 ml of oil base (olive oil) to prepare homogeneous mixture of 2% and 5% hair oil, respectively. Then, the prepared formulation was taken for further study to evaluate hair growth promoting activity.

## Animals

Adult Wistar male albino rats (120–150 g) were used for the study. The experimental protocols used in this study were approved by the Institutional Animal Ethical Committee under CPCSEA (Regd. No 926/ab/06/CPCSEA). All animals were fed with standard laboratory diet and water *ad libitum*. They were housed in an air-conditioned room with 12:12 h light and dark cycle. The room temperature ( $24 \pm 2^\circ\text{C}$ ) and humidity (about 60%) were controlled automatically.

## Primary skin irritation test

Small quantity of 2 % and 5% prepared formulation was applied topically over the preselected area of all the animals and animals were observed for 48 hours for any symptoms of toxicity.

## Drug treatment

The animals were divided into four groups (six animals in each group). Group I was kept as control group (only olive oil), Group II was treated as standard (2% Minoxidil), Group III and IV were treated with 2% and 5% of formulation respectively. Hairs of 1-cm<sup>2</sup> area (dorsal portion) of all animals of each group were removed using hair remover cream to remove all hairs. The treated as well as standard groups (2% Minoxidil) were treated with respective formulations on the denuded area of albino rats once a day for 30 days. The denuded area of rats of each group were unprotected due to free access of light and atmosphere.

## ***In vivo* hair growth analysis**

On completion of 30<sup>th</sup> day treatment, the hair growth status of all the groups was observed visually and data were recorded.

### **Qualitative analysis**

The qualitative analysis of hair growth was evaluated by visual observation of two abovementioned parameters. Initiation time (minimum time required to initiate hair growth) and completion time (minimum time required to cover the denuded skin region with new hair completely). Hair growth initiation and completion time was recorded for each group of animals and compared with control group [20-21].

### **Quantitative analysis**

After 30 days of treatment, skin biopsies were taken from the shaved area and the specimens preserved in 10% formalin buffer [22-23]. Then the tissues were stained and allowed to microscopical analysis having ocular micrometer facility for determination of hair follicles in anagenic and telogenic phase.

### **Hair length analysis**

The hair was plucked randomly from the shaved area of rats from each group. After 30<sup>th</sup> day of the treatment, the length of ten hairs was measured and the average length was determined and compared with animals from control group [24].

### **Hair weight analysis**

After 30 days, the rats of all the groups were sacrificed by cervical dislocation. Dorsal skin area (1 cm<sup>2</sup>) with hairs and without hairs was cut and weighed with the high precision analytical balance. After measuring, hair weight was calculated by subtracting weight of skin from weight of skin with hair [24].

## ***In vitro* hair growth analysis**

### ***Hair follicular analysis***

The hair follicles were isolated from the neonates of albino rat. The neonates were sacrificed by cervical dislocation and the dorsal portion of the skin was dissected

out and washed thoroughly in phosphate buffered saline. The skin was cut into small segments and individually placed in a Petri dish containing PBS. The segments were chopped thoroughly until the intact follicles came out from the skin. The intact hair follicles were isolated using a fine Pasteur pipette in binocular microscope. Individual, freshly isolated hair follicles were placed in separate wells of 96-well plates containing 150  $\mu$ l of Dulbecco's Modified Eagles Medium. Finally 1.5  $\mu$ l of 2% and 5% solution of methanolic extracts of *E. scaber* in dimethyl sulfoxide were added to the corresponding wells and the plates were maintained at 37°C. After 24 h and 72 h of treatment, the status of hair follicle length was measured using a binocular microscope equipped with an eyepiece measuring graticule [25].

## Statistical analysis

Data were expressed as mean  $\pm$  SEM. All data were analyzed statistically using one-way analysis of variance (ANOVA), followed by post-hoc Dunnet's test. Values of  $p < 0.05$  were considered statistically significant.

## RESULTS

### Phytochemical analysis

Phytochemical analysis revealed that methanolic extract of *E. scaber* contains flavonoids, steroids, tannins, terpenes and phenolic compounds.

### Skin irritation analysis

MESL (2% and 5% oil formulation) did not show any sign of irritation i.e. no erythema or edema and no loss of hairs was observed at the site of application till 48 hours.

### *In vivo* hair growth measurement

#### *Qualitative measurement of hair growth*

The times taken for hair growth initiation and completion were observed on shaved area at the end of the course in all the groups. As shown in table 1, the initiation of hair growth was observed on 11<sup>th</sup> day in group-I, 7<sup>th</sup> day in group-II, 10<sup>th</sup> day in group-III and 8<sup>th</sup> day in group-IV. Similarly, the completion of hair growth was found on 26<sup>th</sup> day in group-I, 22<sup>nd</sup> day in group-II, 25<sup>th</sup> and 24<sup>th</sup> day for group-III & IV respectively. MESL (2% and 5% formulation) showed significant ( $p < 0.01$ ) hair growth activity as compared to control group.

Table 1.

Effect of MESL hair formulation on qualitative analysis of hair growth in rats.

S. No.	Treatment	Hair growth (days)	
		Initiation time	Completion time
1	Control	11±0.577	26±0.774
2	Standard (2% minoxidil solution)	7±0.365 **	20±0.632 **
3	2% MESL	10±0.577 *	24±0.632 *
4	5% MESL	8±0.577 *	22±0.774 *

All values were expressed as mean ±SEM; n=6 for each treatment group; MESL: methanolic extracts of *Elephantopus scaber* leaves; \* p <0.01 and \*\* p<0.001 compared with control.

### Quantitative measurement of hair growth

A considerable difference in cyclic phases of hair growth was observed in groups treated with minoxidil and MESL formulation. Cyclic phases of hair growth were markedly effected by the standard drug minoxidil and MESL. After 30 days of treatment, the anagenic population of hair follicles in group I, II, III and IV was found to be 51%, 69%, 59% and 65% respectively. Similarly, the telogenic population of hair follicles in group I, II, III and IV was found to be 49%, 31%, 41% and 35% respectively. Thus, both formulations in treated groups showed significant ( $p<0.01$ ) growth as compared to control. The results are summarized in table 2.

Table 2.

Effect of MESL hair formulation on the percentage of hair growth in different phases after 30 days of application in rats.

S. No.	Treatment	% of Telogen	% of Anagen
1	Control	49±1.751	51±1.732
2	Standard (2% Minoxidil)	31±1.807 **	69±1.932 **
3	2% MESL oil	41±1.932 **	59±1.932 *
4	5% MESL oil	35±1.183 **	65±2.017 **

All values were expressed as mean ±SEM; n=6 for each treatment group; MESL: methanolic extracts of *Elephantopus scaber* leaves; \* p<0.01 and \*\* p<0.001 compared with control.

### Hair length measurement

The length of hair follicles was measured and observed that in group I, II, III and IV the length of hair follicles was 9±0.56 mm, 16±0.577 mm, 12±0.83 mm and 14±0.47 mm, respectively. MESL treated group showed remarkable effect

( $p < 0.01$ ) on length of hair follicles after 30-day treatment. The results are summarized in table 3.

Table 3.

Effect of MESL hair formulation on hair length after 30 days of application in rats

S. No.	Treatment	Hair length (mm)
1	Control	9±0.557
2	Standard (2% Minoxidil)	16±0.577**
3	2% MESL	12±0.830
4	5% MESL	14±0.471**

All values were expressed as mean ± SEM; n=6 for each treatment group; MESL: methanolic extracts of *Elephantopus scaber* leaves; \*  $p < 0.01$  and \*\*  $p < 0.001$  compared with control.

### Hair weight measurement

The weight of newly grown hairs in the test groups and standard group were measured and compared with control group. The weight of hair was significantly ( $p < 0.05$ ) increased, found to be 53 and 58 mg/cm<sup>2</sup> area of dorsal skin for 2% and 5% of MESL hair oil formulation, respectively. Whereas, minoxidil was found to be 53 mg/cm<sup>2</sup> area of dorsal skin of rat (tab. 4).

Table 4.

Effect of MESL hair formulation on hair weight after 30 days of application in rats

S. No.	Treated groups	Weight of hair (mg/cm <sup>2</sup> )
1	Control group	49±2.781
2	Standard (2% Minoxidil)	59±1.807**
3	2% MESL	53±0.577
4	5% MESL	58±1.202**

All values were expressed as mean ± SEM; n=6 for each treatment group; MESL: methanolic extracts of *Elephantopus scaber* leaves; \*\*  $p < 0.001$  compared with control.

### In vitro hair growth measurement

As shown in table 5, the growth of hair follicle was observed under binocular microscope at higher magnification. The MESL in both dose groups showed a significant ( $p < 0.05$ ) increase in length over 72 h in culture against control group (0.13 mm/d). The rate of growth was found to be 0.28±0.02 mm and 0.37±0.02 mm per 72 h in 2% and 5% MESL treated group, respectively.

Table 5.

Effect of hair formulation MESL on *in vitro* hair follicle culture

S. No.	Treated groups	Increase in hair follicle length (mm)	
		After 24 h	After 72 h
1	Control group	0.09±0.018	0.13±0.014
2	DMSO	0.04±0.008	0.08±0.014
3	2% MESL	0.13±0.017*	0.28±0.023*
4	5% MESL	0.17±0.020*	0.37±0.017*

All values were expressed as mean ± SEM; n=6 for each treatment group; MESL: methanolic extracts of *Elephantopus scaber* leaves; \* p<0.01 compared with control.

## DISCUSSION

Methanolic extract of *Elephantopus scaber* hair oil formulation on topical application stimulate the hair growth initiation and completion time. It was observed that the hair growth initiated from the shaved area at the start of the 2<sup>nd</sup> week in all groups, including control, and in the whole denuded area has been covered at the end of the course, in comparison to the control. This may be due to gentle rubbing of the shaved skin while application of extracts. This enhances the blood circulation in local area. Thus, it may exert some effect on hair growth. Androgenetic alopecia is a dihydrotestosterone-mediated process characterized by continuous miniaturization of androgen-reactive hair follicles and accompanied by perifollicular fibrosis of follicular units in histological examination [26, 6]. Retention of late anagenic follicles as well as increase in follicular length and prevention of miniaturization. Therefore, it may be attributed to 5- $\alpha$ -reductase inhibitory activities. The groups treated with formulation produced a greater effect on the length of hair as compared to control group. This may be due to the premature switching of follicles from telogen to anagen phase of hair growth cycle [25]. The *in vitro* study revealed that the leaf extract has direct impact on hair follicles and thus may improve the hair growth. It is expected that several fatty acids, e.g. palmitic, oleic, linoleic, linolenic and arachidonic acids, as well as mixture of these acids show a significant anti-androgenic effect owing to their testosterone 5- $\alpha$ -reductase inhibitory activity [27, 24]. Flavonoids and triterpenoids possess hair growth promoting activity by strengthening the capillary wall of smaller blood vessels supplying hair follicles, improve blood circulation to nourish the hair follicles and thereby promote the hair growth [28, 25]. Flavonoids play a great role by stimulating telogen to anagen phase, a process involved in hair growth, and also cause expressions of some growth factors, such as insulin-like growth factor-1 (IGF-1), vascular endothelial growth factors (VEGF), keratinocyte growth factors (KGF) and hepatocyte growth factors (HGF),

all of which has a stimulatory effects on hair growth [29-31]. Minerals and certain amino acids may be the cause for the significant hair growth activity which not only shows remarkable activity but also devoid of potential side effects as compared to synthetic drugs [32].

Methanolic extract and its water insoluble fraction of green tea showed the presence of phenolic compounds. The EGCG (epigallocatechin-3-gallate), a major constituent of polyphenols of green tea was reported to be useful in the treatment of androgenetic alopecia by selectively inhibiting 5- $\alpha$ -reductase activities [33].

## CONCLUSIONS

In conclusion, the experimental evidence obtained in present laboratory animal study indicates that methanolic extract of *Elephantopus scaber* leaves possesses hair growth promoting effect. This may be due to the presence of triterpenes, polyphenols and flavonoids in the plant extracts. Further pharmacological and biochemical investigations needed to establish the mechanism of action and isolation of phytoconstituents from the extract may provide new directions for the better treatment of alopecia or hair loss.

## ACKNOWLEDGEMENTS

Authors are grateful to Ayushmati Education and Social Society, Bhopal, Madhya Pradesh, India for financial support during the study.

## REFERENCES

1. Vera HP. Treatment of hair loss. *New Engl J Med* 1999; 341:964–973.
2. Bhalearo SS, Salanki NH. Therapeutics approaches to the management of common baldness. *Indian Drugs* 2002; 39:567–573.
3. Sharma M, Banerjee PS. Preparation and evaluation of mixed herbal hair oil formulation. *Phcog J* 2009; 1:154–160.
4. Purwal L, Gupta SPBN, Pande MS. Development and Evaluation of Herbal Formulations for Hair Growth. *E-Journal of Chemistry* 2008; 5:34–38.
5. Kevin JME, Rodney S. Hair physiology and its disorders. *Drug Discovery Today: Disease Mechanisms* 2008; 5: e163–e171.
6. Yoo HG, Kim JS, Lee SR, Pyo HK, Moon HI, Lee JH, Kwon OS, Chung JH, Kim KH, Eun HC, Cho KH. Perifollicular fibrosis: pathogenetic role in androgenetic alopecia. *Biol Pharm Bull* 2006; 29:1246–1250.
7. Bhaumik S, Jyothi MD, Khar A. Differential modulation of nitric oxide production by curcumin in host macrophages and NK cells. *FEBS Lett* 2000; 483:78–82.
8. Paul PHB, Po-Ming H, Hui C, Dominic Chan TW, Bo-Mu W, Thomas CWM, Chun-Tao C. Sesquiterpenes lactone from *Elephantopus scaber*. *Phytochemistry* 1997; 44: 113–116.
9. De Silva LB, Hearth WHMW, Jennings RC, Mahendran M, Wannigamma GE. A new sesquiterpene lactone from *Elephantopus scaber*. *Phytochemistry* 1982; 21:1173–1175.

10. Rajkapoor B, Jayakar B, Anandan R. Antitumor activity of *Elephantopus scaber* Linn. Against Dalton's Ascitic Lymphoma. *Indian J Pharm Sci* 2002; 64:71–73.
11. Poli A, Nicolau M, Simoes CM, Nicolau RM, Zanin M. Preliminary pharmacologic evaluation of crude whole plant extracts of *Elephantopus scaber*. Part I: in vivo studies. *J Ethnopharmacol* 1992; 37:71–76.
12. Tsai CC, Lin CC. Anti-inflammatory effects of Taiwan folk Medicine 'Teng Khai U' on carrageenan and adjuvant induced paw edema in rats. *J Ethnopharmacol* 1999; 64: 85–89.
13. Lin CC, Tsai CC, Yen MH. The evaluation of hepatoprotective effects of Taiwan folk medicine 'Teng Khai U'. *J Ethnopharmacol* 1995; 45:113–123.
14. Rao GB, Rao YV, Pavan S, Dasari VSP. Qualitative and quantitative phytochemical screening and in vitro anti oxidant and anti microbial activities of *Elephantopus scaber* Linn. *Recent Research in Science and Technology* 2012; 4(4):15–20.
15. Burkhkl H. Dictionary of the Economic Products of the Malay Peninsula. Vol-I. Singapore: Government Printing Press, 1935, 909.
16. Kiritikar KD, Basu BD. *Indian Medicinal Plants*. New Delhi, India: Periodical Experts Books Agency, 1991.
17. Lin JK, Tsai SH, Lin-Shiau SY. Anti inflammatory and anti tumor effects of flavonoids and flavanoids. *Drug Future* 2001; 26: 145–157.
18. Chopra RN, Nayak SL, Chopra IC. *Glossary of Indian Medicinal Plants*. Delhi: CSIR, 1956.
19. Morton J. *Atlas of Medicinal Plants of Middle America, Bahamas to Yucatan*. Springfield, IL: Thomas, 1981, 420.
20. Adhirajan N, Dixit VK, Chandrakasan G. Development and evaluation of herbal formulations for hair growth. *Indian drugs* 1999; 38: 559–563.
21. Roy RK, Thakur M, Dixit VK. Development and evaluation of polyherbal formulation for hair growth promoting activity. *J Cosmet Dermatol* 2007; 6:108–112.
22. Roy RK, Thakur M, Dixit VK. Hair growth promoting activity of *Eclipta alba* in male albino rats. *Arch Dermatol Res* 2008; 300: 357–364.
23. Datta K, Singh AT, Mukherjee A, Bhat B, Ramesh B, Burman AC. *Eclipta alba* extract with potential for hair growth promoting activity. *J Ethnopharmacol* 2009; 124: 450–456.
24. Yoon JI, Al-Reza SM, Kang SC. Hair growth promoting effect of *Zizyphus jujuba* essential oil. *Food Chem Toxicol* 2010; 48: 1350–1354.
25. Adhirajan N, Ravi Kumar T, Shanmugasundaram N, Babu M. In vivo and in vitro evaluation of hair growth potential of *Hibiscus rosa-sinensis* Linn. *J Ethnopharmacol* 2003; 88: 235–239.
26. Roy RK, Thakur M, Dixit VK. Effect of *Citrullus coclocynthis* Schrad. on hair growth activity of albino rats. *Aust J Med Herbal* 2007; 6: 108–112.
27. Liang T, Liao S. Growth suppression of hamster flank organs by topical application of c-linolenic and other fatty acid inhibitors of 5 $\alpha$ -reductase. *J Invest Dermatol* 1997; 109: 152–157.
28. Kobayashi N, Suzuki R, Koide C, Suzuki T, Matsuda H, Kubo M. Effect of leaves of *Ginkgo biloba* on hair regrowth in C3H strain mice. *Yakugaku Zasshi* 1993; 113: 718–724.
29. Takahashi T, Kamiya T, Hasegawa A, Yokoo Y. Procyanidin oligomers selectively and intensively promote proliferation of mouse hair epithelial cells in vitro and activate hair follicle growth in vivo. *J Invest Dermatol* 1999; 112: 310–316.
30. Roh SS, Kim CD, Lee MH, Hwang SL, Rang MJ, Yoon YK. The hair growth promoting effect of *Sophora flavescens* extract and its molecular regulation. *J Dermatol Sci* 2002; 30: 43–49.
31. Awe EO, Makinde JM. The hair growth promoting effect of *Russelia equisetiformis* (Schlect & Chan). *J Nat Prod* 2009; 2: 70–73.
32. Banerjee PS, Sharma M, Nema RK. Preparation, evaluation and hair growth stimulating activity of herbal hair oil. *J Chem Pharm Res* 2009; 1: 261–267.
33. Kwon OS, Han JH, Yoo HG, Chung JH, Cho KH, Eun HC, Kim KH. Human hair growth enhancement in vitro by green tea epigallocatechin-3-gallate (EGCG). *Phytomedicine* 2007; 14: 551–555.

WSTĘPNE BADANIE WPŁYWU WYCIĄGU METANOLOWEGO Z *ELEPHANTOPUS SCABER* LINN. NA PROCES WSPOMAGANIA WZROSTU OWŁOSIENIA U SZCZURÓW

HIMANSHU BHUSAN SAHOO<sup>2</sup>, RAKESH SAGAR<sup>1\*</sup>, SUBRAT KUMAR BHATTAMISRA<sup>3</sup>,  
AMRITA BHAIJI<sup>2</sup>

<sup>1</sup> Department of Pharmacognosy and Phytochemistry  
Dr. H.S. Gour Central University  
Sagar, 470003  
Madhya Pradesh, India

<sup>2</sup> Department of Pharmacology and Toxicology  
Vedica College of Pharmacy  
Bhopal 462021  
Madhya Pradesh, Indie

<sup>3</sup> Department of Pharmacology  
Roland Institute of Pharmaceutical Sciences  
Berhampur 760010  
Orissa, Indie

\*autor, do którego należy kierować korespondencję: e-mail: drrsagar@gmail.com

**Streszczenie**

Badano wpływ wyciągu metanolowego z liści *Elephantopus scaber* na wzrost włosów u szczurów laboratoryjnych. Przez 30 dni badano wzrost włosów u szczurów Wistar po podaniu 2% i 5% preparatu sporządzonego z użyciem ekstraktu metanolowego. Substancją kontrolną był minoksidil (roztwór 2 %). Preparat z ekstraktu metanolowego przygotowany na bazie oleju istotnie ( $p < 0,01$ ) zwiększał inicjację wzrostu włosów i przyspieszał jego zakończenie. Podczas biopsji odnotowano też zależną od dawki poprawę stanu skóry, długości włosów i zwiększenie masy włosów. Podczas badaniu *in vitro* wykazano istotną ( $p < 0,05$ ) poprawę tworzenia się nowych mieszków włosowych. Powyższe wyniki wskazują, że wyciąg metanolowy z *E. scaber* może istotnie zwiększać porost włosów.

**Słowa kluczowe:** łysienie, wzrost włosów, *Elephantopus scaber*, minoksidil