

Ethological study of flower visiting insects in two parks, Kolkata, India

Subhodeep Sarkar¹, Abantika Nandy², Soumendra Nath Talapatra^{2,*}

¹Department of Environmental Science, Asutosh College
S. P. Mukherjee Road, Kolkata, India
*Phone: 91-33-2461-5445

²Department of Environmental Science, University of Calcutta
51/1 & 2 Hazra Road, Kolkata 700019, India

*E-mail address: ecologylive@yahoo.co.in

ABSTRACT

Flower visiting insects attract by the flowers colour, shape, size and fragrance as pollinator. This is a mutual relationship between flowers and insects. The present study aims to know behavioural response or ethology of flower visiting insect in relation to particular flower and/or probable atmospheric changes in two managed parks, Kolkata, India. The study areas were selected as per heavily-populated neighborhoods, nearby office buildings, nearby roads and continuous vehicular movements, human interactions as visitors. The study was carried out at 2 sampling stations viz (i) Elliot park and (ii) Agri-Horticulture Society. The flower species were selected viz. *Helianthus annuus* (sunflower), *Petunia sp.* (petunia) and *Buganvilia spectabilis* (Buganvilia) planting above mentioned areas because these species are more common among other species. In each flower, behavioural response or ethology of visiting insects were studied by visual observation and total 10 flowers of each species were observed randomly. The present results clearly indicate that various insects were majorly showed foraging and feeding behaviour and only mating behaviour was found in two species. This study is a preliminary assessment of flower visiting insects' ethology but further researches are needed in relation to pollination efficiencies of flower visiting insects in the particular flower and air pollution load nearby area by using instruments. It was concluded that insect visitors are showing foraging and feeding behavior but only two species were showed mating behaviour, which may be due to the vehicular air pollution because two parks are located nearby roads and continuous vehicular movements were observed.

Keywords: Insect behavior; Vehicular pollution; Flower visiting insects; Insect ethology

1. INTRODUCTION

Behavioral study or ethology of animal is the study of adaptive behavior in relation to social and environmental circumstances and the natural selection shapes behavior have been

reported by Bird and O'Connell (2006). It deals with the ways in which behavior may be adaptive by allowing an animal to increase or even maximize its reproductive success. The branch of ecology as ethology examines the adaptive significance of behavior, or how behavior may increase survival and reproduction in species. In case of insects, there was majorly observed searching or foraging behavior. Besides this, three different kinds of interaction have been observed between flower and insect: (1) pollinating behavior, (2) feeding behavior, (3) mating behavior. Pollination is one of the most fascinating aspects of interaction between flowering plant (s) and insect (s). Generally the extent of interdependence is regulated by phenology, floral characters as well as by form, structure and behavior of the pollinators. During evolution flowers have developed various strategies for attracting insects (Ram and Mathur, 1984).

Earliest terrestrial insects likely were scavengers may have initially used plants by feeding directly on leaves, roots, spores or pollen. Plant feeding really takes off, when flowering plants develop. There are some categories of feeding interaction between flowering plant and insect. These are (1) leaf chewer (eat entire leaves or large portions of leaves), (2) plant miner and borers (mining involves burrowing into leaves or just below the surface of stems, fruits or roots, boring involves burrowing more deeply into stems, roots, or fruits), (3) sap sucker (pierce plant tissue to obtain liquid from either phloem or xylem), (4) seed predator (some eat the entire seed, but many feed on the inside of the seed by boring into it), (5) nectar or pollen feeder and (6) gall former (galls are an abnormal or pathological growth in plants, done as a response to some invader).

Some insect individuals carried visible pollen loads, females seemed to spend more time on flowers than males, and male spent greater proportion of their time searching for mates (Schlinger, 1956) and the mating takes place within the flower patches. (Borkent and Schlinger, 2008).

Matteson et al. (2012) have documented the presence of herbaceous flowering plants and flower-visiting insects (e.g. Diptera, Coleoptera, Lepidoptera, Hymenoptera) across a wide range of urban green spaces' and residential and commercial blocks. According to Bell (1990), searching behavior is an active movement by which insects seek resources viz. food, mates, oviposition and nesting sites and refugia. It is an important kind of behavior, was already observed in the insects because these resources are absolutely essential for their growth, development, and maintenance in an individual and for ensuring the success of future generations. The searching mechanisms are efficient and accurate assessment mechanisms in which it can be determined the crucial part for an individual's chances of survival and reproduction. It was also documented the searching behavior incurred costs in addition to the energy used for locomotion. Insects visit flowers for several reasons, the predominant reason being for food (Alamu et al., 2013). The mating behavior of insect species is a fundamental issue because it helps to know the conservation of important species or destruction of pest species. The mating behavior is a very complex phenomenon involving several factors (Murvosh et al., 1964). It was also reported that insects of various groups visit flowers in search of mates (Proctor et al., 1996; Sugiura, 2007).

Many studies have already been carried out in discrete urban habitats (e.g. city parks, meadow remnants, community and private gardens) by many research have affirmed the importance of floral resources for floral-feeding insects such as bees and other arthropods (Ram and Mathur, 1984; Cane 2005; Hernandez et al. 2009; Matteson et al., 2012, Alamu et al., 2013). However, few studies have documented across the spectrum of urban habitats, including residential, commercial, and different types of green spaces, to enable an evaluation of the effects of land use heterogeneity on biotic communities within cities (Hennig and

Ghazoul 2011; Sattler et al. 2010; Wojcik 2011; Chowdhury et al., 2014). Chowdhury et al., (2014) have studied the diversity of flower visiting insects in managed park but no one has attempted before the ethological aspects flower visiting insects in the managed park.

The present study aims to know behavioural response or ethology of flower visiting insect in relation to particular flower in two managed parks, Kolkata, India.

2. MATERIALS AND METHODS

The study was carried out at 2 sampling stations viz (i) Elliot park at Jawaharlal Nehru road, (latitude = 22° 32' N and longitude = 88° 20' E) and (ii) Agri-horticulture Society at Alipore Road (latitude = 22° 31' N and longitude = 88° 19' E), situated at 5.18 meters above sea level. The study was carried in dry season.

The flower species were selected viz. *Helianthus annuus*, *Petunia sp.* and *Bougainvillea sp.* growing near roadside of above mentioned areas because these species are more common among other species. These plant species produce flowers at different times during the year and thus, they are good sources of pollen and nectar to the different types of insects that forage on flowers. The direct and indirect influences on flower visiting insects in two urban habitats, ethological assessment was done on specific floral and insect distributions by transects randomly within these two parks and observed the insect pollinators on the basis of foraging, feeding and mating behavior.

The sampling design was involved identifying insects onto the specific flowering species with the help of literatures, books, museum specimens, internet etc. and also by visual identification. Behavioural study was done by visual observation, image capture, etc. All the activities of insects during the visit onto flower were recorded and tabulated in Table 1. These two sampling stations mainly two parks were selected on the basis of human interference, moderate and heavy traffic density and continuous vehicular movement nearby roads as per visualization.

3. RESULTS

In the present study, the behavioral responses or ethology of flower visiting insects were observed in two managed parks, namely Elliot park and Agri-Horticulture Society. In these two parks, the flower visiting insects ethological observation was made on three particular flowering species viz. *Petunia sp.*, *Helianthus annuus* and *Bougainvillea spectabilis*, which were very common in these two parks. The ethology of insects on these flowering species, 10 species of individual flowering species were randomly observed and identified individual insect species and their behavioural response were tabulated in Table 1. In comparison with Agri-horticulture society, various insects viz. small red and black ant, fire ant, dragon fly, lady bug beetle, honeybee, large black ant, bumble bee, hoverfly were observed in Elliot park while in Agri-horticulture society, it was observed some of the above mentioned species alongwith psyche butterfly, leaf beetle, blue/green bottle fly, stick insect and crab spider.

In the present ethological study, it was observed several flower visiting insects and they have showed behavioural responses onto the flower. These insect species viz. the small red and black ants were found in *Petunia sp.* *Bougainvillea sp.* in Elliot park, and was absent in sunflower and was mainly showed nectar searching behavior and crawling behavior on the

petal. But in Agri-horticulture society, ant was present only in sunflower and absent in other flowers and showed the same activity like Elliot Park. Dragonfly was very less in number in Elliot Park than Agri-horticulture society.

Table 1. Flower visiting insect behavior or ethological study between two parks.

SI No.	Flower species	ELLIOT PARK			AGRI-HORTICULTURE SOCIETY		
		Insect species	Behavioral response		Insect species	Behavioral response	
			Foraging and Feeding behavior	Mating behavior		Foraging and Feeding behavior	Mating behavior
1.	<i>Petunia sp.</i>	Small black ant <i>Monomorium sp.</i>	A small ant was at the base of the flower and crawled on the petal	No behavior	Common Evening Brown Butterfly, <i>Melanitis sp.</i>	This butterfly sat on a leaf for some time and wagged it's wing in a linear direction.	No behavior
	Dragonfly <i>Agriocnemis sp.</i>	A dragonfly sat on a petal for long time and sucked petal juice. It changed the petal after some time period and it also nodded its tail after certain time interval.	No behavior.				
2.	<i>Petunia sp.</i>	No insect found	No behavior	No behavior	No insect found	No behavior	No behavior
3.	<i>Petunia sp.</i>	Lady bug Beetle <i>Coccinella transversalis</i>	It was sitting on the petal for a long time.	No behavior	Dragonfly, <i>Ceragrion coromandelianum</i>	The dragonfly sits on leaf and petal for very short time and changes the leaf very quickly.	No behavior

4.	<i>Petunia sp.</i>	No insect found	No behavior	No behavior	Leaf beetle, <i>Phyllotreta sp.</i>	It was sitting for a long time on the leaf and at seems to be collecting food from the leaf.	No behavior
5.	<i>Petunia sp.</i>	No insect found	No behavior	No behavior	No insect found	No behavior	No behavior
6.	<i>Petunia sp.</i>	Honey bee <i>Apis sp.</i>	It was collecting nectar from the base region of the flower and was flying around the flower.	No behavior	Hoverfly, <i>Melanostoma fasciatum</i>	The fly was sitting for a long time on the white petal. It seems to be sucking petal juice or making some activity by its mouth parts.	No behavior
7.	<i>Petunia sp.</i>	No insect found	No behavior	No behavior	No insect found	No behavior	No behavior
8.	<i>Petunia sp.</i>	No insect found	No behavior	No behavior	No insect found	No behavior	No behavior

12.	<i>Helianthus annuus</i>	honey bees. <i>Apis sp.</i>	Same activity.	No behavior	Honey bee. <i>Apis sp.</i>	Collecting & searching nectar from the disc floret region	No behavior
11.	<i>Helianthus annuus</i>	honey bees. <i>Apis sp.</i>	flying on a circular pattern over the flower. crawling and searching & collecting of nectar at disc floret region.	No behavior	Hoverfly , <i>Melanostoma fasciatum</i>	A fly was sitting on leaf and making some activity with its mouthparts.	No behavior
					Lady bug beetle <i>Coccinella transversalis</i>	It was sucking juice from young green stem.	No behavior
10.	<i>Petunia sp.</i>	Hoverfly , <i>Melanostoma fasciatum</i>	A fly was sitting on the petal for a long time. It seems to be sucking juice from the petal.	No behavior	No insect found	No behavior	No behavior
9.	<i>Petunia sp.</i>	Fire ant <i>Solenopsis sp.</i>	It was crawling on the leaf and petal and seems to be that ant want to collect nectar.	No behavior	No insect found	No behavior	No behavior

13.	<i>Helianthus annuus</i>	honey bees. <i>Apis sp.</i>	Same activity	No behavior	No insect found.	No behavior.	No behavior
14.	<i>Helianthus annuus</i>	honey bees. <i>Apis sp.</i>	Same activity	No behavior	Dragon fly <i>Ceriagrion coromandelianum</i>	It was sitting on the green stem and sucking juice from it.	No behavior
15.	<i>Helianthus annuus</i>	honey bees. <i>Apis sp.</i>	Same activity	No behavior	Honey bee <i>Apis sp.</i>	Collecting & searching nectar from the disc floret region	No behavior
16.	<i>Helianthus annuus</i>	honey bees. <i>Apis sp.</i>	Same activity	No behavior	Dragon fly <i>Ceriagrion coromandelianum</i>	Dragonfly sitting on the leaf and it seems to be sucking juice from the leaf. It touches it's tip of the tail by its leg by bending the tail to the leg. It also moves its head and eyes.	No behavior
17.	<i>Helianthus annuus</i>	No insect found	No behavior	No behavior	No insect found.	No behavior.	No behavior

21.	<u>Bougainvillea spectabilis</u>	Lady Bug <u>Coccinella transversalis</u>	Continuously moving it's sensory organ and also crawling on the leaf and flower.	No behavior	No insect found	No behavior.	No behavior
20.	<i>Helianthus annuus</i>	bumble bee <i>Bombus sp.</i>	Sucking of nectar by mouthparts & move very fast from one flower to another.	No behavior	Lady bug beetle <u>Coccinella transversalis</u>	Sitting on the green stem for a long time and it seems to be sucking juice from stem.	No behavior
					Large black ant, <i>Camponotus sp.</i>	Searching food on leaf and green stem.	No behavior
19.	<i>Helianthus annuus</i>	honey bees. <i>Apis sp.</i>	Same activity	No behavior	Lady bug beetle <u>Coccinella transversalis</u>	sitting on the upper surface of the leaf at sunlight and sucking juice from stem.	No behavior
18.	<i>Helianthus annuus</i>	honey bees. <i>Apis sp.</i>	Same activity	No behavior	Large black ant, <i>Camponotus sp.</i>	Crawling on the leaf and green stem likely searching food.	No behavior

22.	<u>Bougainvillea spectabilis</u>	Lady Bug <u>Coccinella transversalis</u>	Continuously moving it's sensory organ and also crawling on the leaf and flower.	No behavior	bumble bee <i>Bombus sp.</i>	Simply pollinating behavior.	No behavior
23.	<u>Bougainvillea spectabilis</u>	No insect found	No behavior	No behavior	Blue green bottle fly <i>Calliphora sp.</i>	No behavior	Mating behavior on the upper surface of the leaf
24.	<u>Bougainvillea spectabilis</u>	No insect found	No behavior.	No behavior	green marsh hawk dragonfly, <i>Orthetrum sabina</i>	Sitting on the leaf & moves its head.	No behavior
25.	<u>Bougainvillea spectabilis</u>	Large black ant, <i>Camponotus sp.</i>	Searching for food on the petal and was crawling on the flower petal and base.	No behavior	psyche butterfly , <i>Leptosia sp.</i>	Simply pollinating behavior.	No behavior
26.	<u>Bougainvillea spectabilis</u>	Hoverfly <i>Melanostoma fasciatum</i>	Sitting on the flower petal for a long time and it seems to be sucking juice from the petal.	No behavior	Common Evening Brown Butterfly, <i>Melanitis sp.</i>	Simply pollinating behavior.	No behavior

27.	<u>Bougainvillea spectabilis</u>	No insect found.	No behavior.	No behavior	Dragon fly <i>Ceriatrigon coromandelianum</i>	sitting on the leaf & sucking juice from the leaf.	No behavior
28.	<u>Bougainvillea spectabilis</u>	Hoverfly <i>Melanostoma fasciatum</i>	Sitting on the flower and sucking nectar from the stigma by using it's mouth parts.	No behavior	Stick insect <i>Phasmida sp.</i>	Sucking petal juice from its mouthparts.	No behavior
29.	<u>Bougainvillea spectabilis</u>	Bumble bee <i>Bombus sp.</i>	Moving from flower to flower very fast. It was clearly pollinating behavior.	No behavior	Crab Spider <i>Misumenoides sp.</i>	It may be sucking juice from the petal.	No behavior
30.	<u>Bougainvillea spectabilis</u>	Hoverfly <i>Melanostoma fasciatum</i>	Sitting on flower for a short time period and was moving it's wing in a particular pattern.	No behavior	No insect found	No behavior	No behavior

This species was only found in *Petunia sp.* in Elliot Park while it was present in all three flowering species in Agri-horticulture society. It was mainly showed feeding behavior by sucking juice from flower petal. Lady bug beetle was showed mainly feeding behavior and searching for food but a particular and continuous movement of their sensory organ was observed in Elliot park which was absent in Agri-horticulture society. Honeybees were showed a preference in sunflower as their source of food in both of the park, but there was absent in other flower of Agri-horticulture society while honey bees were showed feeding and searching behavior in petunia in Elliot park. Bumble bee was present in both the parks

and was showed feeding behavior in the sunflower of Elliot park and it was also showed pollinating behavior clearly in Bougainvillea flower in both parks. Hoverfly was also observed resting for a very long time on the flower and it was seemed to be sucking petal juice. They were present in Elliot Park, while in Agri-horticulture society, only housefly was found but there was no hoverfly. Butterfly was absent in Elliot Park, in comparison with Agri-horticulture society. It was showed clearly pollinating behavior on bougainvillea flower and was also moved its wings in a particular linear direction while sitting on *Petunia sp.* A crab spider was showed simply feeding and foraging behavior on bougainvillea at Agri-horticulture society. Mating behavior was observed very rare in two managed parks. No insect was showed any kind of mating behavior in Elliot Park, while in Agri-horticulture society two blue/green bottle flies were showed clearly mating phase on the upper surface of the bract of Bougainvillea flower.

4. DISCUSSION AND CONCLUSION

The present study of behavioral responses or ethology of flower visiting insects in two managed park namely Elliot Park and Agri-Horticulture Society indicate that there were variable changes among the behavioral responses of same species in two different parks and also the species diversity was more or less different in two parks. The diversity variation has already been studied by Chowdhury et al., (2014).

Generally the green spaces' (e.g. cemeteries, parks, gardens) study have showed an important research area within cities to determine urban ecology. However, study of biotic communities besides office and residential buildings is important because the area, where humans spend most of their time, they interact with nature and where most ecosystem services are likely interacting on a daily basis (Pickett and Cadenasso 2008). The difference shown in two different parks may be vary on certain factors such as human interference, as Elliot park is a public park, vehicular air pollution as Elliot park is just beside the road side and landscape change. Insect outbreaks have been observed in the surroundings of polluted industrial areas and along highways (Holopainen, 2002). According to Grimm et al., (2000), human actions dramatically altered the functioning of ecosystems of which humans are a part, and equally, humans are a part of virtually all ecosystems and have been so for millennia. Nowhere has this human participation been more intense than in cities, suburbs, and exurbs and in the supporting hinterlands. Researchers at the University of Virginia in Charlottesville report that three common constituents of smog destroy floral scents released by flowers to attract bees and other pollinators (Potera, 2008), which may affect the insect behavioral responses. According to McFrederick et al., (2009), chemical signals play important roles in ecological interactions but are vulnerable to perturbation by air pollution. In polluted air masses, signals may travel shorter distances before being destroyed by chemical reactions with pollutants. Pollination, attraction of natural enemies of plant pests, aggregation pheromones, and mate attraction are likely to be affected (McFrederick et al., 2009).

It was observed in the present results (Table 1) that when comparison was made between Elliot park and Agri-horticulture society, various insects viz. small red and black ant, fire ant, dragon fly, lady bug beetle, honey bee, large black ant, bumble bee, hoverfly were observed in Elliot park while in Agri-horticulture society, it was observed some of the above mentioned species alongwith psyche butterfly, leaf beetle, blue/green bottle fly, stick insect and crab spider.

In this present study it was observed that insects were preferred particular flower for their bright colour, shape, nectar, fragrance, food sources etc. All the honey bees were observed in sunflower and petunia flower and were showed feeding and searching behavior. This was an evidence of other research works (Winston, 1991; Celebrezze and Paton, 2004; Ushimaru et al., 2007; Whelan et al., 2009) that honey bees are good pollinators.

The small red and black ants were mainly showed nectar searching behavior and crawling behavior on the petal. Dragonflies were less in number and were mainly showed feeding behavior by sucking juice from flower petal. Lady bug beetle was showed mainly feeding behavior and searching for food but a particular and continuous movement of their sensory organ. Hoverfly was also observed resting for a very long time on the flower and it was seemed to be sucking petal juice. A crab spider was showed simply feeding and foraging behavior. The stick insect was showed sucking of petal juice as feeding behavior. Similar observations were established of flower visiting insects on the basis of foraging and feeding behavior (Wehner et al., 1983; Bell et al., 1985; Bell, 1990; Alamu et al., 2013).

Butterflies were showed clearly pollinating behavior and also moved its wings in a particular linear direction while sitting on flowering species, the behavior may be an attraction to the partner as mating behavior (Singer, 1982). No insect was showed any kind of mating behavior in Elliot Park, but only in Agri-horticulture society two blue/green bottle flies were showed clearly mating phase on the upper surface of the bract of Bougainvillea flower. The mating behavior is depending upon many environmental factors (Mondor et al., 2004). Pheromones are utilized by insects for several purposes, including alarm signaling (Kislow and Edwards, 1972; Blatt et al., 1998; Hunt et al., 2003), kin recognition (Dani et al., 2001; Vander Meer and Alonso, 2002) and sexual communication (Pickett et al., 1992; Landolt and Phillips, 1997). It was also reported that insects of various groups visit flowers in search of mates (Proctor et al., 1996; Sugiura, 2007) but the present results have documented mating behavior only in bottle flies among all the present insects, which may be due to the affect of atmospheric changes by automobile air pollution (McFrederick et al., 2009) and there may be the chance of vehicular air pollution affect (CPCB, 2009; Citizen's Report, 2011) because these two parks are located nearby roads. In other way, the behavioral differences among the insects of two managed park may be on the size, colour or fragrance of these studied flower.

In the present study it was concluded that the flower visiting insects were observed majorly foraging and feeding behavior while mating behavior was found only two insect species among all other species, which may be the effects of individual and/or combination of automobile air pollutants (McFrederick et al., 2009). It was already reported that Indian cities are susceptible to automobile air pollution (Trivedi et al., 2003; CPCB, 2009; Citizen's report, 2011), though there no attempt has been made on physico-chemical properties of present air pollutants. This study is a preliminary assessment of flower visiting insects and their eco-ethological study. Many studies on flower visiting insects have already been done in greenspace, garden etc. in other parts of globe (Bell, 1990; Matteson et al., 2012, Chowdhury et al., 2014) but less work have documented in eco-ethological aspects in managed parks (Borkent and Schlinger, 2008). In addition, further researches are needed in relation to pollination efficiencies of flower visiting insects in the particular flower and air pollution load by using instruments. It was observed that three flowering species viz. sunflower, petunia and bougainvillea are very common both the parks but insect visitors are showing foraging and feeding behavior but only two species were showed mating behaviour, which may be due to the vehicular air pollution because two parks are located nearby roads and continuous vehicular movements were observed.

Acknowledgement

The authors convey their gratitude to the Department of Environmental Science, University of Calcutta, for providing the necessary infrastructure for doing this study.

References

- [1] Alamu O.T., Amao A.O., Oke O.A., Suleiman R.A., *Int J Adv Agri Res* 1 (2013) 87-91.
- [2] Bell W.J., *Ann Rev Entomol* 35 (1990) 447-67.
- [3] Bell W.J., Tortorici C., Roggero R.J., Kipp, L.R., Tobin, T.R., *Anim Behav* 33 (1985) 436-448.
- [4] Bird D.W., O'Connell J.F., *J Archaeol Res* 14(2) (2006) 143-188.
- [5] Blatt S.E., Borden J.H., Pierce H.D.Jr., Gries R., Gries G., *J Chem Ecol* 24 (1998) 1013-1031.
- [6] Borkent C.J., Schlinger E.I., *The Canadian Entomologist* 140(2) (2008) 250-256.
- [7] Dani F.R., Jones G.R., Destri S. et al., *Anim Behav* 62 (2001) 165-171.
- [8] Cane, J.H., Bees' needs challenged by urbanization. – In: Johnson, E. A. and Klemens, M. W. (eds), *Nature in fragments: the legacy of sprawl*. Columbia Univ. Press, (2005), pp. 109-124.
- [9] Celebrezze T., Paton D.C., *Austral Ecol* 29 (2004) 129-136.
- [10] Chowdhury S., Sarkar S., Nandy A., Talapatra S.N., *International Letters of Natural Sciences* 10 (2014) 58-68.
- [11] Citizen's Report, *Centre of Science and Environment*, (2011) 1-106.
- [12] CPCB, Central Pollution Control Board, New Delhi, (2009).
<http://www.cpcb.nic.in/bulletin/del/2009.html>.
- [13] Grimm N.B., Grove J.M., Pickett S.T.A., Redman C.L., *BioScience* 50 (2000) 571-584.
- [14] Hennig E.I., Ghazoul J., *Perspectives in Plant Ecology, Evolution and Systematics* 13 (2011) 137-150.
- [15] Hernandez J.L., Frankie G.W., Thorp R.W., *Cities and the Environment* 2(1) (2009) 15.
<http://escholarship.bc.edu/cate/vol2/iss1/3>
- [16] Holopainen J.K., *Entomologia Experimentalis et Applicata* 104 (2002) 137-142.
- [17] Hunt G.J., Wood K.V., Guzman-Novoa E., Lee H.D., Rothwell A.P., Bonham, C.C., *J Chem Ecol* 29(2) (2003) 453-463.
- [18] Kislow C.J., Edwards L.J., *Nature* 235 (1972) 108-109.
- [19] Landolt P.J., Phillips T.W., *Ann Rev Entomol* 42 (1997) 371-391.
- [20] Matteson K.C., Grace J.B., Minor, E.S., *Oikos* (2012) 01-13.
- [21] McFrederick Q.S., Fuentes J.D., Roulston T'ai., Kathilankal J.C., Lerdau M., *Oecologia* 160 (2009) 411-420.

- [22] Mondor E., Tremblay M., Awmack C., Lindroth R.L., *Global Change Biology* 10 (2004) 1820-1824.
- [23] Murvosh C.M., Fye R.L., Labrecque G.C., *Ohio J Science* 64(4) (1964) 264-271.
- [24] Potera C., *Environ Health Perspective* 116(8) (2008) A334.
- [25] Proctor M., Yeo P., Lack A., The natural history of pollination. Timber Press, Portland, OR (1996).
- [26] Pickett J.A., Wadhams L.J., Woodcock C.M., Hardie J., *Ann Rev Entomol* 37 (1992) 67-90.
- [27] Pickett S.T.A., Cadenasso M.L., *J Ecol* 96 (2008) 8-12.
- [28] Ram H., Mathur G., *Proc Indian Acad Sci (Anim Sci)* 93(4) (1984) 359-363.
- [29] Sattler S.E., Funnell-Harris D.L., Pedersen J.F., *Food Chem* 58 (2010) 3611-3616.
- [30] Schlinger E.I., *Proceedings of the United States National Museum* 106 (1956) 359-375.
- [31] Singer M.C., *Oecologia* 52 (1982) 230-235.
- [32] Sugiura S., Abe T., Yamaura Y., Makino S., *Naturwissenschaften* 94 (2007) 703-707.
- [33] Trivedi S., Agrawal M., Rajput M., *Indian J Air Poll Cont.* 3 (2003) 44-53.
- [34] Ushimaru A., Watanabe T., Nakata K., *American J Bot* 94(2) (2007) 249-258.
- [35] Vander Meer R.K., Alonso L.E., *Behav Ecol Sociobiol* 51 (2002) 122-130.
- [36] Wehner R., Harkness R.D., Schmid-Hempel P., Foraging strategies in individually searching ants *Cataglyphis bicolor* (Hymenoptera: Formicidae). In Information Processing in Animals, ed. M. Lindauer. Vol. I. New York: Gustav Fischer Verlag (1983).
- [37] Whelan R.J., Ayre D.J., Beynon F.M., *Annals Bot* 103 (2009) 1395-1401.
- [38] Winston M.L., The Biology of the Honey Bee, Harvard University Press, (1991) pp. 294, ISBN 970674074095.
- [39] Wojcik V.A., *J Pollin Ecol* 4(7) (2011) 48-56.

(Received 28 May 2014; accepted 04 June 2014)