

Article ID: 193117
DOI: 10.5586/aa/193117

Publication History
Received: 2024-04-24
Accepted: 2024-09-09
Published: 2024-10-22

Handling Editor
Anna Bieniek; University of Warmia and Mazury in Olsztyn, Poland; <https://orcid.org/0000-0002-5903-1405>

Authors' Contributions
RMH-H: Research concept and design; AAE-A: Collection and/or assembly of data; AAE-A: Data analysis and interpretation; RMH-H, AAE-A: Writing the article; RMH-H: Critical revision of the article; RMH-H: Final approval of the article



Funding
This research received no external funding.

Competing Interests
No competing interests have been declared.

Copyright Notice
© The Author(s) 2024. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits redistribution, commercial and noncommercial, provided that the article is properly cited.

SHORT COMMUNICATION

Effects of green seaweed (*Ulva onhoi*) on the reproductive development of tomato (*Solanum lycopersicum*) plants

Adrian Alejandro Espinosa-Antón ,
Rosalba Mireya Hernández-Herrera *

Departamento de Botánica y Zoología, Centro Universitario de Ciencias Biológicas y Agropecuarias (CUCBA), Universidad de Guadalajara, Zapopan, 45200, México

* To whom correspondence should be addressed. Email: rosalmir@yahoo.com

Abstract

Algae-derived products have great potential as crop biostimulants due to their multiple beneficial effects at different stages of plant development. Green seaweeds of the genus *Ulva* are well suited for this purpose because they are widely distributed and grow rapidly in a wide range of conditions. In this study, a greenhouse experiment was conducted to evaluate the effects of dry seaweed powder (DSP) and liquid seaweed extract (LSE) of *Ulva ohnoi* on the reproductive development of tomato plants. The experiment included three treatments: (1) plants treated with 5 g of DSP, (2) plants treated with 250 mL of LSE, and (3) control plants (without algae). The reproductive parameters, chlorophyll content, and mineral composition were measured during the flowering and early fruiting periods. The application of DSP was the most effective treatment in promoting early flowering and significantly increased the number of buds (103%), flower clusters (55%), flowers (61%), and fruits (45%) per plant. Furthermore, the DSP-treated plants exhibited an enhancement in the levels of chlorophyll and nutrients in the plants and fruits. The results of the current work show that the application of *U. ohnoi* in its natural form (dry powder) stimulates the reproductive development of crop tomato. This represents a sustainable and natural alternative to synthetic inputs that growers can incorporate into horticultural production to improve yield attributes.

Keywords

plant biostimulant; tomato productivity; nutrient status; reproductive organs

1. Introduction

Tomato (*Solanum lycopersicum* L.) is one of the most popular horticultural crops in the world in terms of human consumption and other industrial uses (O. Ali et al., 2019). Furthermore, tomato production represents an important source of income for rural and peri-urban producers in several countries worldwide (Suchithra et al., 2022). In 2022, Mexico was the eighth largest producer of tomatoes in the world, producing nearly 3.46 million tons (Government of Mexico, 2024; <https://www.gob.mx>). As with other crops, the productive fitness of tomato is largely dependent on the plant's ability to form and mature flowers and fruits (Hussain et al., 2021). Therefore, enhancing reproductive development through the use of natural products is a sustainable strategy to increase crop yield and quality and minimize dependence on synthetic chemical inputs that impact the environment (Ma et al., 2022).

Seaweeds represent a diverse and abundant marine resource along the world's coastlines, with great potential as a source of biofertilizers and biostimulants for modern horticulture (Di Stasio et al., 2017). Several studies have demonstrated the beneficial effects of seaweed-derived products on the vegetative growth and yield parameters of tomato plants (O. Ali et al., 2019; Hussain et al., 2021). However, there is limited information on the phenotypic response of plants treated with these products to

reproductive development traits, and most of these studies focus on the evaluation of the biostimulant properties of brown seaweed-derived formulations. In particular, seaweed extracts have been demonstrated to stimulate early flowering, increased flower formation, and improved fruit set in a variety of crop plants, including bean, pepper, and tomato (N. Ali et al., 2016; O. Ali et al., 2019; Dookie et al., 2020; El-Yazied et al., 2012). These positive effects during the reproductive development phase inevitably led to an improvement in crop yields. More recently, Hussain et al. (2021) observed an increase in the number of flowers and fruits in response to the application of a commercial brown seaweed extract, and Dookie et al. (2020) reported increased expression levels of flowering genes, following the application of alkaline extracts derived from *Ascophyllum nodosum* or *Sargassum* sp.

In turn, few studies have evaluated the agricultural potential of green algal species (Ma et al., 2022). In a previous study, we reported that treatment with the green alga *Ulva ohnoi* (i.e., dry seaweed powder and liquid seaweed extract) improved the vegetative growth attributes of tomato plants (Espinosa-Antón et al., 2023). Therefore, we hypothesize that the application of *U. ohnoi* will accelerate the initiation of flowering and enhance flower and fruit formation by improving the nutritional status and growth characteristics during the vegetative phase. The objective of this study was to evaluate the effects of drench application of the dry seaweed powder and a liquid seaweed extract of *U. ohnoi* on the reproductive development of tomato.

2. Material and methods

2.1. Obtention of dry seaweed powder and preparation of seaweed liquid extracts

Ulva ohnoi biomass was obtained from the company Productos Marinos de las Californias S. DE R.L. DE C.V., Ensenada, Mexico. A liquid extract at 0.2% was obtained as in Espinosa-Antón et al. (2023). The application of 5 g of the dry seaweed powder every 15 days and 250 mL of the liquid seaweed extract at 0.2% every 7 days was previously established by Espinosa-Antón et al. (2023).

2.2. Greenhouse experiment

The experiments were conducted using certified *Solanum lycopersicum* L. cv. “Rio Fuego” seeds (Kristen Seed, San Diego, CA, USA).

The tomato plants were transplanted into individual 5 L pots with vermiculite, peat moss, and soil in a ratio of 1:1:1 (v/v) and grown in greenhouse conditions with 16 h light regime at 30 °C and 8 h dark regime at 16 °C, and relative humidity 70 to 85% for 75 days after transplanting (DAT). The plants were watered daily with 250 mL of water and fertilized every two weeks with 150 mL of N–P–K solution (20:20:20) until the end of the experiment. The plants were arranged in a completely randomized design, with each treatment row (six plants) replicated two times ($n = 12$). Treatments: (1) plants treated with 5 g of dry seaweed powder (DSP) every 15 days; (2) plants treated with 250 mL of liquid seaweed extract at 0.2% (LSE) every 7 days; and (3) control plants (without seaweed). Both DSP and LSE treatments were applied directly to the substrate surface with the drench method one week after transplanting and every 7 or 15 days thereafter until the end of the experiment.

2.3. Data collection from the reproductive development stage of the tomato crop

Data were collected during the reproductive development stage of the tomato crop (30 to 70; DAT), corresponding to the flowering and early fruiting periods. All the plants were monitored each day to determine when the first flowers emerged and opened. The reproductive parameters evaluated on the treated plants included the number of flower clusters, buds, flowers, and fruits (Figure 1).




Figure 1 Schematic representation of parameters evaluated in the tomato crop. (A) chlorophyll content in leaves; (B) flower bud clusters; (C) flowers; (D) development of fruits; (E) ripe fruit for chemical analysis.

Table 1 Effect of dry seaweed powder (DSP) and liquid seaweed extract (LSE) of *U. ohnoi* on the reproductive development of tomato plants.

Treatments	Flower clusters/plant	Flower buds/plant	Flowers/plant	Fruits/plant
Control	5.50 ± 1.62 ^a	5.58 ± 2.8 ^a	10.88 ± 3.6 ^a	21.33 ± 6.8 ^a
DSP	8.50 ± 2.71 ^b	11.33 ± 3.7 ^b	17.08 ± 4.5 ^b	31.00 ± 8.7 ^b
LSE	5.08 ± 1.62 ^a	5.75 ± 3.2 ^a	10.75 ± 3.6 ^a	25.67 ± 8.8 ^a

Values represent the mean ± standard deviation ($n = 12$ plants). Different letters in columns (a–c) indicate significant differences compared to the control, Dunnett's test ($p < 0.05$).

2.4. Chemical analysis of tomato plants

Chlorophyll content in leaves at 30, 50, and 70 DAT was measured with a non-intrusive method using a SPAD 502 Plus portable chlorophyll meter (Minolta; Spectrum Technologies Inc., Aurora, IL, USA) and expressed as equivalent to SPAD units. On the other hand, at 50 DAT, leaf samples were collected from the same 12 plants from each treatment to determine the nutrient composition. Nitrogen, potassium, calcium, magnesium, sulfur, and iron were then determined using a 240FS AA atomic absorption spectrometer (PerkinElmer, Santa Clara, CA, USA), while phosphorus was determined by colorimetry (Espinosa-Antón et al., 2023). The parameters of tomato fruits, such as total soluble solids and pH, were measured using a portable refractometer and a HI 2211 pH meter (Hanna Instruments). All mineral measurements were performed with three replicates.

Data were analyzed using one-way ANOVA and Dunnett's multiple comparison of means ($p < 0.05$).

3. Results

3.1. Reproductive development stage of the tomato crop

In this study, at 30 DAT, flower bud formation was observed in the three experimental groups, indicating the beginning of the reproductive stage. However, the first opened flowers were observed at 35 DAT in the DSP-treated plants, indicating a reduction of five days in the start of the flowering process compared to the control plants. Similarly, the treatment with DSP resulted in an increase in the number of flower clusters (55%), buds (103%), and flowers (61%) at 50 DAT and the number of fruits per plant (45%) at 70 DAT compared to the control plants (Table 1). In contrast, the LSE-treated plants did not show significant differences in the reproductive parameters compared to the control plants.

Table 2 Minerals content in foliar tissue of tomato plants treated with dry seaweed powder (DSP) or liquid seaweed extract (LSE) of *Ulva ohnoi*.

Nutrients (%)	Control	DSP	LSE
Nitrogen	0.92 ± 0.08 ^a	1.33 ± 0.15 ^b	0.98 ± 0.02 ^a
Potassium	1.21 ± 0.05 ^a	1.44 ± 0.04 ^b	1.24 ± 0.07 ^a
Phosphorous	3.53 ± 0.03 ^a	4.23 ± 0.03 ^b	4.41 ± 0.01 ^c
Calcium	1.44 ± 0.01 ^a	1.56 ± 0.02 ^b	1.61 ± 0.02 ^c
Magnesium	0.51 ± 0.01 ^a	0.67 ± 0.02 ^b	0.53 ± 0.01 ^a
Sulfur	1.11 ± 0.07 ^a	1.38 ± 0.01 ^b	1.29 ± 0.13 ^a
Iron	0.0012 ± 0.8 ^a	0.0014 ± 0.9 ^b	0.0015 ± 0.8 ^c

Values represent mean ± standard deviation from a mix of foliar tissue of four plants ($n = 3$). Different letters in rows (a–c) indicate significant differences compared to the control, Dunnett's test ($p < 0.05$).

Table 3 Dry matter (DM), total soluble solid (° Brix) and pH in tomato fruits treated with dry seaweed powder (DSP) or liquid seaweed extract (LSE) of *Ulva ohnoi*.

	DM	° Brix	pH
Control	5.88 ± 0.54 ^a	3.98 ± 0.67 ^a	4.26 ± 0.78 ^b
DSP	5.94 ± 0.78 ^a	4.08 ± 0.68 ^b	4.33 ± 0.99 ^c
LSE	6.07 ± 0.74 ^a	4.15 ± 0.48 ^b	4.19 ± 0.81 ^a

3.2. Chemical analysis of tomato plants

The results obtained from the chemical analysis of plants from the DSP treatment showed a significant ($p < 0.05$) increase in the foliar content of N (45%), K (19%), P (20%), Ca (8%), Mg (31%), S (24%), and Fe (17%), while the SLE treatment increased the content of P (25%) and Fe (25%) compared to the control plants (Table 2). In the same way, the DSP treatment enhanced the dry matter, total soluble solid, and pH in the tomato fruits (Table 3).

Likewise, the DSP treatment significantly increased the chlorophyll content at 30 DAT (3.5 units), 50 DAT (8.1 units), and 70 DAT (6.2 units), while the SLE treatment had a similar effect at 70 DAT (6.2 units) compared to the control plants (Figure 2).

4. Discussion

Our study shows that the treatment with the green algae *Ulva ohnoi* in its natural form (dry powder) improved the reproductive development parameters of the tomato plants. Similarly, greenhouse tomato plants exhibited an increase in the number of flowers and fruits after soil application of the dry biomass of the green algae *Acutodesmus dimorphus* and *Chlorella vulgaris* and the brown algae *Sargassum johnstonii* (García-González & Sommerfeld, 2016; Suchithra et al., 2022). More broadly, other studies have also shown that application of seaweed extracts induces early flowering and increased fruit set in tomato and pepper plants (N. Ali et al., 2016; O. Ali et al., 2019; Hussain et al., 2021; Suchithra et al., 2022).

The observed results may be partially related to the biostimulant and biofertilizer properties of the algal products/extracts. This is due to the presence of bioactive compounds in algae, including various types of polysaccharides, amino acids, phenolic compounds, and plant growth regulators, which stimulate the absorption and assimilation of nutrients essential for crop growth and development (Di Stasio et al., 2017). In particular, the reproductive phase represents a critical developmental transition point during which plants reallocate their metabolism and require specific amounts of certain nutrients (Hussain et al., 2021). For the flowering stage, the tomato requires large amounts of P (responsible for the number of flowers and buds that form) and K (promotes flower initiation), while during the fruiting stage, the most needed elements are K (stimulates flower maturation and fruit formation) and Ca (promotes adequate




Figure 2 Relative chlorophyll content in the leaves of tomato plants treated with dry seaweed powder (DSP) and liquid seaweed extract (LSE) of *Ulva ohnoi*. The vertical bars represent mean \pm standard error ($n = 12$ plants). The asterisk (*) in the bars indicates significant differences in comparison to the control group based on Dunnett's test ($p < 0.05$).

fruit growth) (Bodale et al., 2021). The aforementioned information is in accordance with the results of our study on the nutritional status of *U. ohnoi*-treated plants during the reproductive phase, as expressed through foliar mineral and chlorophyll content (Table 2, Figure 2).

Another possible explanation for our findings is related to our previous research, which demonstrated the promotion of microbial growth in the soil of tomato plants following the application of DSP and SLE of *U. ohnoi* (Espinosa-Antón et al., 2023). In this regard, Hussain et al. (2021) suggested that the observed increase in the number of flower clusters, flowers, and fruits in tomato plants grown in soil treated with seaweed extracts was due to an increase in beneficial bacterial families in the rhizosphere at flower initiation. On the other hand, it is possible that the bioactive compounds present in *U. ohnoi* induced the overexpression of key genes in reproductive morphogenesis (Dookie et al., 2020).

This study provides the first evidence of the biostimulant effect of *U. ohnoi* powder on tomato reproductive development, justifying its use to improve yield and quality in greenhouse horticulture crops.

References

- Ali, N., Farrell, A., Ramsubhag, A., & Jayaraman, J. (2016). The effect of *Ascophyllum nodosum* extract on the growth, yield and fruit quality of tomato grown under tropical conditions. *Journal of Applied Phycology*, 28, 1353–1362. <https://doi.org/10.1007/s10811-015-0608-3>
- Ali, O., Ramsubhag, A., & Jayaraman, J. (2019). Biostimulatory activities of *Ascophyllum nodosum* extract in tomato and sweet pepper crops in a tropical environment. *Plos One*, 14, Article e0216710. <https://doi.org/10.1371/journal.pone.0216710>
- Bodale, I., Mihalache, G., Achiței, V., Teliban, G. C., Cazacu, A., & Stoleru, V. (2021). Evaluation of the nutrients uptake by tomato plants in different phenological stages using an electrical conductivity technique. *Agriculture*, 11(4), Article 292. <https://doi.org/10.3390/agriculture11040292>
- Di Stasio, E., Roupheal, Y., Colla, G., Raimondi, G., Giordano, M., Pannico, A., El-Nakhel, C., & de Pascale, S. (2017). The influence of *Ecklonia maxima* seaweed extract on growth, photosynthetic activity and mineral composition of *Brassica rapa* L. subsp. *sylvestris* under nutrient stress conditions. *European Journal of Horticulture Science*, 82(6), 286–293. <https://doi.org/10.17660/eJHS.2017/82.6.3>
- Dookie, M., Ali, O., Ramsubhag, A., & Jayaraman, J. (2020). Flowering gene regulation in tomato plants treated with brown seaweed extracts. *Scientia Horticulture*, 276, Article 109715. <https://doi.org/10.1016/j.scienta.2020.109715>

- El-Yazied, A., El-Gizawy, A. M., Ragab, M. I., & Hamed, E. S. (2012). Effect of seaweed extract and compost treatments on growth, yield and quality of snap bean. *Journal of American Science*, 8(6), 1–20.
- Espinosa-Antón, A. A., Zamora-Natera, J. F., Zarazúa-Villaseñor, P., Santacruz-Ruvalcaba, F., Sánchez-Hernández, C. V., Alcántara, E. A., Torres-Morán, M. I., Velasco-Ramírez, P., & Hernández-Herrera, R. M. (2023). Application of seaweed generates changes in the substrate and stimulates the growth of tomato plants. *Plants*, 12(7), Article 1520. <https://doi.org/10.3390/plants12071520>
- Garcia-Gonzalez, J., & Sommerfeld, M. (2016). Biofertilizer and biostimulant properties of the microalga *Acutodesmus dimorphus*. *Journal of Applied Phycology*, 28, 1051–1061. <https://doi.org/10.1007/s10811-015-0625-2>
- Government of Mexico. (2024, February 20). *Agri-Food Panorama 2023* [Panorama Agroalimentario 2023]. <https://www.gob.mx/siap/acciones-y-programas/panorama-agroalimentario-258035>
- Hussain, H. I., Kasinadhuni, N., & Arioli, T. (2021). The effect of seaweed extract on tomato plant growth, productivity and soil. *Journal of Applied Phycology*, 33, 1305–1314. <https://doi.org/10.1007/s10811-021-02387-2>
- Ma, C., Song, W., Yang, J., Ren, C., Du, H., Tang, T., Quin, S., Liu, Z., & Cui, H. (2022). The role and mechanism of commercial macroalgae for soil conditioner and nutrient uptake catalyzer. *Plant Growth Regulation*, 97(3), 455–476. <https://doi.org/10.1007/s10725-022-00819-8>
- Suchithra, M. R., Muniswami, D. M., Sri, M. S., Usha, R., Rasheeq, A. A., Preethi, B. A., & Dineshkumar, R. (2022). Effectiveness of green microalgae as biostimulants and biofertilizer through foliar spray and soil drench method for tomato cultivation. *South African Journal of Botany*, 146, 740–750. <https://doi.org/10.1016/j.sajb.2021.12.022>