



World News of Natural Sciences

An International Scientific Journal

WNOFNS 27 (2019) 1-10

EISSN 2543-5426

Charcoal production: A promoter of deforestation in Nigeria

M. O. O. Isese

Forestry Research Institute of Nigeria, P.M.B 5054 Jericho Hills, Ibadan, Nigeria

E-mail address: Switmykel@mail.com

ABSTRACT

Charcoal production in Nigeria is a growing source of concern for government and forestry institutions because of its perceived impact on the environment. The agreed impact is deforestation, i.e., the clearance of indiscriminate cutting of forest trees. Forest degradation associated with charcoal production is difficult to monitor and overlooked and under-represented in forest cover change and carbon emission estimates. Emissions of greenhouse gases a consequence of deforestation from charcoal production in tropical ecosystems in 2009 are estimated at 71.2 million t for carbon dioxide and 1.3 million t for methane. Trees used for charcoal production are *Vitellaria paradoxa*, *Ficus sur*, *Tectona grandis*, *Khaya ivorensis*, *Bombax buonopozense* just to mention a few. This review aims to further discredit the act of indiscriminate forest tree falling for charcoal production which have proofed by various works as a promoter of deforestation.

Keywords: Deforestation, Charcoal production, Forest degradation, greenhouse gases, Tree plant, *Vitellaria paradoxa*, *Ficus sur*, *Tectona grandis*, *Khaya ivorensis*, *Bombax buonopozense*

1. INTRODUCTION

Charcoal is a fuel produced by carbonization of biomass. Although investment in charcoal production from forest plantations is increasing in tropical regions, mostly, biomass for charcoal production is obtained from natural forests in which natural regeneration is the main source of forest recovery. Charcoal is a key source of energy in African urban centers where 80% percent of the population uses it as the main source of energy for cooking (Zulu and Richardson, 2013). Population projections show an unprecedented increase in the urban

population in African cities, increasing from 30% in 2000 to reach 60% by year 2050 (Boko et al. 2007, Montgomery 2008, UN-HABITAT, 2010). The energy needs of the growing urban population will pose an increasing pressure on forest resources of rural areas (Barnes et al. 2004, Arnold et al., 2006, Grimm et al., 2008).

In many Sub-Saharan Africa, there is low energy consumption among household and in many other developing countries and this makes them too dependent upon wood fuels for their energy requirements (Arnold & Persson, 2005). The growing demand for charcoal in these countries has resulted in localized deforestation in vulnerable areas. The problem of fuel in Nigeria is becoming unbearable, resulting in constant deforestation to meet domestic needs.

Deforestation and forest degradation are the principal causes of forest cover change and account for a large proportion of global carbon emissions (Achard et al. 2007, van der Werf et al., 2009). Deforestation, defined as the long-term or permanent conversion of land from forest use to other non-forest uses (GOFCEGOLD 2009), represents an abrupt and rapid change in land cover. Population projections show an unprecedented increase in the urban population in African cities, increasing from 30% in 2000 to reach 60% by year 2050 (Boko et al. 2007, Montgomery 2008, UN-HABITAT 2010).

The energy needs of the growing urban population will pose an increasing pressure on forest resources of rural areas (Barnes et al. 2004, Arnold et al., 2006, Grimm et al., 2008). The charcoal sector in Africa is weakened by a lack of reliable information, because a very small fraction of charcoal produced is assessed and recorded.

Therefore, the actual magnitude of use and impacts on forest degradation or rural livelihoods has been a subject of debate (Zulu, 2013). The attempt to capture the impact of charcoal production on deforestation in Africa has been addressed by many scholars (Chidumaya & Gumbo, 2013). In Tanzania, charcoal production accounted for 30,000 ha per year of degraded closed woodland in 2005 and 2006 (Abdallah & Monela, 2007, Schafsma et.al, 2012). In Malawi, about 15,000 ha of forest have been cut down yearly due to charcoal production (Kambewa et.al, 2007).

This paper reviews traditional charcoal production methods and assesses this process from a deforestation view point. We base the paper on a literature review; it seeks to provide a broader perspective of the concerns surrounding charcoal production in Nigeria.

2. IMPACT OF CHARCOAL PRODUCTION ON DEFORESTATION

Deforestation is the clearing up of forest areas and converting them to other purposes like Agriculture, Urbanization and wood-fuel which includes charcoal production. This charcoal production contributes to deforestation for the main source of charcoal production is trees and to produce charcoal trees must be cut down.

About 31% of the world land surface is covered by forest, while in Nigeria it falls from 16.6% in 1996 to 7.7% in 2015, According to Food and Agricultural Organization of United Nations as of 2005 Nigeria has highest deforestation rate in the world at 12.2% equivalent of 11,089,000 hectares had been deforested and between (2000 to 2005). The link between charcoal production and deforestation have been demonstrated by several studies since the early 90s and this linkage is due to fact that deforestation frequently occurs in areas with intense charcoal production (Mwampamba et. al, 2013).

The common reason is that deforestation was always a result of agricultural expansion and logging. The need for new agricultural lands causes more damage than the charcoal production itself. Charcoal production is considered as merely a by-product of the deforestation process [French, 1986, Combes et. al., 2009]. Michael Daley, associate professor of environmental science at Lasell College in Newton, Massachusetts, the number one major problem caused by deforestation is the impact on the global carbon cycle, when this trees are been cut down those carbon they supposed to absorb as food through photosynthesis process will be trapped in the atmosphere and this contributes to the increases on carbon dioxide accumulated in the atmosphere. Moreover there are more emission also created when felled trees releases the carbon they had been storing and rot or burn on the forest floor, so not only that this trees act as carbon dioxide mitigation but cutting them down also increases carbon dioxide emission through carbon cycle process. Forest makes up about 30% of total land cover and are of incredible value to life on earth, they are carbon bank and play very important role in climate control, climate mitigation and effects of global warming, also act as watershed and with many raw materials that human depends on joined with the bio-diversities it contains. Trees lowers temperature through transpiration of water and shading of surface and also reduces heat sink while carbon dioxide makes trees healthier which in turn serve as climate change mitigation which are major causes of global warming.

In almost all countries where charcoal is produced there have been reports highlighting concern about deforestation and forest degradation that accompanies the production process (Hofstad et al., 2009). Forest degradation refers to less obvious changes in the woody canopy cover while deforestation is the more or less complete loss of forest cover that is often associated with forest clearance (Grainger, 1999). Degradation therefore represents the temporary or permanent reduction in the density, structure, species composition or productivity of vegetation cover. The most predominant species used for charcoal production in Nigeria are Teak and Shea tree as shown in Table 1; the local communities do not use Eucalyptus spp. for charcoal production, in areas where the preferred trees are scarce, several fruit trees are used to produce charcoal. Therefore, it is recommended that Eucalyptus could be an alternative tree for afforestation plans to curb deforestation.

Table 1. Tree name and species used for Charcoal production in Nigeria

Tree name	Species	Family
Shea Tree	<i>Vitellaria paradoxa</i>	Sapotaceae
Bush Fig Tree	<i>Ficus sur</i>	Moraceae
Kapok Tree	<i>Bombax buonopozense</i>	Bombacaceae
Mahogany	<i>Khaya ivorensis</i>	Meliaceae
Teak	<i>Tectona grandis</i>	Verbenaceae

Source: Fieldwork 2019.

3. GREEN HOUSE GASES EMISSION A BYPRODUCT OF DEFORESTATION: THE ROLE OF CHARCOAL

Charcoal from most earth-based kilns is produced in an oxygen-poor environment that results in the formation of products of incomplete combustion, such as methane. Charcoal production therefore affects global warming through the production and emission of greenhouse gases, such as carbon dioxide (CO₂) and methane (CH₄). Although carbon monoxide is one of the products of incomplete combustion during charcoal making, it is not listed as a greenhouse gas by the IPCC as it is considered being short-lived in the atmosphere (IPCC, 2007). Kammen and Lew (2005) have shown that emissions during charcoal production have a greater global warming contribution than emissions from charcoal burning. The emission factors for charcoal production have been calculated in a number of studies (Adam, 2009; Akagi et al., 2010; Pennise et al., 2001; Smith et al., 1999; Ward et al., 1999). To calculate emission quantities from charcoal production in tropical ecosystems, we used FAO estimates of wood-charcoal production in 2009 available at www.faostat.fao.org/faost. The values represented in Table 2 are the values of carbon emissions in Nigeria between 1960 and 2014. Charcoal-based energy demand has been identified as a mechanism of forest cover change in Africa (Hosonuma et al., 2012, Siteo et al., 2016). The emissions associated with this mechanism represent a key component of the large CO₂ emissions and emission uncertainties at the continental level (Ciais et al. 2011).

Table 2. Nigeria carbon Emission from 1960-2014

Year	Carbon dioxide emission (kt)
2014	96,280.0
2010	91,517.3
2005	106,068.0
2000	76,057.2
1995	35,841.3
1990	39,196.6
1985	69,893.0
1980	68,154.9
1975	47,396.0
1970	21,540.0
1965	11,763.7
1960	3,406.6

Source : Worldbank

4. POPULATION GROWTH AND ECONOMIC GROWTH: MAJOR FACTORS LIMITING CHARCOAL FUEL TRENDS

The key factors that affect the outlook of demand and supply include the current state of forest resources and their uses are population and income. Each of these factors affects both demand and supply of wood. It should be recognized that these factors differ in their effects on demand for fuelwood and demand for round wood although factors such as population growth and economic growth are important for both. The demand for industrial roundwood is derived from the demand for forest products. Therefore, an understanding of the dynamics of the supply and demand process of these markets can be brought to operation in understanding future developments for this component of wood demand and supply. This situation is contrary to the fuelwood demand structure and the process of supply is quite different from those four industrial products. Three billion people depend on wood to meet basic energy needs and much of the production of fuelwood is based on gathering from forests and scattered trees by individual households which promote forest degradation.

Population growth has historically been a major factor influencing wood consumption, and it is not likely that the close relationship between population growth and growth in consumption will change significantly in the foreseeable future. The world population increased from 2.4 billion people in 1950 to 5.5 billion people in the mid-1990s. The world population grows at a rate of 2.0% per year in the early 1970s and is currently increasing at a rate of 1.7% per year.

On a per capita basis, world consumption of timber, both fuelwood and industrial roundwood, declined slightly over the past four decades to 0.6 m³ per person from 0.7 m³ per person (Brooks et al., 1996). There has been changes in patterns of consumption (among developing and developed countries) and in the composition between fuelwood and industrial round wood, and among industrial products), but the net result has been little change in global per capita consumption.

Economic growth also have a particular impact on the demand for fuelwood, because the marginal propensity to increase consumption of commercial energy in developing countries is very high and highest among the poorest countries with currently high dependence on wood and biomass. The strong preference for convenience, efficiency and cleanliness in cooking and heating fuel may be expected to lead to the substitution of commercial fuels as income rises. Energy is of particular importance in the context of wood demand as it constitutes both a basic need for human civilization and an essential component in economic activity and development.

As already noted, charcoal production contributes to deforestation through the growing demand for wood biomass energy (Defries et al., 2010) as well as through fragmentation of habitats and forest degradation at the local level (Nellemann and Corcoran, 2010). Over the years, charcoal production has contributed to the loss of resilience in tropical forest ecosystems (Thompson et al., 2009).

Further, the value of total forest removals is increasingly being questioned while values of the contribution of employment and services have been further marginalized (FAO, 2010b). Also affected are market based mechanisms for the promotion of the forest sector e.g., ecotourism, sale of certified forest products, payments for environment services (Crowe and ten Kate, 2010). Charcoal making has been viewed as a leakage under REDD+ mechanisms thereby affecting any co-benefits that may be realized from such schemes (UNFCCC, 2010).



Fig. 1. Processes of charcoal production by traditional earth kiln. Upper images (II) show the process of log piling into clamp, and the lower left image (III) shows covering of the clamp with dung and soil lumps, while the lower right image (V) shows the harvesting of charcoal and packaging it into bags. The steps number (I, IV) is not represented in this figure.

Recommendations

- Since it could be difficult to completely eradicate tree felling for charcoal production due to the given the prevailing poverty rate in Nigeria, government should enforce tree planting policies to encourage afforestation.
- Public enlightenment and awareness should be carried out to educate the general public on the importance of agro-forestry. Forest and environmental conservation should be embarked upon by all tiers of government.
- Government should monitor tree cutters so as to ensure that not just single species of tree is being cut down. This will help to prevent the extinction of such endangered species

- Forestry research organizations should be funded for more sustainable research into the reduction of growth years of indigenous trees for a faster regeneration.
- Mechanisms for methane capture should be developed and adopted so as to collect the emissions during charcoal production process. This gas can be used to generate energy while protecting the environment from pollution and global warming.
- The charcoal producers through their associations should embark on a vast tree planting project and reduce dependence on natural forest degradation to protect the forest regions.

5. CONCLUSION

Due to high demand for cleaner energy, the demand for charcoal has extended from national to international market. This has further increased the dangers of deforestation in the country. It has been established that Charcoal production is a driver of deforestation around the world; hence to curb this disaster prohibition of charcoal production from unauthorized felling of trees must be enforced.

References

- [1] F. Achard, R. DeFries, H. Eva, M. Hansen, P. Mayaux and H-J. Stibig. Pan-tropical monitoring of deforestation. *Environ. Res. Lett.* 2007, 2, 045022
- [2] J.C. Adam. Improved and more environmentally friendly charcoal production system using a low-cost retort–kiln (Eco-charcoal). *Renew Energy* 2009, 34: 1923-5.
- [3] S.K Akagi, R.J Yokelson, C. Wiedinmyer, M.J Alvarado, J.S Reid, T. Karl, et al. Emission factors for open and domestic biomass burning for use in atmospheric models. *Atmos Chem Phys Discuss* 2010, 10: 27523-602.
- [4] M. Arnold and R. Persson. Reassessing the fuelwood situation in developing countries. *Int for Rev* 2003, 5: 379–83.
- [5] J.E.M Arnold, G. Köhlin and R. Persson. Woodfuels, livelihoods, and policy interventions: changing perspectives. *World Dev* 2006, 34, 596-611.
- [6] D.F Barnes, K. Krutilla and W.F. Hyde. *The Urban Household Energy Transition: Social and Environmental Impacts in the Developing World* (Washington, DC: RFF Press), 2004.
- [7] M. Crowe, K. Ten Kate. *Biodiversity offsets: policy options for government*. Washington, D.C.: Forest Trends; 2010.
- [8] M. Boko, I. Niang, A. Nyong, C. Vogel, A. Githeko, M. Medany, B. Osman-Elasha, R. Tabo and P. Yanda. *Africa Climate Change 2007: Impacts, Adaptation and Vulnerability*. Contribution of Working Group II to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change ed ML Parry et al (Cambridge: Cambridge University Press), 2007.

- [9] P. Ciais, A. Bombelli, M. Williams, S.L Piao, J. Chave, C.M Ryan, M. Henry, P. Brender and R. Valentini. The carbon balance of Africa: synthesis of recent research studies. *Phil. Trans. R. Soc. A*, 2011, 369, 2038-57
- [10] R.S DeFries, T. Rudel, M. Uriarte, M. Hansen. Deforestation driven by urban population growth and agricultural trade in the twenty-first century. *Nat Geosci* 2010, 3: 178-81.
- [11] FAO. (2010b). Criteria and indicators for sustainable woodfuels. Rome: Food and Agriculture Organization of the United Nations.
- [12] GOFC-GOLD, (2009). Reducing greenhouse gas emissions from deforestation and degradation in developing countries: a sourcebook of methods and procedures for monitoring, measuring and reporting GOFC-GOLD Report Version COP14-2 GOFC-GOLD Project Office, Natural Resources Canada, Alberta, Canada
- [13] N.B. Grimm et al. *Global change and the ecology of cities Science* 2008, 319, 756-60
- [14] A. Grainger. Constraints on modeling the deforestation and degradation of tropical open woodlands. *Glob Ecol Biogeogr* 1999, 8: 179-90.
- [15] Hosonuma N .et al, (2012). An assessment of deforestation and forest degradation drivers in developing countries. *Environ. Res. Lett.* 7, 044009
- [16] O. Hofstad, G. Kohlin, J. Namaalwa. How can emissions from woodfuel be reduced? In: A. Angelsen, M. Brockhaus, M. Kanninen, E. Sills, W.D Sunderlin, S. Wertz-Kanounnikoff, editors. Realizing REDD+: national strategy and policy options. Bogor, Indonesia: Center for International Forestry Research 2009. p. 237-49.
- [17] IPCC. Climate change: the physical science basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change. Available from <http://www.ipcc.ch> , 2007
- [18] D.M Kammen and D.J Lew, Review of technologies for the production and use of charcoal. Renewable and Appropriate Energy Laboratory report, Energy and Resources Group & Goldman School of Public Policy. Berkeley, CA, USA: University of California, 2005, pp. 1-19.
- [19] M.R Montgomery. The urban transformation of the developing world. *Science* 2008, 319, 761-4
- [20] Mwampamba, T.H.; Ghilardi, A.; Sander, K.; Chaix, K.J. Dispelling common misconceptions to improve attitudes and policy outlook on charcoal in developing countries. *Energy Sustain. Dev.* 2013, 17, 75–85
- [21] French, D. confronting an unsolvable problem: Deforestation in Malawi. *World Dev.* 1986, 14, 531–540
- [22] Combes Motel, P.; Pirard, R.; Combes, J.L. A methodology to estimate impacts of domestic policies on deforestation: Compensated Successful Efforts for “avoided deforestation” (REDD). *Ecol. Econ.* 2009, 68, 680-691

- [23] C. Nellemann, E. Corcoran, editors. Dead planet, living planet - biodiversity and ecosystem restoration for sustainable development. A rapid response assessment Arendal: United Nations Environment Programme, UNEP/GRID-Arendal, 2010.
- [24] D.M Pennise, K.R Smith, J.P Kithinji, M.E Rezende, T.J Raad, J. Zhang, et al., . Emissions of greenhouse gases and other airborne pollutants from charcoal making in Kenya and Brazil. *J Geophys Res* 2001, 106, 24143-55
- [25] K.R Smith, D.M Pennise, P. Khummongkol, V. Chaiwong, K. Ritgeen, and J. Zhang, et al., Greenhouse gases from small-scale combustion in developing countries: charcoal-making kilns in Thailand. Report EPA-600/R-99-109. Research Triangle Park, N.C.: U.S. Environmental Protection Agency, 1999.
- [26] A. Siteo, I. Remane, R. Ribeiro, M.P Falcão, R. Mate, J. Nhamirre, S. Walker, L. Murray and J. Joana Melo. Identificação e análise dos agentes e causas directas e indirectas de desmatamento e degradação florestal em Moçambique Final Report Maputo (in Portuguese), 2016.
- [27] UNFCCC. Outcome of the Ad Hoc Working Group on Long-term Cooperative Action under the Convention. United Nations Framework Convention on Climate Change; http://unfccc.int/meetings/cop_16/items/5571.php , 2010.
- [28] UN-HABITAT, (2010). The State of African cities 2010: Governance, inequality and urban land markets (Nairobi: UN-HABITAT)
- [29] G.R Van der Werf, D.C Morton, R.S DeFries, L. Giglio, J.T Randerson, G.J Collatz and P.S Kasibhatla. Estimates of fire emissions from an active deforestation region in the southern Amazon based on satellite data and biogeochemical modeling. *Biogeosciences* 2009, 6, 235-49
- [30] D.E Ward, E. Chidumayo, S.P Baker, R.A Susott. Charcoal production and smoke emissions from earthen kilns in Zambia. Final report submitted to Environmental Research Laboratory, 1 College Station Road, Athens, GA, USA; 1999. World Bank Energy Sector Management Assistance Programme (World Bank/ESMAP). Sustainable woodfuel supplies from the dry tropical woodlands. ESMAP technical paper 013. Washington, DC: World Bank/ESMAP, 2001.
- [31] L.C Zulu and R.B Richardson. Charcoal, livelihoods, and poverty reduction: Evidence from sub-Saharan Africa. *Energy Sustain. Dev.* 2013, 17, 127-37.
- [32] Chidumayo, E.N.; Gumbo, D.J. The environmental impacts of charcoal production in tropical ecosystems of the world: A synthesis. *Energy Sustain. Dev.* 2013, 17, 86-94.
- [33] Abdallah, J.; Monela, G. Overview of Miombo woodlands in Tanzania. *In Proceedings of the First MITMIOMBO Project Workshop, Morogoro, Tanzania*, 6-12 February 2007; Volume 50, pp. 9-23.
- [34] Schaafsma, M.; Morse-Jones, S.; Posen, P.; Swetnam, R.D.; Balmford, A.; Bateman, I.J.; Burgess, N.D.; Chamshama, S.A.O.; Fisher, B.; Green, R.E.; et al. Towards transferable functions for extraction of Non-timber Forest Products: A case study on charcoal production in Tanzania. *Ecol. Econ.* 2012, 80, 48-62.

- [35] Kambewa, P.S.; Mataya, B.F.; Sichinga, W.K.; Johnson, T.R. Charcoal: The Reality - A Study of Charcoal Consumption, Trade and Production in Malawi; International Institute for Environment and Development: London, UK, 2007; ISBN 978-1-84369-678-0.