

CHLOROPHYLL CONTENT OF APHID-INFESTED SEEDLING LEAVES OF FIFTEEN MAIZE GENOTYPES

HUBERT SYTYKIEWICZ*, PAWEŁ CZERNIEWICZ, IWONA SPRAWKA,
AND ROBERT KRZYŻANOWSKI

*Department of Biochemistry and Molecular Biology,
University of Natural Sciences and Humanities in Siedlce,
Prusa 12, 08-110 Siedlce, Poland*

Received May 10, 2013; revision accepted August 20, 2013

We measured the total chlorophyll (Chl *a+b*) content in seedling leaves of fifteen maize cultivars infested by two studied aphid species (oligophagous *Rhopalosiphum padi* L., monophagous *Sitobion avenae* F.) 7 and 14 days after the beginning of infestation, using a SPAD-502 chlorophyll meter. Chlorophyll loss was more severe in *R. padi*-infested than in *S. avenae*-infested plants. Chlorophyll depletion was greater after long-term (14 days) than after short-term aphid infestation in the investigated host systems. Seedlings of Złota Karłowa and Tasty Sweet were more damaged by aphid feeding; Ambrozja and Płomyk plants were less damaged by aphid feeding.

Key words: Total chlorophyll (Chl *a+b*) content, SPAD units, maize, *Rhopalosiphum padi*, *Sitobion avenae*.

INTRODUCTION

Maize is one of the most important crop species, widely cultivated throughout the world. *Zea mays* is increasingly used in bioethanol and biogas production and in the pulp and paper industry. Maize can grow under different climatic conditions, but high temperature and humidity can accelerate the population growth of many phytophagous insects that infest this host (Fornalé et al., 2012; Herrmann, 2013; Reddy et al., 2013; Semenčenko et al., 2013). Aphids (Hemiptera, Aphidoidea) represent the major group of arthropods heavily infesting *Z. mays* cultivars (Dahmardeh et al., 2010; Khatoun et al., 2010, Lewis et al., 2010). In Poland, maize plants are colonized by four aphid species: *Metopolophium dirhodum* Walk., *Rhopalosiphum maidis* F., *Rhopalosiphum padi* L. and *Sitobion avenae* F. (Pieńkosz et al., 2005; Strażyński, 2008). This group of highly specialized phloem-feeding hemipterans has coevolved multi-level adaptations facilitating exploitation of the resources provided by their host plants (Czerniewicz et al., 2011a; Sprawka et al., 2011; Chrzanowski et al., 2012; Sprawka et al., 2012; 2013a). These insects cause a wide spectrum of detrimental effects in attacked organs, including mechanical disruption of penetrated tissues, depletion of pho-

toassimilates, and intensification of many intracellular processes (Sempruch et al., 2010; Sytykiewicz et al., 2011a; Goławska et al., 2012). Aphid salivary secretions contain a variety of hydrolytic enzymes and biologically active substances that modulate metabolic reactions within the host. In some cases, aphid infestation contributes to the activation of premature senescing process and programmed cell death in plant organs (Carolan et al., 2009; Anstead et al., 2010; Sytykiewicz et al., 2011b; Sprawka et al., 2013b). Colonization of the different host systems may lead to transmission of a wide range of pathogenic plant viruses, such as barley yellow dwarf virus (BYDV), bean leafroll virus (BLRV), beet western yellows virus (BWYV), chickpea chlorotic stunt virus (CpCSV), faba bean necrotic yellows virus (FBNYV), maize dwarf mosaic virus (MDMV), soybean dwarf virus (SbDV) and sugarcane mosaic virus (SCMV) (Makkouk and Kumari, 2009; Higashi and Bressan, 2012; Stewart et al., 2012; Zielińska et al., 2012; Ferriol et al., 2013).

Changes in the total chlorophyll (Chl *a+b*) content of foliar tissues is an important indicator of disturbed chloroplast development and impaired photosynthetic capacity in plants exposed to a broad spectrum of biotic and abiotic stressors. Many studies have shown that aphid infestation can trigger

*e-mail: hubert.sytykiewicz@uph.edu.pl

severe Chl breakdown within the host. An imbalance between biosynthesis and catabolic turnover of green pigments in plant tissues indicates profound inhibition of photosynthesis process (Heng-Moss et al., 2003; Smith et al., 2005; Botha et al., 2006; Diaz-Montano et al., 2007; Goławska et al., 2010). Portable leaf-clip-type chlorophyll meters perform rapid, repetitive and nondestructive estimates of Chl concentrations in living plants (Hawkins et al., 2009; Jinwen et al., 2009; Coste et al., 2010; Gholizadeh et al., 2011; Shrestha et al., 2012; Ghosh et al., 2013). The use of these reliable and advanced instruments for measurement of green pigment content is not restricted to controlled laboratory bioassays but can be extended to experiments in the plants' natural environment (Uddling et al., 2007; Ruiz-Espinoza et al., 2010; Neto et al., 2011; Cerovic et al., 2012). SPAD-502 chlorophyll meters have been used to obtain accurate measurements in arbitrary SPAD units reflecting absolute foliar Chl concentrations (Arunyanark et al., 2008; Marengo et al., 2009; Mielke et al., 2010; Ling et al., 2011; Shrestha et al., 2012). A significant linear relationship between SPAD readings and leaf nitrogen status in tissues of monocotyledonous species has been documented (Rostami et al., 2008; Swain and Sandip, 2010; Liu et al., 2012; Ghosh et al., 2013). To the best of our knowledge there are no published data from studies using a SPAD-502 meter to assess aphid-affected changes in the Chl content of tissues of maize plants representing a variety of genotypes. A wide range of maize cultivars presumably should reflect different levels of susceptibility to damage from aphid colonization and consequently different degrees of Chl decrement in infested leaves. In this study we compared the total Chl ($a+b$) content of seedling leaves of fifteen *Z. mays* genotypes infested by two aphid species, oligophagous *R. padi* and monophagous *S. avenae*. We also monitored the increase in size of the aphid populations on *Z. mays* plants. The relationship between the total Chl content of seedling leaves (nondestructive SPAD readings) and Chl $a+b$ concentrations measured spectrophotometrically in leaf extracts of uninfested (control) maize plants is estimated in this study.

MATERIALS AND METHODS

PLANT MATERIAL

Seeds of the fifteen *Z. mays* genotypes used in this study (Ambrozja, Anawa, Delitop, Dobosz, Eleganza, Gucio, Makila, Nana, Narew, Płomyk, Rataj, Tasty Sweet, Touran, Waza, Złota Karłowa) were obtained from Syngenta (Warsaw, Poland),

Hodowla Roślin Smolice, Grupa IHAR (Smolice, Poland), KWS Polska (Poznań, Poland), Hodowla Roślin Snowidza (Snowidza, Poland) and local garden shops. The seeds were germinated and seedlings were grown in a climate chamber at $22\pm 2^\circ\text{C}/16\pm 2^\circ\text{C}$ (day/night), $65\pm 5\%$ relative humidity, $100\ \mu\text{M m}^{-2}\ \text{s}^{-1}$ light intensity and a long-day photoperiod (L16:D8). Maize seedlings were individually planted in round plastic pots (10 cm diam., 9 cm high) filled with general-purpose horticultural substrate widely used for greenhouse experiments, with no supplementary fertilization.

APHIDS

Wingless parthenogenetic females (apterae) of the bird cherry-oat aphid (*Rhopalosiphum padi* L.) and the grain aphid (*Sitobion avenae* F.) were collected from infested cereal plants growing in Siedlce district. The aphid colonies were maintained for a year on seedlings of winter wheat cv. Tonacja in the Department of Biochemistry and Molecular Biology, University of Natural Sciences and Humanities in Siedlce. The aphids were reared in an environmental chamber under the same conditions as described for seedling cultivation. Apterous 1- or 2-day-old adults of the two aphid species were used for the bioassays.

INFESTATION EXPERIMENT

Leaves of 14-day-old seedlings of each maize cultivar were infested with adult apterous females of *R. padi* or *S. avenae* – 10 (I-10) or 40 (I-40) individuals per plant. Uninfested seedlings were the control. Aphid-stressed and control maize plants were isolated in gauze-covered plastic cylinders (20×50 cm). The density of the insect populations developing on the *Z. mays* plants was monitored 7 and 14 days post infestation (dpi).

NONDESTRUCTIVE CHLOROPHYLL MEASUREMENT

The total chlorophyll (Chl $a+b$) content of seedling leaves of the fifteen maize genotypes was measured in situ with a SPAD-502 meter (Minolta Corp., Ramsey, NJ). This instrument measures SPAD units, defined as the ratio of light transmittance at two peak wavelengths (650 and 940 nm) (Goławska et al., 2010; Mielke et al., 2010; Filipović et al., 2013). The total Chl content values given here are means of 25 SPAD readings from different locations of both leaf sides. In making the measurements we avoided the vein areas of the leaf blades. The relative concentration of green pigments in foliar tissues of aphid-damaged and control maize plants was estimated at 7 and 14 dpi.

TABLE 1. Total chlorophyll content (SPAD units, mean \pm SD) in leaves of *R. padi*-stressed maize seedlings

Maize genotype	7 dpi			14 dpi		
	U	I-10	I-40	U	I-10	I-40
Ambrozja	27.6 \pm 2.1	27.5 \pm 2.0	26.8 \pm 1.5	30.1 \pm 2.9	29.2 \pm 2.3	27.5 \pm 1.8
Anawa	26.8 \pm 1.8	26.7 \pm 1.5	26.1 \pm 1.2	29.3 \pm 2.7	29.0 \pm 2.0	28.2 \pm 2.3
Delitop	28.4 \pm 3.5	27.4 \pm 2.2	25.6 \pm 1.8	35.5 \pm 3.8	33.7 \pm 2.9	30.4 \pm 2.7*
Dobosz	25.7 \pm 1.3	24.3 \pm 1.4	22.8 \pm 0.9*	27.3 \pm 1.8	24.6 \pm 1.2*	22.3 \pm 0.9*
Eleganza	24.6 \pm 1.1	23.6 \pm 1.1	22.5 \pm 0.8	28.7 \pm 2.3	27.2 \pm 1.6	25.3 \pm 1.4*
Gucio	27.5 \pm 2.3	26.0 \pm 1.7	23.2 \pm 1.3*	33.5 \pm 3.1	31.4 \pm 2.8	27.9 \pm 1.9*
Makila	28.9 \pm 3.1	26.9 \pm 1.9	25.3 \pm 1.0*	32.8 \pm 2.7	30.3 \pm 2.6	27.5 \pm 1.6*
Nana	27.2 \pm 2.5	25.7 \pm 1.4	23.2 \pm 1.4*	30.5 \pm 2.9	28.4 \pm 1.4	25.9 \pm 1.0*
Narew	22.1 \pm 0.9	22.1 \pm 0.8	22.1 \pm 0.8	25.6 \pm 1.3	25.6 \pm 1.3	25.6 \pm 1.3
Plomyk	26.2 \pm 1.7	26.0 \pm 1.8	25.1 \pm 1.4	29.4 \pm 2.5	28.8 \pm 1.7	27.5 \pm 1.2
Rataj	25.3 \pm 1.4	25.3 \pm 1.2	25.3 \pm 1.6	28.9 \pm 2.2	28.9 \pm 1.9	28.9 \pm 2.4
Tasty sweet	28.4 \pm 3.5	25.5 \pm 1.0*	22.5 \pm 1.2*	30.1 \pm 2.8	23.3 \pm 1.0*	20.5 \pm 0.9*
Touran	27.5 \pm 2.6	25.8 \pm 1.7	24.0 \pm 1.1*	32.7 \pm 3.0	30.1 \pm 2.5	28.1 \pm 2.1*
Waza	26.0 \pm 1.5	25.4 \pm 1.5	24.2 \pm 1.5	29.8 \pm 2.1	28.5 \pm 1.5	27.4 \pm 1.4
Złota Karłowa	27.9 \pm 2.8	24.6 \pm 1.1*	20.3 \pm 0.7*	28.4 \pm 2.4	23.9 \pm 1.2*	19.1 \pm 0.8*

* $P \leq 0.01$ (Student's t-test) – significant difference in leaf total Chl between aphid-stressed and uninfested seedlings; dpi – days post initial infestation; U – uninfested plants; I-10 – seedlings initially infested with 10 aphids; I-40 – seedlings initially infested with 40 aphids.

TISSUE-DESTRUCTIVE CHLOROPHYLL QUANTIFICATION

Freshly collected control samples (10 g) of seedling leaves of five maize genotypes (Ambrozja, Plomyk, Tasty Sweet, Waza, Złota Karłowa) were homogenized at 4°C for 5 min in 40 cm³ ice-cold 80% (v/v) acetone. The extracts were filtered through two layers of gauze and then centrifuged at 5000 \times g for 10 min. The pellet was discarded and the supernatants were analyzed. Chlorophyll *a* and *b* content was determined by measurement of absorbance at 644, 662 and 750 nm using a Hewlett-Packard UV-Vis spectrophotometer (model 8453). The levels of Chl *a* and *b* in the tested samples were estimated by the following empirical formulas (Ihl et al., 2000):

$$\text{Chl } a \text{ (nmol cm}^{-3}\text{)} = 11.30 \times (A_{662} - A_{750}) - 1.11 \times (A_{644} - A_{750})$$

$$\text{Chl } b \text{ (nmol cm}^{-3}\text{)} = 18.07 \times (A_{644} - A_{750}) - 2.87 \times (A_{662} - A_{750})$$

Total Chl (*a*+*b*) content in the seedling leaves is expressed in $\mu\text{g g}^{-1}$ fresh weight (FW).

STATISTICAL ANALYSIS

The experiments were completely randomized and performed in three independent replicates. Each bioassay variant (aphid infestation level – exposure time) and control (nonstressed plants) comprised 10 maize seedlings of similar height. All data given are means \pm SD. The significance of differences in total Chl content (SPAD units) between aphid-stressed and uninfested seedling leaves was ana-

lyzed with Student's t-test. A linear regression model was used to test the correlation between relative SPAD values and spectrophotometrically determined Chl *a*+*b* concentrations ($\mu\text{g g}^{-1}$ FW) in acetone extracts of seedling leaves of uninfested maize plants. The significance of differences in the number of aphids on the maize cultivars was checked by one-way ANOVA followed by Duncan's multiple range test (DMRT). Significance was deemed at $p \leq 0.01$. The statistical calculations were done with STATISTICA 9.0 (StatSoft Poland).

RESULTS

The results on total Chl content (SPAD units) in seedling leaves of fifteen tested maize cultivars colonized by two aphid species (*Rhopalosiphum padi*, *Sitobion avenae*) are presented in Tables 1 and 2. Long-term aphid infestation (14 dpi) and higher initial number of insects (40 per plant, I-40) resulted in significantly greater diminution of chlorophyll than short-term (7-day) infestation and lower initial number of insects (10 per plant, I-10). This was true for both aphid species, but the extent of chlorophyll depletion was markedly higher in *R. padi*-stressed than in *S. avenae*-stressed plants: 14-day *R. padi* feeding caused more than 10% reduction of foliar Chl levels in seedlings of nine cultivars (Delitop, Dobosz, Eleganza, Guccio, Makila, Nana, Tasty Sweet, Touran, Złota Karłowa), whereas *S. avenae*

TABLE 2. Total chlorophyll amount (SPAD units, mean \pm SD) in leaves of *S. avenae*-infested maize seedlings

Maize genotype	7 dpi			14 dpi		
	U	I-10	I-40	U	I-10	I-40
Ambrozja	27.6 \pm 3.5	27.1 \pm 2.0	25.9 \pm 1.1	30.1 \pm 2.7	29.4 \pm 1.7	28.0 \pm 1.5
Anawa	26.8 \pm 2.6	26.8 \pm 1.9	26.6 \pm 1.9	29.3 \pm 2.4	29.2 \pm 1.5	28.6 \pm 1.9
Delitop	28.4 \pm 4.2	27.2 \pm 2.5	26.3 \pm 1.5	35.5 \pm 3.2	33.9 \pm 2.8	32.4 \pm 2.8
Dobosz	25.7 \pm 3.6	24.1 \pm 1.2	22.7 \pm 0.8*	27.3 \pm 1.5	25.2 \pm 0.7	23.7 \pm 1.1*
Eleganza	24.6 \pm 2.2	24.0 \pm 0.9	23.1 \pm 0.9	28.7 \pm 1.8	27.9 \pm 0.9	26.5 \pm 1.5
Gucio	27.5 \pm 3.9	25.9 \pm 1.3	24.2 \pm 0.9*	33.5 \pm 3.0	31.2 \pm 2.5	28.2 \pm 1.6*
Makila	28.9 \pm 4.7	27.7 \pm 2.2	26.3 \pm 1.2	32.8 \pm 2.9	30.9 \pm 2.0	28.5 \pm 2.1*
Nana	27.2 \pm 2.3	26.4 \pm 1.7	25.7 \pm 1.0	30.5 \pm 2.8	29.2 \pm 1.8	27.5 \pm 1.4
Narew	22.1 \pm 1.4	22.1 \pm 0.8	22.1 \pm 0.7	25.6 \pm 1.2	25.6 \pm 1.2	25.6 \pm 1.3
Płomyk	26.2 \pm 1.7	26.0 \pm 1.5	25.2 \pm 1.2	29.4 \pm 2.7	29.0 \pm 1.4	28.2 \pm 1.6
Rataj	25.3 \pm 3.8	25.3 \pm 1.2	25.3 \pm 1.3	28.9 \pm 2.6	28.9 \pm 1.1	28.9 \pm 1.9
Tasty sweet	28.4 \pm 4.3	26.2 \pm 1.6	23.8 \pm 0.9*	30.1 \pm 2.8	27.3 \pm 0.9*	21.5 \pm 0.8*
Touran	27.5 \pm 4.1	26.1 \pm 1.3	25.0 \pm 1.0	32.7 \pm 3.0	30.4 \pm 1.8	27.9 \pm 1.4*
Waza	26.0 \pm 3.2	25.1 \pm 1.0	24.2 \pm 0.9	29.8 \pm 2.7	28.5 \pm 1.2	27.3 \pm 1.0
Złota Karłowa	27.9 \pm 2.9	25.4 \pm 1.1	23.1 \pm 0.8*	28.4 \pm 2.3	24.9 \pm 0.6*	19.4 \pm 0.6*

* $P \leq 0.01$ (Student's t-test) – significant difference in leaf total Chl between aphid-stressed and uninfested seedlings; dpi – days post initial infestation; U – uninfested plants; I-10 – seedlings initially infested with 10 aphids; I-40 – seedlings initially infested with 40 aphids.

TABLE 3. Abundance of *R. padi* populations developing on the investigated maize cultivars

Maize genotype	7 dpi		14 dpi	
	I-10	I-40	I-10	I-40
Złota Karłowa	31 \pm 2.20 ^a	84 \pm 4.56 ^a	122 \pm 8.54 ^a	217 \pm 11.52 ^a
Tasty sweet	25 \pm 1.37 ^b	67 \pm 3.78 ^b	98 \pm 7.29 ^b	164 \pm 9.38 ^b
Dobosz	14 \pm 0.85 ^c	52 \pm 3.15 ^c	61 \pm 5.2 ^c	115 \pm 6.53 ^c
Gucio	13 \pm 0.62 ^c	46 \pm 2.94 ^d	55 \pm 4.9 ^d	98 \pm 5.9 ^d
Makila	10 \pm 0.38 ^{cd}	35 \pm 2.72 ^e	33 \pm 2.68 ^e	90 \pm 4.74 ^{de}
Touran	8 \pm 0.21 ^d	28 \pm 2.16 ^f	30 \pm 2.10 ^e	76 \pm 3.92 ^f
Nana	8 \pm 0.14 ^d	23 \pm 1.50 ^{fg}	27 \pm 1.19 ^f	69 \pm 3.61 ^g
Eleganza	7 \pm 0.19 ^{de}	21 \pm 1.32 ^g	23 \pm 1.05 ^g	52 \pm 2.30 ^h
Delitop	6 \pm 0.09 ^e	24 \pm 1.13 ^{fg}	20 \pm 0.94 ^g	45 \pm 2.14 ^{hi}
Płomyk	6 \pm 0.08 ^e	19 \pm 0.32 ^g	2 \pm 0.19 ⁱ	14 \pm 1.27 ^k
Waza	5 \pm 0.06 ^e	22 \pm 0.94 ^g	16 \pm 4.70 ^{gh}	31 \pm 1.95 ^j
Ambrozja	5 \pm 0.07 ^e	13 \pm 0.28 ^h	3 \pm 0.24 ⁱ	9 \pm 0.86 ^l
Anawa	2 \pm 0.04 ^f	8 \pm 0.17 ⁱ	–	–
Narew	–	–	–	–
Rataj	–	–	–	–

Values are means \pm SD. Means bearing the same letter within column do not differ significantly by Duncan's MRT test at $P \leq 0.01$, dpi – days post initial infestation; dash indicates the absence of aphids; I-10 – seedlings initially infested with 10 aphids; I-40 – seedlings initially infested with 40 aphids.

infestation led to such a Chl decline in tissues of six cultivars (Dobosz, Guccio, Makila, Tasty Sweet, Touran, Złota Karłowa). The decrement in green pigment levels was highest in insect-stressed seedlings of

Złota Karłowa and Tasty Sweet (~30% decrease at 14 dpi, I-40). The Chl content of the Rataj and Narew cultivars was unaffected by aphid infestation. The other investigated cultivars varied in the extent of chloro-

TABLE 4. Abundance of *S. avenae* populations developing on the investigated maize cultivars

Maize genotype	7 dpi		14 dpi	
	I-10	I-40	I-10	I-40
Złota Karłowa	17±1.38 ^a	38±2.19 ^a	86±5.17 ^a	109±7.25 ^a
Tasty sweet	14±0.65 ^b	33±1.57 ^b	57±4.39 ^b	96±6.47 ^b
Dobosz	9±0.58 ^c	29±1.35 ^c	44±3.12 ^c	85±6.12 ^c
Gucio	7±0.52 ^d	24±1.18 ^d	30±2.85 ^d	52±4.83 ^d
Makila	7±0.49 ^d	20±0.83 ^e	23±2.67 ^e	34±2.71 ^e
Touran	6±0.42 ^{de}	16±0.62 ^g	19±1.34 ^{ef}	30±2.38 ^e
Nana	5±0.36 ^e	19±0.71 ^{ef}	17±0.92 ^f	25±1.94 ^f
Eleganza	4±0.30 ^f	13±0.57 ^h	11±0.76 ^g	22±1.73 ^{fg}
Delitop	4±0.21 ^f	11±0.48 ⁱ	15±0.80 ^f	19±1.24 ^g
Waza	4±0.27 ^f	6±0.23 ^{jk}	10±0.64 ^g	17±0.96 ^g
Płomyk	3±0.18 ^g	8±0.35 ^j	1±0.03 ^h	6±0.49 ^h
Ambrozja	2±0.15 ^h	5±0.19 ^k	1±0.08 ^h	1±0.09 ⁱ
Anawa	1±0.07 ⁱ	3±0.18 ^l	–	–
Narew	–	–	–	–
Rataj	–	–	–	–

Values are means ±SD. Means bearing the same letter within column do not differ significantly by Duncan's MRT test at $P \leq 0.01$. dpi – days post initial infestation; dash indicates the absence of aphids; I-10 – seedlings initially infested with 10 aphids; I-40 – seedlings initially infested with 40 aphids.

phyll loss in response to aphid feeding. Long-term *R. padi* colonization (14 dpi, I-40) caused Chl reduction ranging from 3.7% (Anawa) to 18.3% (Dobosz); the decline in *S. avenae*-attacked seedlings ranged from 2.4% (Anawa) to 13.2% (Dobosz).

The *R. padi* populations on the *Z. mays* seedlings reached higher abundance than the *S. avenae* populations (Tabs. 3, 4). Adult apterous females of both aphid species produced no offspring on Narew and Rataj plants, and there were no aphids on seedlings of cv. Anawa at 14 dpi, indicating this genotype's high resistance to these aphids. Aphid density per plant at 14 dpi was low on Ambrozja (9 *R. padi*, 1 *S. avenae*) and Płomyk (14 *R. padi*, 6 *S. avenae*). The other tested cultivars differed greatly in aphid abundance. For both aphid species at 14 dpi it was highest on Złota Karłowa (217 *R. padi*, 109 *S. avenae*) and Tasty Sweet (164 *R. padi*, 96 *S. avenae*).

We used the uninfested controls to check the accuracy of the SPAD readings versus spectrophotometric measurements of total Chl content in tissues of all tested *Z. mays* varieties. Table 5 presents the correlations between mean SPAD values and chlorophyll *a+b* amounts ($\mu\text{g}\cdot\text{g}^{-1}$ FW) in uninfested seedling leaves. Statistical analyses showed a strong positive correlation between the noninvasive SPAD readings and spectrophotometrically determined chlorophyll concentrations in acetone extracts prepared from seedlings of the studied *Z. mays* cultivars ($p \leq 0.01$).

TABLE 5. Correlation between total chlorophyll content (SPAD units) and chlorophyll *a+b* amount ($\mu\text{g}\cdot\text{g}^{-1}$ FW) in uninfested (control) seedling leaves of selected maize cultivars

Maize genotype	R ²
	SPAD-Chl abs
Ambrozja	0.98
Anawa	0.93
Delitop	0.95
Dobosz	0.91
Eleganza	0.94
Gucio	0.92
Makila	0.95
Nana	0.93
Narew	0.96
Płomyk	0.95
Rataj	0.96
Tasty sweet	0.94
Touran	0.92
Waza	0.95
Złota Karłowa	0.90

SPAD-Chl abs represents the relationship between SPAD values and spectrophotometrically determined chlorophyll *a+b* concentrations in acetone extracts of seedling leaves of uninfested maize plants; FW – fresh weight; R² – coefficient of determination (linear regression model).

DISCUSSION

There are many recent reports on the destructive effects of aphid infestation on the functioning of various mono- and dicotyledonous plant systems (e.g., Czerniewicz et al., 2011b; Sempruch et al., 2011; Chen et al., 2012; Goławska et al., 2012; Łukasik et al., 2012; Sempruch et al., 2012; Bak et al., 2013; Riedell et al., 2013). One such effect is chlorosis (Heng-Moss et al., 2003; Smith et al., 2005; Botha et al., 2006; Diaz-Montano et al., 2007; Goławska et al., 2010). Highly bioactive effector molecules in aphid salivary gland secretions may cause severe damage to infested plants (Cooper et al., 2010; Cooper et al., 2011; Nicholson et al., 2012; Rao et al., 2013). Although *Z. mays* is one of the most important model plants and a valuable crop, there has been no published assessment of aphid-evoked changes in the total chlorophyll concentration of seedlings representing a wide range of genotypes. In this study we assessed the tolerance of a diverse range of maize varieties to infestation by aphids (*Sitobion avenae*, *Rhopalosiphum padi*). *S. avenae* is a monophagous and monoecious insect infesting a large array of plants of the family Poaceae (Gao and Liu, 2013; Li et al., 2013; Svobodová et al., 2013), whereas the life cycle of oligophagous *R. padi* involves seasonal migration between two groups of taxonomically distant host plants. The primary hosts of *R. padi* are a few members of the genus *Prunus*, and the secondary hosts include a broad range of cereals and grasses (Stoetzel and Miller, 2001; Aslan and Uygun, 2005; Łukasik 2009; Coulette et al., 2013).

In the maize cultivars we examined the extent of chlorophyll depletion in response to aphid feeding depended on the genotype. Oligophagous *R. padi* caused more Chl *a+b* loss than monophagous *S. avenae* in the stressed seedlings. Riedell and Blackmer (1999) reported similar results in wheat cv. Sharp infested by greenbug (*Schizaphis graminum* Rond.) and Russian wheat aphid (*Diuraphis noxia* Mordv.): Chl *a+b* content was lower under aphid feeding than in the nonstressed control, and the loss was greater under *S. graminum* than under *D. noxia* feeding. More green pigment was lost from mature (fully expanded) wheat leaves than from young leaves. In our work the extent of Chl loss in foliar tissues depended on the initial abundance of hemipterans on the maize seedlings and the duration of exposure to feeding. Similarly, Heng-Moss et al. (2003) found that prolonged *D. noxia* feeding on wheat leaves caused more Chl loss (~5-fold decrease after 13 days) than after short-term infestation. In their study, Russian wheat aphid feeding caused significant loss of chlorophyll content in susceptible Betta wheat but aphid-resistant Betta-*Dn1* wheat showed

only slightly lower total Chl than uninfested plants. Chl levels in *D. noxia*-tolerant Betta-*Dn2* and control plants were quite similar, but 11 and 13 days after the initial infestation the aphid-infested Betta-*Dn2* leaves showed higher Chl content than the uninfested control. Haile et al. (1999) reported that *D. noxia*-tolerant wheat line PI262660 completely recovered its photosynthetic capacity 7 days after aphids were removed from the colonized plants; there was no such compensation reaction in tissues of susceptible Arapahoe wheat or introduction line PI137738, showing a high level of antibiosis. In experiments by Smith et al. (2005), *D. noxia*-stressed leaves of wheat cv. Wichita showed a ~53% decline in total Chl content (SPAD units) versus control leaves; isogenic lines PI 372129 and P243781 showed Chl loss of 53% and 21% respectively. In leaves of two *Triticum aestivum* cultivars, Tugela-DN (resistant) and Tugela (susceptible), Botha et al. (2006) found that *D. noxia* feeding caused a gradual decrement of total chlorophyll versus the uninfested controls during 16 days of colonization. The decrease was greater in the susceptible cultivar than in Tugela-DN. After 7 days of infestation by the pea aphid (*Acyrtosiphon pisum* Harr.), Chl *a+b* content was lower in leaves of four legume hosts: alfalfa (*Medicago sativa* L. var. Radius), clover (*Trifolium pratense* L. var. Bona), pea (*Pisum sativum* L. var. Tulipan) and vetch (*Vicia faba* L. var. Jaga) (Goławska et al., 2010). Interestingly, in that experiment the decline was lower after 17 days than after 7 days.

Even a small population of the soybean aphid (*Aphis glycines* Mats.) significantly inhibited photosynthetic processes in attacked soybean plants (Macedo et al., 2003): there was high suppression of gas exchange in plant tissues under an intermediate level of aphid infestation (20–49 aphids per leaflet), and medium and high *A. glycines* densities (≥ 50 aphids per leaflet) caused an increase of F_o (nonvariable fluorescence) values, indicating damage to the PSII reaction system. In susceptible *T. aestivum* cultivars colonized by *D. noxia*, Burd and Elliott (1996) found elevated F_o levels, which may disrupt electron transfer in the PSII system and reduce D1 protein synthesis. Soybean aphid feeding altered the quenching coefficients (qP, coefficient of photochemical fluorescence quenching; qN, coefficient of non-photochemical fluorescence quenching) in tissues of infested soybean plants, according to Macedo et al. (2003), who suggested that aphid colonization may suppress the xanthophyll cycle, which participates in protecting photosystem II under stress conditions by modulating the thylakoid membrane pH gradient. Haile and co-workers (1999) reported substantial reduction of photosynthetic efficiency in wheat plants attacked by *D. noxia*. Some authors have speculated that the Chl loss in aphid-injured plants

may be connected with elevated biosynthesis of different defensive secondary metabolites such as saponins (Haile et al., 1999; Goławska et al., 2010).

Feeding by *S. graminum* biotype E on leaves of winter wheat cv. Sturdy and Largo for one hour resulted in plasmolysis and initial degeneration of cell organelles of the parenchyma; also observed was accumulation of plastoglobules and chloroplast swelling (Morgham et al., 1994). González et al. (2002) reported ultrastructural damage to chloroplasts in leaves of *Sorghum halepense* L. infested with *Sipha flava* Forb.; light microscopy showed increased chloroplast volume, loss of grana structures and aggregation of starch granules. Rapid and uncontrolled release of high amounts of photoactive chlorophyll molecules from thylakoid membranes can unbalance the intracellular redox state, and the photodynamic action of these pigments may induce massive oxidative damage of organelles. Some researchers have shown increased chlorophyllase activity in plants subjected to aphid colonization (Ni et al., 2002; Ciepiela et al., 2005; Sytykiewicz 2007), wounding, and the action of phytopathogens (Stangarlin and Pascholati, 2000; Kariola et al., 2005). Chlorophyllase (chlorophyll chlorophyllidohydrolase, EC 3.1.1.14) is a chloroplast membrane biocatalyst responsible for hydrolytic turnover of free chlorophyll substrates into the corresponding chlorophyllide forms (Sytykiewicz et al., 2013). Plant species have evolved a variety of mechanisms participating in chlorophyll degradation and chlorophyllase regulation (Lee et al., 2010). A wide spectrum of isozymes perform different biological functions in plants. For example, isoform 1 of chlorophyllase alleviates excessive levels of reactive oxygen species generated in leaves of the model plant *Arabidopsis thaliana* stressed with the pathogenic microorganisms *Erwinia carotovora* and *Alternaria brassicicola* (Kariola et al., 2005).

The saliva produced by wingless *R. padi* females contains a variety of biocatalysts (e.g., cellulase, polygalacturonase, pectinesterase, proteolytic enzymes) participating in cleavage of chemical compounds of the cell wall and cytoplasmic membranes, leading to their disruption; *R. padi* infestation of winter wheat cv. Bezostaja led to the occurrence of large areas of hydrolysis in the foliar mesophyll (Urbańska and Niraz, 1990). Those results are in accordance with observations from Sytykiewicz (2007) indicating that *R. padi* can feed from the sieve tubes as well as parenchyma cells in bird cherry leaves. The intracellular mode of aphid stylet penetration through plant tissues can be associated with tonoplast disintegration, with subsequent release of toxic constituents from vacuoles (Urbańska et al., 2006). The salivary sheaths of *S. avenae*, on the other hand, were found to be deposited intercellularly within the epidermis and mesophyll of winter

wheat cv. Sakva seedlings (Urbańska 2010). The absence of salivary diffusion into the cells surrounding the aphid stylet pathway and the presence of aphid saliva within vascular bundles confirm that *S. avenae* uses a typical phloem-feeding mode of stylet penetration. Matsiliza and Botha (2002) noted intercellular stylet penetration by *Sitobion yakini* (East.) in the epidermis and parenchyma of barley (*Hordeum vulgare* L.) var. Clipper, and found that it fed only on thin-walled sieve elements and not the thick-walled vascular bundles. Speculating from the sparse published data, we suggest that the more severe depletion of Chl from maize leaves infested with oligophagous *R. padi* than with monophagous *S. avenae* may be due to the difference in the aphids' modes of stylet penetration. The species-specific chemical composition of aphid salivary secretions should also differently affect the content of photosynthetic pigments in colonized plants.

In this work we confirmed a strong positive correlation between total Chl content (relative SPAD units) in seedling leaves and Chl *a+b* concentrations in acetone leaf extracts of the tested maize cultivars. We also demonstrated that the SPAD-502 chlorophyll meter reliably reflects the amount of foliar Chl in *Zea mays* plants. A significant advantage of this technique is that it can quantify Chl nondestructively in plant tissues *in situ*. Our results provide a platform for further work to estimate a broad range of photosynthetic capacity parameters and to assess the relative expression of genes involved in chlorophyll biosynthesis and catabolic turnover in tissues of aphid-susceptible and tolerant maize plants. The detailed mechanisms underlying aphid-triggered modulation of the host's metabolism remain unclear and await more study.

ACKNOWLEDGMENTS

This research was funded by the National Science Center (Poland) under grant no. N N310 733940.

REFERENCES

- ANSTEAD J, SAMUEL P, SONG N, WU C, THOMPSON GA, and GOGGIN F. 2010. Activation of ethylene-related genes in response to aphid feeding on resistant and susceptible melon and tomato plants. *Entomologia Experimentalis et Applicata* 134: 170–181.
- ARUNYANARK A, JOGLOY S, AKKASAENG C, VORASOOT N, KESMALA T, NAGESWARA RAO RC, WRIGHT GC, and PATANOTHAI A. 2008. Chlorophyll stability is an indicator of drought tolerance in peanut. *Journal of Agronomy and Crop Science* 194: 113–125.
- ASLAN MM, and UYGUN N. 2005. Aphids (Homoptera: Aphididae) of Kahramanmaraş Province, Turkey. *Turkish Journal of Zoology* 29: 201–209.

- BAK A, MARTINIÈRE A, BLANC S, and DRUCKER M. 2013. Early interactions during the encounter of plants, aphids and arboviruses. *Plant Signaling & Behavior* 8: 6, e24225. DOI: 10.4161/psb.24225.
- BOTHA A-M, LACOCK L, VAN NIEKERK C, MATSIOLOKO MT, DU PREEZ FB, LOOTS S, VENTER E, KUNERT KJ, and CULLIS CA. 2006. Is photosynthetic transcriptional regulation in *Triticum aestivum* L. cv. 'Tugela DN' a contributing factor for tolerance to *Diuraphis noxia* (Homoptera: Aphididae)? *Plant Cell Reports* 25: 41–54.
- BURD JD and ELLIOTT NC. 1996. Changes in chlorophyll a fluorescence induction kinetics in cereals infested with Russian wheat aphid (Homoptera: Aphididae). *Journal of Economic Entomology* 89: 1332–1337.
- CAROLAN JC, FITZROY CF, ASHTON PD, DOUGLAS AE, and WILKINSON TL. 2009. The proteome of the pea aphid saliva characterized by LC/MS-MS. *Proteomics* 9: 2457–2467.
- CEROVIC ZG, MASDOUMIER G, GHOZLEN NB, and LATOUCHE G. 2012. A new optical leaf-clip meter for simultaneous non-destructive assessment of leaf chlorophyll and epidermal flavonoids. *Physiologia Plantarum* 146: 251–260.
- CHEN X, VOSMAN B, VISSER RGF, VAN DER VLUGT RAA, and BROEKGAARDEN C. 2012. High throughput phenotyping for aphid resistance in large plant collections. *Plant Methods* 8: 33. DOI:10.1186/1746-4811-8-33.
- CHRZANOWSKI G, LESZCZYŃSKI B, CZERNIEWICZ P, SYTYKIEWICZ H, MATOK H, KRZYZANOWSKI R, and SEMPRUCH C. 2012. Effect of phenolic acids from black currant, sour cherry and walnut on grain aphid (*Sitobion avenae* F.) development. *Crop Protection* 35: 71–77.
- CIEPIELA AP, SYTYKIEWICZ H, and CZERNIEWICZ P. 2005. Changes in the activity of chlorophyllase and Mg-dechelataase in bird cherry leaves induced by *Rhopalosiphum padi* L. feeding. *Aphids and Other Hemipterous Insects* 11: 25–31.
- COOPER WR, DILLWITH JW, and PUTERKA GJ. 2010. Salivary proteins of Russian wheat aphid (Homoptera: Aphididae). *Environmental Entomology* 39: 223–231.
- COOPER WR, DILLWITH JW, and PUTERKA GJ. 2011. Comparisons of salivary proteins from five aphid (Homoptera: Aphididae) species. *Environmental Entomology* 40: 151–156.
- COSTE S, BARALOTO C, LEROY C, MARCON E, RENAUD A, RICHARDSON AD, ROGGY J-C, SCHIMANN H, UDDLING J, and HÉRAULT B. 2010. Assessing foliar chlorophyll contents with the SPAD-502 chlorophyll meter: a calibration test with thirteen tree species of tropical rainforest in French Guiana. *Annals of Forest Science* 67: 607. DOI: 10.1051/forest/2010020.
- COULETTE Q, COUTY A, LASUE P, RAMBAUD ZC, and AMELINE A. 2013. Colonization of the biomass energy crop miscanthus by the three aphid species, *Aphis fabae*, *Myzus persicae*, and *Rhopalosiphum padi*. *Journal of Economic Entomology* 106: 683–689.
- CZERNIEWICZ P, LESZCZYŃSKI B, CHRZANOWSKI G, SEMPRUCH C, and SYTYKIEWICZ H. 2011a. Effects of host plant phenolics on spring migration of bird cherry-oat aphid (*Rhopalosiphum padi* L.). *Allelopathy Journal* 27: 309–316.
- CZERNIEWICZ P, SYTYKIEWICZ H, LESZCZYŃSKI B, CHRZANOWSKI G, SEMPRUCH C, and SPRAWKA I. 2011b. Wpływ zawartości roślinnych związków azotowych na wiosenne migracje mszycy czeremchowo-zbożowej (*Rhopalosiphum padi* L.). *Progress in Plant Protection* 51: 409–412.
- DAHMARDEH M, GHANBARI A, SYAHSAR BA, and RAMRODI M. 2010. The role of intercropping maize (*Zea mays* L.) and cowpea (*Vigna unguiculata* L.) on yield and soil chemical properties. *African Journal of Agricultural Research* 5: 631–636.
- DIAZ-MONTANO J, REESE JC, SCHAPAUGH WT, and CAMPBELL LR. 2007. Chlorophyll loss caused by soybean aphid (Homoptera: Aphididae) feeding on soybean. *Journal of Economic Entomology* 100: 1657–1662.
- FERRIOL I, RUBIO L, PÉREZ-PANADÉS J, CARBONELL EA, DAVINO S, and BELLUURE B. 2013. Transmissibility of *Broad bean wilt virus 1* by aphids: influence of virus accumulation in plants, virus genotype and aphid species. *Annals of Applied Biology* 162: 71–79.
- FILIPOVIĆ A, POLJAK M, and ŠKOBIĆ D. 2013. Response of chlorophyll a, SPAD values and chlorophyll fluorescence parameters in leaves of apricot affected some abiotic factors. *Journal of Food Science and Engineering* 3: 19–24.
- FORNALÉ S, CAPELLADES M, ENCINA A, WANG K, IRAR S, LAPIERRE C, RUEL K, JOSELEAU J-P, BERENGER J, PUIGDOMÈNECH P, RIGAU J, and CAPARRÓS-RUIZ D. 2012. Altered lignin biosynthesis improves cellulosic bioethanol production in transgenic maize plants down-regulated for cinnamyl alcohol dehydrogenase. *Molecular Plant* 5: 817–830.
- GAO S, and LIU D. 2013. Differential performance of *Sitobion avenae* (Homoptera: Aphididae) clones from wheat and barley with implications for its management through alternative cultural practices. *Journal of Economic Entomology* 106: 1294–1301.
- GHOZIZADEH A, AMIN MSM, ANUAR AR, AIMRUN W, and SABERIOON MM. 2011. Temporal variability of SPAD chlorophyll meter readings and its relationship to total nitrogen in leaves within a Malaysian Paddy Field. *Australian Journal of Basic and Applied Sciences* 5: 236–245.
- GHOSH M, SWAIN DK, JHA MK, and TEWARI VK. 2013. Precision nitrogen management using chlorophyll meter for improving growth, productivity and N use efficiency of rice in subtropical climate. *Journal of Agricultural Science* 5: 253–266.
- GOLAWSKA S, KRZYŻANOWSKI R, and ŁUKASIK I. 2010. Relationship between aphid infestation and chlorophyll content in Fabaceae species. *Acta Biologica Cracoviensia Series Botanica* 52: 76–80.
- GOLAWSKA S, ŁUKASIK I, WÓJCICKA A, and SYTYKIEWICZ H. 2012. Relationship between saponin content in alfalfa and aphid development. *Acta Biologica Cracoviensia Series Botanica* 54: 1–8.
- GONZÁLES WL, RAMÍREZ CC, OLEA N, and NIEMEYER HM. 2002. Host plant changes produced by the aphid *Sipha flava*: consequences for aphid feeding behaviour and growth. *Entomologia Experimentalis et Applicata* 103: 107–113.
- HAILE FJ, HIGLEY LG, NI X, and QUISENBERRY SS. 1999. Physiological and growth tolerance in wheat to Russian wheat aphid (Homoptera: Aphididae) injury. *Environmental Entomology* 28: 787–794.
- HAWKINS TS, GARDINER ES, and COMER GS. 2009. Modeling the relationship between extractable chlorophyll and SPAD-

- 502 readings for endangered plant species research. *Journal for Nature Conservation* 17: 123–127.
- HENG-MOSS TM, NI X, MACEDO T, MARKWELL JP, BAXENDALE FP, QUISENBERRY SS, and TOLMAY V. 2003. Comparison of chlorophyll and carotenoid concentrations among Russian wheat aphid (Homoptera: Aphididae)-infested wheat isolines. *Journal of Economic Entomology* 96: 475–481.
- HERRMANN A. 2013. Biogas production from maize: current state, challenges and prospects. 2. Agronomic and environmental aspects. *BioEnergy Research* 6: 372–387.
- HIGASHI CH, and BRESSAN A. 2012. Influence of a propagative plant virus on the fitness and wing dimorphism of infected and exposed insect vectors. *Oecologia* 172: 847–856.
- IHL M, MARTIN AS, and BIFANI V. 2000. Chlorophyllase and quality of 'Granny Smith' apples during storage. *Gartenbauwissenschaft* 65: 266–271.
- JINWEN L, JINGPING Y, PINPIN F, JUNLAN S, DONGSHENG L, CHANGSHUI G, and WENYUE C. 2009. Responses of rice leaf thickness, SPAD readings and chlorophyll a/b ratios to different nitrogen supply rates in paddy field. *Field Crops Research* 114: 426–432.
- KARIOLA T, BRANDER G, LI J, and PALVA ET. 2005. Chlorophyllase 1, a damage control enzyme, affects the balance between defense pathways in plants. *Plant Cell* 17: 282–294.
- KHATOON T, HUSSAIN K, MAJEED A, NAWAZ K, and NISAR MF. 2010. Morphological variations in maize (*Zea mays* L.) under different levels of NaCl at germinating stage. *World Applied Sciences Journal* 8: 1294–1297.
- LEE GC, CHEPYSHKO H, CHEN HH, CHU C, CHOU YF, AKOH CC, and SHAW JF. 2010. Genes and biochemical characterization of three novel chlorophyllase isozymes from *Brassica oleracea*. *Journal of Agricultural and Food Chemistry* 58: 8651–8657.
- LEWIS MF, LORENZANA RE, JUNG HJG, and BERNARDO R. 2010. Potential for simultaneous improvement of corn grain yield and stover quality for cellulosic ethanol. *Crop Science* 50: 516–523.
- LI F, KONG L, LIU Y, WANG H, CHEN L, and PENG J. 2013. Response of wheat germplasm to infestation of English grain aphid (Hemiptera: Aphididae). *Journal of Economic Entomology* 106: 1473–1478.
- LING Q, HUANG W, and JARVIS P. 2011. Use of a SPAD-502 meter to measure leaf chlorophyll concentration in *Arabidopsis thaliana*. *Photosynthesis Research* 107: 209–214.
- LIU ZA, YANG JP, and YANG ZC. 2012. Using a chlorophyll meter to estimate tea leaf chlorophyll and nitrogen contents. *Journal of Soil Science and Plant Nutrition* 12: 339–348.
- ŁUKASIK I. 2009. Effect of host plant alternation on some adaptive enzymes of the bird cherry-oat aphid *Rhopalosiphum padi* (L.). *Journal of Pest Science* 82: 203–209.
- ŁUKASIK I, GOŁAWSKA S, and WÓJCICKA A. 2012. Effect of cereal aphid infestation on ascorbate content and ascorbate peroxidase activity in triticale. *Polish Journal of Environmental Studies* 21: 1937–1941.
- MACEDO TB, BASTOS CS, HIGLEY LG, OSTLIE KR, and MADHAVAN S. 2003. Photosynthetic responses of soybean to soybean aphid (Homoptera: Aphididae) injury. *Journal of Economic Entomology* 96: 188–193.
- MAKKOUK KM, and KUMARI SG. 2009. Epidemiology and integrated management of persistently transmitted aphid-borne viruses of legume and cereal crops in West Asia and North Africa. *Virus Research* 141: 209–218.
- MARENCO RA, ANTEZANA-VERA SA, and NASCIMENTO HCS. 2009. Relationship between specific leaf area, leaf thickness, leaf water content and SPAD-502 readings in six Amazonian tree species. *Photosynthetica* 47: 184–190.
- MATSILIZA B, and BOTHA CEJ. 2002. Aphid (*Sitobion yakini*) investigation suggests thin-walled sieve tubes in barley (*Hordeum vulgare*) to be more functional than thick-walled sieve tubes. *Physiologia Plantarum* 115: 137–143.
- MIELKE M, SCHAFFER B, and LI C. 2010. Use of a SPAD meter to estimate chlorophyll content in *Eugenia uniflora* L. leaves as affected by contrasting light environments and soil flooding. *Photosynthetica* 48: 332–338.
- MORGHAM AT, RICHARDSON PE, CAMPBELL RK, BURD JD, EIKENBARY RD, and SUMNER LC. 1994. Ultrastructural responses of resistant and susceptible wheat to infestation by greenbug biotype E (Homoptera: Aphididae). *Annals of the Entomological Society of America* 87: 908–917.
- NETO C, CARRANCA C, CLEMENTE J, and DE VARENNES A. 2011. Assessing the nitrogen nutritional status of young non-bearing 'Rocha' pear trees grown in a Mediterranean region by using a chlorophyll meter. *Journal of Plant Nutrition* 34: 627–639.
- NI X, QUISENBERRY SS, HENG-MOSS T, MARKWELL J, HIGLEY L, BAXENDALE F, SARATH G, and KLUCAS R. 2002. Dynamic change in photosynthetic pigments and chlorophyll degradation elicited by cereal aphid feeding. *Entomologia Experimentalis et Applicata* 105: 43–53.
- NICHOLSON SJ, HARTSON SD, and PUTERKA GJ. 2012. Proteomic analysis of secreted saliva from Russian Wheat Aphid (*Diuraphis noxia* Kurd.) biotypes that differ in virulence to wheat. *Journal of Proteomics* 75: 2252–2268.
- PIEŃKOSZ A, LESZCZYŃSKI B, and WARZECHA R. 2005. Podatność kukurydzy na mszycę zbożowe. *Progress in Plant Protection* 45: 989–992.
- RAO SA, CAROLAN JC, and WILKINSON TL. 2013. Proteomic profiling of cereal aphid saliva reveals both ubiquitous and adaptive secreted proteins. *PLoS One* 8: e57413. DOI: 10.1371/journal.pone.0057413.
- REDDY KR, HENRY WB, SEEPAL R, LOKHANDE S, GAJANAYAKE B, and BRAND D. 2013. Exogenous application of glycinebetaine facilitates maize (*Zea mays* L.) growth under water deficit conditions. *American Journal of Experimental Agriculture* 3: 1–13.
- RIEDEL WE, and BLACKMER TM. 1999. Leaf reflectance spectra of cereal aphid damaged wheat. *Crop Science* 39: 1835–1840.
- RIEDEL WE, BECKENDORF EA, and CATANGUI MA. 2013. Soybean aphid injury effects on shoot N components in *Glycine max*. *Crop Science* 53: 232–239.
- ROSTAMI M, KOOCHEKI AR, NASIRI MAHALLATI M, and KAFI M. 2008. Evaluation of chlorophyll meter (SPAD) data for prediction of nitrogen status in corn (*Zea mays* L.). *American-Eurasian Journal of Agricultural and Environmental Science* 3: 79–85.

- RUIZ-ESPINOZA F, MURILLO-AMADOR B, GARCIA-HERNANDEZ LJ, FENECH-LARIOS L, RUEDA-PUENTE OE, TROYO-DIEGUEZ E, KAYA C, and BELTRAN-MORALES A. 2010. Field evaluation of the relationship between chlorophyll content in basil leaves and a portable chlorophyll meter (SPAD-502) readings. *Journal of Plant Nutrition* 33: 423–438.
- SEMNČENKO VV, MOJOVIĆ LV, ĐUKIĆ-VUKOVIĆ AP, RADOSAVLJEVIĆ MM, TERZIĆ DR, and MILAŠINOVIĆ ŠEREMEŠIĆ MS. 2013. Suitability of some selected maize hybrids from Serbia for the production of bioethanol and dried distillers' grains with solubles. *Journal of the Science of Food and Agriculture* 93: 811–818.
- SEMPRUCH C, LESZCZYŃSKI B, WÓJCICKA A, MAKOSZ M, MATOK H, and CHRZANOWSKI G. 2010. Changes in activity of lysine decarboxylase within winter triticale in response to grain aphid feeding. *Acta Biologica Hungarica* 61: 512–515.
- SEMPRUCH C, MICHALAK A, and LESZCZYŃSKI B. 2011. Effect of grain aphid (*Sitobion avenae* Fabricius, 1775) feeding on content of free amino acids within selected parts of triticale plants. *Aphids and Other Hemipterous Insects* 17: 137–144.
- SEMPRUCH C, HORBOWICZ M, KOSSON R, and LESZCZYŃSKI B. 2012. Biochemical interactions between triticale (Triticosecale; Poaceae) amines and bird cherry-oat aphid (*Rhopalosiphum padi*; Aphididae). *Biochemical Systematics and Ecology* 40: 162–168.
- SHRESTHA S, BRUECK H, and ASCH F. 2012. Chlorophyll index, photochemical reflectance index and chlorophyll fluorescence measurements of rice leaves supplied with different N levels. *Journal of Photochemistry and Photobiology B: Biology* 113: 7–13.
- SMITH CM, BOYKO E, and STARKEY S. 2005. Differential expression of genes in wheat, *Triticum aestivum* L. controlling resistance to the Russian wheat aphid, *Diuraphis noxia* (Mordvilko). *IOBC-WPRS Bulletin*, 28: 11–20.
- SPRAWKA I, GOŁAWSKA S, CZERNIEWICZ P, and SYTYKIEWICZ H. 2011. Insecticidal action of phytohemagglutinin (PHA) against the grain aphid, *Sitobion avenae*. *Pesticide Biochemistry and Physiology* 100: 64–69.
- SPRAWKA I, GOŁAWSKA S, GOŁAWSKI A, CZERNIEWICZ P, and SYTYKIEWICZ H. 2012. Antimetabolic effect of phytohemagglutinin to the grain aphid *Sitobion avenae* Fabricius. *Acta Biologica Hungarica* 63: 342–353.
- SPRAWKA I, GOŁAWSKA S, GOŁAWSKI A, CHRZANOWSKI G, CZERNIEWICZ P, and SYTYKIEWICZ H. 2013a. Entomotoxic action of jackbean lectin (Con A) in bird cherry-oat aphid through the effect on insect enzymes. *Journal of Plant Interactions*. DOI 10.1080/17429145.2013.848947
- SPRAWKA I, GOŁAWSKA S, PARZYCH T, GOŁAWSKI A, CZERNIEWICZ P, and SYTYKIEWICZ H. 2013b. Induction of apoptosis in the grain aphid *Sitobion avenae* (Hemiptera:Aphididae) under the influence of phytohaemagglutinin PHA. *Applied Entomology and Zoology*. DOI: 10.1007/s13355-013-0214-2
- STANGARLIN JR, and PASCHOLATI SF. 2000. Activities of ribulose-1,5-bisphosphate carboxylase-oxygenase (Rubisco), chlorophyllase, β -1,3-glucanase and chitinase and chlorophyll content in bean cultivars (*Phaseolus vulgaris*) infected with *Uromyces appendiculatus*. *Summa Phytopathologica* 26: 34–42.
- STEWART LR, BOUCHARD R, REDINBAUGH MG, and MEULIA T. 2012. Complete sequence and development of a full-length infectious clone of an Ohio isolate of maize dwarf mosaic virus (MDMV). *Virus Research* 165: 219–224.
- STOETZEL MB, and MILLER GL. 2001. Aerial feeding aphids of corn in the United States with reference to the root-feeding *Aphis maidiradicis* (Homoptera: Aphididae). *Florida Entomologist* 84: 83–98.
- STRAŻYŃSKI P. 2008. Aphid fauna (Hemiptera, Aphidoidea) on maize crops in Wielkopolska – species composition and increase in number. *Aphids and Other Homopterous Insects* 14: 123–128.
- SVOBODOVÁ E, TRNKA M, DUBROVSKÝ M, SEMERÁDOVÁ D, EITZINGER J, STIPÁNEK P, and ZALUD Z. 2013. Determination of areas with the most significant shift in pests' persistence in Europe under climate change. *Pest Management Science* DOI: 10.1002/ps.3622. [Epub ahead of print].
- SWAIN DK, and SANDIP SJ. 2010. Development of SPAD values of medium- and long-duration rice variety for site-specific nitrogen management. *Journal of Agronomy* 9: 38–44.
- SYTYKIEWICZ H. 2007. Biochemiczne i anatomiczne aspekty żerowania mszycy czeremchowo-zbożowej (*Rhopalosiphum padi* L.) na żywicielu pierwotnym. Ph.D. dissertation, Akademia Podlaska, Wydział Rolniczy, Siedlce.
- SYTYKIEWICZ H, GOŁAWSKA S, and CHRZANOWSKI G. 2011a. Effect of the bird cherry-oat aphid, *Rhopalosiphum padi* L. feeding on phytochemical responses within the bird cherry. *Polish Journal of Ecology* 59: 329–338.
- SYTYKIEWICZ H, CZERNIEWICZ P, SPRAWKA I, GOŁAWSKA S, CHRZANOWSKI G, and LESZCZYŃSKI B. 2011b. Induced proteolysis within the bird cherry leaves evoked by *Rhopalosiphum padi* L. (Hemiptera, Aphidoidea). *Acta Biologica Hungarica* 62: 316–327.
- SYTYKIEWICZ H, SPRAWKA I, CZERNIEWICZ P, SEMPRUCH C, LESZCZYŃSKI B, and SIKORA M. 2013. Biochemical characterisation of chlorophyllase from leaves of selected *Prunus* species – A comparative study. *Acta Biochimica Polonica* 60: 457–465.
- UDDLING J, GELANG-ALFREDSSON J, PIKKI K, and PLEJEL H. 2007. Evaluating the relationship between leaf chlorophyll concentration and SPAD-502 chlorophyll meter readings. *Photosynthesis Research* 91: 37–46.
- URBAŃSKA A. 2010. Histochemical analysis of aphid saliva in plant tissue. *Electronic Journal of Polish Agricultural Universities, Biology* 13: #26. (<http://www.ejpau.media.pl/volume13/issue4/art-26.html>)
- URBAŃSKA A, and NIRAZ S. 1990. Anatomiczne i biochemiczne aspekty żerowania mszyc zbożowych. *Zeszyty Problemowe Postępów Nauk Rolniczych* 392: 201–213.
- URBAŃSKA A, LESZCZYŃSKI B, and MATOK H. 2006. How does gramine affect probing behaviour of grain aphid. *Electronic Journal of Polish Agricultural Universities, Biology* 9: #2 (<http://www.ejpau.media.pl/volume9/issue1/art-02.html>).
- ZIELIŃSKA L, TRZMIEL K, and JEŻEWSKA M. 2012. Ultrastructural changes in maize leaf cells infected with maize dwarf mosaic virus and sugarcane mosaic virus. *Acta Biologica Cracoviensia Series Botanica* 54: 97–104.