

DIGENEAN FAUNA OF THE GREAT CORMORANT *PHALACROCORAX CARBO SINENSIS* (BLUMENBACH, 1798) IN THE BRACKISH WATERS OF THE VISTULA LAGOON AND THE GULF OF GDAŃSK (POLAND)

GERARD KANAREK, JILJI SITKO*, LESZEK ROLBIECKI AND JERZY ROKICKI

Division of Invertebrate Zoology, University of Gdańsk, al. Marsz. Piłsudskiego 46, 81-378 Gdynia, Poland, E-mail: kanarek@sat.ocean.univ.gda.pl; *Moravian Ornithological Station, Bezručova 10, 751 52 Přerov, Czech Republic

ABSTRACT. The great cormorant *Phalacrocorax carbo sinensis* (Blumenbach, 1798) is one of the most important birds in the Vistula Lagoon and the Gulf of Gdańsk area. As a typical piscivore, the bird plays a significant part in life cycles of parasites in aquatic ecosystems. Within January 2000-June 2001, a complete helminthological examination was performed on great cormorants collected in the nesting area at Kały Rybackie on the Vistula Spit (80 specimens) and in the Vistula Lagoon (10 specimens). The infection prevalence, mean intensity, and intensity range were 92.2%, 376.5, and 1-4524, respectively. The presence of 9 digenean species (*Paryphostomum radiatum* (Dujardin, 1845), *Petasiger exaeretus* Dietz, 1909, *P. phalacrocoracis* (Yamaguti, 1939), *Mesorchis pseudoechinatus* (Olsson, 1876), *Metorchis xanthosomus* (Creplin, 1846), *Cryptocotyle concavum* (Creplin, 1825), *Hysteromorpha triloba* (Rudolphi, 1819), *Tylodelphys clavata* (Nordmann, 1832), and *Holostephanus dubinini* Vojtek et Vojtkova, 1968) was recorded, *P. phalacrocoracis* being the most common parasite (prevalence 92.2%, mean intensity 323.8). *Tylodelphys clavata* proved a parasite new for the great cormorant moreover the records of *P. phalacrocoracis*, *H. triloba*, and *H. dubinini* are the first in Poland, while *C. concavum* and *M. pseudoechinatus* were for the first time recorded in the great cormorant in Poland.

Key words: brackish waters, Digenea, great cormorant, *Phalacrocorax carbo sinensis*, Poland.

INTRODUCTION

Of the five subspecies of the cormorant *Phalacrocorax carbo* (Linnaeus, 1758), the great cormorant *Phalacrocorax carbo sinensis* (Blumenbach, 1798) occurs in central Europe, including Poland (Cramp and Simmons 1977, Tomiałojć 1990, Przybysz 1997). Great cormorants are large piscivorous birds growing to the length of 100 cm and attaining the wing span of 160 cm. Great cormorants inhabiting central Europe migrate regularly to overwinter in the Mediterranean, Bay of Biscay, and North Sea areas (Cramp and Simmons 1977, Przybysz 1997). In the 1980s, the *Ph. carbo sinensis* population grew rapidly in abundance throughout Europe, thereby rendering the great cormorant a common species, and even a dominant in numerous avian faunas. Such population growth resulted in a number of adverse effects,

including those incurred by fisheries (fish consumption) and forestry (damage to trees caused by strongly caustic faeces) (Dobrowolski and Dejtrowski 1997, Przybysz 1997, Szramka 2000).

The knowledge of the great cormorant parasitic fauna in Poland is fragmentary only. Some publications dealt with cestodes (Rybicka 1958, Korpaczewska 1963a, b), nematodes (Okulewicz 1989, Okulewicz and Rokicki 1998, Żuchowska 2000), and mallophagans (Złotorzycka 1990). The great cormorant trematode fauna was studied solely by Sulgostowska (1958, 1960a, b, 1963) who examined a few birds only.

This paper deals, for the first time in Poland, with digeneans parasitising the great cormorants feeding in brackish waters of the Vistula Lagoon and the Gulf of Gdańsk.

MATERIALS AND METHODS

Within January 2000-June 2001, a total of 90 great cormorants including 78 juveniles at their first year of life and 12 adult, sexually mature birds were examined. As the gonads were poorly developed in the juveniles, only the adults were sexed. The juveniles were collected during breeding seasons (mid-May until late June) in 2000 (52 birds) and 2001 (26 birds) from the nesting colony at Kały Rybackie on the Vistula Spit (Fig. 1). They were those individuals unable to fly that fell out of nests. The adults examined had been found dead on the Vistula Spit or in




Fig. 1. Study area; thick circle – breeding colony at Kały Rybackie

fishing nets set up in the Vistula Lagoon (1 bird in January 2000, 1 bird in March 2000 and 1 in March 2001, 4 birds in April 2000, and 5 birds in April 2001).

The birds were subjected to a complete helminthological examination, carried out according to the generally adopted protocol (Dubinina 1971). The intestine, stomach, body cavity, mouth cavity, eyelids, liver, gall bladder, kidneys, ovaries, bursa Fabricii (at juveniles), oesophagus, tracheae, bronchia, heart, tarsi, and the subcutaneous tissue were examined. The parasites were removed from the intestine by decantation, the supernatant being additionally filtered through the miller's gauze. The parasites were fixed in 70% ethanol. Some of the parasites were stained in borax carmine, dehydrated in glacial acetic acid, and cleared in benzyl alcohol. Permanent mounts of some parasites were made by embedding them in Canada balsam; the remaining ones were left in the clearing fluid or in alcohol.

RESULTS

The total digenean prevalence in the great cormorant examined was 92.2%, the mean intensity and the intensity range amounting to 376.5 and 1-4524 individuals, respectively. The intestines and gall bladders (the latter containing only *Metorchis xanthosomus*) of 83 birds were found to contain 9 trematode species which were dominated by *Petasiger phalacrocoracis* (92.2%, 323.8; 1-4500) (Table 1).

The year 2000 materials showed the presence of 8 digenean species (*Paryphostomum radiatum*, *Petasiger exaeretus*, *P. phalacrocoracis*, *Metorchis xanthosomus*, *Cryptocotyle concavum*, *Hysteromorpha triloba*, *Tylodelphys clavata*, *Holostephanus dubinini*), the infection affecting 93.1% of the birds and the infection mean intensity amounting to 75.6. In 2001 the survey yielded 7 species (lack of *C. concavum* and *T. clavata*, the presence of *Mesorchis pseudoechinatus* not recorded in 2000), the infection prevalence and mean intensity being 90.6% and 335.6, respectively.

The infection level was different in the various ages categories: the juveniles hosted 8 digenean species that affected 93.6% of the birds with a mean intensity of 406.5, while the adults hosted 7 species affecting 83.3% of the birds with a mean intensity of 157.5 (Table 1).

DISCUSSION

The great cormorant (*Phalacrocorax carbo sinensis*) is one of the most common birds in the area of the Vistula Lagoon and the Gulf of Gdańsk. It nests at the Vistula Spit's base, forming a breeding colony at Kąty Rybackie; the colony is the largest in Poland and one of the largest in Europe. About 6000 breeding pairs are recorded there each year; the number of birds counted at the end of the breeding season may even exceed 25000 (Stempniewicz 1997).

Table 1. The frequency of *Digenea* in cormorants of different age, feeding on the Vistula Lagoon and Gulf of Gdańsk

| Parasites | No. of parasites [ind.] | | Prevalence [%] | | Mean int. [ind] | | Range of int. [ind.] | |
|----------------------------------|----------------------------|--------|-------------------|--------|--------------------|--------|-------------------------|--------|
| | Juveniles | Adults | Juveniles | Adults | Juveniles | Adults | Juveniles | Adults |
| <i>Paryphostomum radiatum</i> | 323 | 326 | 53.8 | 83.3 | 46.6 | 46.6 | 1-39 | 1-122 |
| <i>Petasiger exaeretis</i> | 4 | 8 | 3.8 | 16.6 | 4 | 4 | 1-2 | 3-5 |
| <i>P. phalacrocoracis</i> | 26020 | 852 | 93.6 | 83.3 | 85.2 | 85.2 | 1-4500 | 1-452 |
| <i>Mesorchis pseudoechinatus</i> | 2 | 0 | 2.6 | 0 | 0 | 0 | 1 | 0 |
| <i>Metorchis xanthosomus</i> | 565 | 11 | 37.2 | 33.3 | 2.8 | 2.8 | 1-128 | 1-5 |
| <i>Cryptocotyle concavum</i> | 392 | 0 | 3.83 | 3.33 | 0 | 0 | 9-68 | 0 |
| <i>Hysteromorpha triloba</i> | 648 | 376 | 53.8 | 50 | 62.7 | 62.7 | 1-149 | 1-255 |
| <i>Tylodelphys clavata</i> | 0 | 2 | 0 | 8.33 | 2 | 2 | 0 | 32 |
| <i>Holostephanus dubinini</i> | 2001 | 7 | 74.4 | 16.6 | 3.5 | 3.5 | 1-239 | 2-5 |
| Total | 29673 | 1579 | 93.6 | 83.3 | 157.7 | 157.7 | 1-4524 | 1-522 |
| | | | | | 376.5 | 376.5 | | 1-4524 |
| | | | | | 406.5 | 406.5 | | 1-522 |
| | | | | | 92.2 | 92.2 | | 1-522 |
| | | | | | 83.3 | 83.3 | | 1-522 |
| | | | | | 93.6 | 93.6 | | 1-522 |
| | | | | | 31252 | 31252 | | 1-4524 |
| | | | | | 1579 | 1579 | | 1-4524 |
| | | | | | 83.3 | 83.3 | | 1-4524 |
| | | | | | 93.6 | 93.6 | | 1-4524 |
| | | | | | 31252 | 31252 | | 1-4524 |
| | | | | | 1579 | 1579 | | 1-4524 |
| | | | | | 83.3 | 83.3 | | 1-4524 |
| | | | | | 93.6 | 93.6 | | 1-4524 |
| | | | | | 31252 | 31252 | | 1-4524 |

The number of digenean species found in this study (9) is fairly high. For comparison, 9 species were also recorded in Ukraine (Smogorzhevskaya 1976), 12 taxa (11 species and 1 taxon identified to the genus level) being recorded in the former East Germany (Reimer 1969) and 11 taxa being found in Moravia (Czech Republic) (Našincová et al. 1993).

Tylodelphys clavata was recorded for the first time in the great cormorant. As, however, the parasite was represented by two immature individuals, and in view of the digenean's larval stages being very common in the intermediate hosts, i.e., fish in the Gulf of Gdańsk and in the Vistula Lagoon (Rolbiecki et al. 1999, Rolbiecki 2003), it can be presumed that the infection was accidental and that the parasites would perish in the bird without achieving their maturity. On the other hand, *Petasiger phalacrocoracis*, *Hysteromorpha triloba*, and *Holostephanus dubinini* proved new for the fauna of Poland, while the records of *Cryptocotyle concavum* and *Mesorchis pseudoechinatus* were the first in the great cormorant of Poland. Apart from *Paryphostomum radiatum*, *Petasiger exaeretus*, *P. phalacrocoracis*, and *Holostephanus dubinini*, the parasites specific for the great cormorant, and the grebe-specific *Tylodelphys clavata*, the parasites identified affect a wide spectrum of hosts, including birds belonging to different orders (Yamaguti 1958, Bykhovskaya-Pavlovskaya 1962, Sonin 1985, 1986, Sulgostowska and Czaplińska 1987). Among the parasites found in this study, *Mesorchis pseudoechinatus* and *Cryptocotyle concavum* had been earlier found in the area, the former in the Laridae and the latter in the Laridae and Anatidae (only in the Gulf of Gdańsk) (Markowski 1933, Malczewski 1964, Sulgostowska and Grytner-Zięcina 1974, Grytner-Zięcina and Sulgostowska 1978). In addition, in the Gulf of Gdańsk *Clangula hyemalis* (Anatidae) had been found to host digeneans (not identified to the species level) of the genera *Tylodelphys*, *Paryphostomum*, and *Petasiger* (Sulgostowska and Grytner-Zięcina 1974). Prior to this study, the great cormorant in Poland (Masurian lakes) had been found to host as few as 3 digenean species: *Paryphostomum radiatum*, *Petasiger exaeretus*, and *Metorchis xanthosomus* (Sulgostowska 1958, 1960a, b, 1963). It has to be remembered, however, that *P. exaeretus* is frequently mistaken for *P. phalacrocoracis*, regarded as a common species (Našincová et al. 1993, 1994).

The parasites identified consist of species typical of both marine (*Mesorchis pseudoechinatus*, *Cryptocotyle concavum*) and freshwater (*Paryphostomum radiatum*, *Petasiger exaeretus*, *P. phalacrocoracis*, *Metorchis xanthosomus*, *Hysteromorpha triloba*, *Tylodelphys clavata*, *Holostephanus dubinini*) ecosystems. It is worth emphasising that the infection is of local origin because all the species, except *Tylodelphys clavata*, were recorded in the juvenile cormorants, fed by their parents with fish caught in the vicinity of the colony. It is noteworthy that life cycles of all the digenean species found involve metacercariae (larvae) dwelling in fish; the birds become infected *via* the trophic chain. The absence of larger, typically

freshwater reservoirs in the vicinity of the Kąty Rybackie colony is indicative of the fact that the digeneans find conditions favourable for their development also in the brackish habitats of the Vistula Lagoon and the Gulf of Gdańsk of average salinities amounting to 2.4 and 7.5‰, respectively (Łomniewski 1958, Majewski 1990).

The relationship between the parasites' presence and the birds' age may be an interesting problem. The present study, involving 78 juveniles and as few as 12 adult great cormorants, does not make it possible to detect any unambiguous trend in age-dependent infection effects. However, the infections were heavier in the juveniles, both generally and with respect to most of the parasitic species (Table 1). This finding is in agreement with observations of other authors (Bykhovskaya-Pavlovskaya 1962, Smogorzhevskaya 1976, Sitko 1993) who noted that young birds were frequently more heavily infected than adults. The authors quoted explain this by referring to a higher food consumption by the juveniles fed by the parents who take up less food themselves. Obviously, this explanation applies to the period of nesting only. When that stage is over, the young birds join their parents in getting food on their own. In addition, at the beginning of their independent life the young, less experienced cormorants, are able to catch less fish than the adults.

REFERENCES

- Bykhovskaya-Pavlovskaya I.E. 1962. Trematody ptic fauny SSSR. Izd. AN SSSR, Moskva-Leningrad.
- Cramp S., Simmons K.E.L. 1977. Handbook of the Middle Asia and North Africa birds. The birds of the western Palearctic, Vol. I. Oxford University Press.
- Dobrowolski K., Dejtrowski R. 1997. Conflict between fishermen and cormorants *Phalacrocorax carbo* in Poland. *Ekologia Polska* 45: 279-283.
- Dubinina M.N. 1971. Parasitologicheskoe issledovanije ptic. Izd. Nauka, Leningrad.
- Grytner-Zięcina B., Sulgostowska T. 1978. Trematodes of *Oidemia fusca* (L.), *Oidemia nigra* (L.), and *Somateria mollissima* (L.), from the Baltic Coast. *Acta Parasitologica Polonica* 25: 121-128.
- Korpaczewska W. 1963a. Tapeworms of aquatic birds in some Mazurian lakes. *Acta Parasitologica Polonica* 11: 315-336.
- Korpaczewska W. 1963b. Formation of a population structure and cestode complexes in water birds. *Acta Parasitologica Polonica* 11: 337-344.
- Łomniewski K. 1958. Zalew Wiślany. Prace geograficzne 15. PAN, Warszawa.
- Majewski A. 1990. Zatoka Gdańska. Wydawnictwo Geologiczne, Warszawa.
- Malczewski A. 1964. Trematoda mew z rodzaju *Larus* L. znad Zalewu Wiślanego. *Wiadomości Parazytologiczne* 10: 563-564.
- Markowski S. 1933. Contributions a l'étude de la faune helminthologique de la presequ'île de Hel. *Fragmenta Faunistica Musei Zoologici Polonici* 2: 107-111.
- Našincová V., Moravec F., Scholtz T. 1993. Trematodes of the common cormorant (*Phalacrocorax carbo*) in Czech Republic. *Acta Societatis Zoologicae Bohemicae* 57: 31-46.
- Našincová V., Scholtz T., Moravec F. 1994. Redescription of *Petasiger exaeretus* Dietz, 1909 and *P. phalacrocoracis* (Yamaguti, 1939) (Trematoda: Echinostomatidae), parasites of cormorants. *Systematic Parasitology* 27: 139-147.

- Okulewicz A. 1989. Redeskrpcja *Capillaria carbonis* (Rudolphi, 1819) (Capillariidae), na podstawie materiału zebranego z kormorana czarnego (*Phalacrocorax carbo*). *Wiadomości Parazytologiczne* 35: 577-583.
- Okulewicz A., Rokicki J. 1998. *Contracaecum micropapillatum* (Stossich, 1819) (Nematoda) – new species to the parasitofauna in Poland. A special Symposium Arranged on Behalf of the Baltic Society for Parasitology and Scandinavian Society for Parasitology, Ecology and Bird-Parasite Interactions, Vilnius, Lithuania 25-28 June, 1998: 61.
- Przybysz J. 1997. Kormoran. Wydawnictwo Lubuskiego Klubu Przyrodników, Świebodzin.
- Reimer L. 1969. Helminthen von Kormoranen von Brutkolonien der Deutschen Demokratischen Republik. *Wissenschaftliche Zeitschrift der Ernst-Moritz-Arndt-Universität Greifswald. Mathematisch-naturwissenschaftliche Reihe* 18: 129-135.
- Rolbiecki L. 2003. Diversity of the parasite fauna of cyprinid (Cyprinidae) and percid (Percidae) fishes in the Vistula Lagoon, Poland. *Wiadomości Parazytologiczne* 49: 125-164
- Rolbiecki L., Rokicki J., Morozińska-Gogol J., Chibani M. 1999. Larval stages of helminths in fish from the Vistula Lagon and the Gulf of Gdańsk in relation to birds occurrence. *Bulletin of the Sea Fisheries Institute* 147: 51-60
- Rybicka K. 1958. Tasiemce ptaków (excl. Anseriformes) jeziora Drużno. *Acta Parasitologica Polonica* 6: 143-178.
- Sitko J. 1993. Ecological relations of trematodes infecting lariform birds in Czech Republic. *Acta scientarium naturalium Academiae scientarium Bohemicae*, Brno 27: 1-93.
- Smogorzhevskaya L.A. 1976. Gel'minty vodoplavayushchikh i bolotnykh ptic fauny Ukrainy. Naukova Dumka, Kiev.
- Sonin M. D. 1985. Opredelitel' trematod ryboyadnykh ptic Palearktiki (brakhilajmidy, klinostomatidy, ciklocelidy, fasciolidy, notokotilidy, plagiorkhidy, shistosomatidy). Izd. Nauka, Moskva.
- Sonin M. D. 1986. Opredelitel' trematod ryboyadnykh ptic Palearktiki (opistorkhidy, renikolidy, strigeidy). Izd. Nauka, Moskva.
- Stempniewicz L. 1997. Ryby, kormorany i las. W: Ocena presji kormorana czarnego *Phalacrocorax carbo sinensis* na ichtiofaunę Zalewu Wiślanego (Red. L. Stempniewicz.). Raport 3: 1-12.
- Sulgostowska T. 1958. Flukes of birds of Drużno Lake. *Acta Parasitologica Polonica* 6: 111-141.
- Sulgostowska T. 1960a. Intestinal trematodes of birds of mesotrophic lakes: Gołdapiwo and Mamry Północne. *Acta Parasitologica Polonica* 8: 85-114.
- Sulgostowska T. 1960b. Extra-intestinal trematodes in birds of the mesotropic lakes: Gołdapiwo and Mamry Północne. *Acta Parasitologica Polonica* 8: 471-492.
- Sulgostowska T. 1963. Trematodes of birds in the biocenosis of the lakes Drużno, Mamry Północne and Święcajty. *Acta Parasitologica Polonica* 11: 239-264.
- Sulgostowska T., Czaplińska D. 1987. Katalog fauny pasożytniczej Polski. Część IV, Pasożyty ptaków. Zeszyt 3, Pierwotniaki i przywry. PWN, Warszawa.
- Sulgostowska T., Grytner-Zięcina B. 1974. Trematodes of *Clangula hyemalis* from the Baltic Coast. *Acta Parasitologica Polonica* 22: 401-413.
- Szramka H. 2000. Skutki ochrony kormoranów w Nadleśnictwie Elbląg. *Las Polski* 3: 12-13.
- Tomiałojć L. 1990. Ptaki Polski, rozmieszczenie i liczebność. PWN, Warszawa.
- Yamaguti S. 1958. Systema Helminthum. Vol. 1. The digenetic trematodes of vertebrates. Interscience Publ., Inc., New York-London.
- Złotorzycka J. 1990. Katalog fauny pasożytniczej Polski. Część IV, Pasożyty ptaków. Zeszyt 3, Pasożytnicze stawonogi. PWN, Warszawa-Wrocław.
- Żuchowska, E. 2000. *Contracaecum rudolphii* Harwich, 1964 (Nematoda: Anisakidae) u kormoranów w Polsce. *Wiadomości Parazytologiczne* 46: 411-412.