

DOI 10.2478/pjvs-2013-0059

Original article

The effect of PCB126, 77, and 153 on the intracellular mobilization of Ca^{+2} in bovine granulosa and luteal cells after FSH and LH surge *in vitro*

J. Mlynarczuk, M. Kowalik

Institute of Animal Reproduction and Food Research, Polish Academy of Sciences,
Tuwima 10, 10-747 Olsztyn, Poland

Abstract

Polychlorinated biphenyls (PCBs) are a group of persistent environmental pollutants that impair cattle reproduction. Among other effects, PCBs can disturb the intracellular mobilization of Ca^{+2} in several cell types. Hence, it is possible that they disrupt the transduction of intracellular signals generated from gonadotropin (FSH/LH) receptors. In steroidogenic ovarian cells, a defect in Ca^{+2} mobilization may have a detrimental influence on two important processes: the secretion of steroids (E2 or/and P4) and their morphological and functional differentiation.

The aim of this study was to determine the influence of PCBs: 126 (dioxin-like) 77 (ambivalent) and 153 (estrogen-like) and a mixture of PCBs (Aroclor 1248) on these processes. Bovine granulosa and luteal cells were incubated for 72 hrs with PCBs (100 ng/ml), followed by Fura 2AM dye, and the fluctuations in intracellular Ca^{+2} mobilization after FSH/LH treatment were determined using an inverted microscope coupled with a CCD camera. The intensity and area of fluorescence excited by UV light were detected in the green spectrum of visible light. Aroclor 1248 and PCBs 153 and 77 significantly decreased ($P < 0.01-0.001$) the effect of FSH on intracellular Ca^{+2} mobilization in granulosa cells. In luteal cells, the most effective PCB on this process was PCB 77. The results revealed adverse effects of PCBs on the mobilization of intracellular Ca^{+2} . Moreover, the estrogen-like congeners were found to more effectively disturb this process than the dioxin-like PCB 126. Hence, it is possible for PCBs to have a negative influence on reproductive processes by affecting calcium mobilization.

Key words: PCB, calcium mobilization, FSH, LH, cattle

Introduction

Polychlorinated biphenyls (PCBs) belong to a class of persistent environmental pollutants. As endocrine disruptors, PCBs may impair many vital processes, including reproduction in humans and animals (Ulbrich et al. 2004, Kotwica et al. 2006, Meeker and Hauser 2010). These substances are often detected in bovine reproductive organs (Thomas et al. 1999, Glynn et al. 2000), including ovaries, where they are accumulated mainly in follicular fluid (Kamarianos et al. 2003). PCBs can disturb ovarian steroidogenesis (Roselli et al. 2000) as well as the secretion of oxytocin (OT) from bovine luteal cells (Młynarczuk and Kotwica 2005) and prostaglandins (PGE₂ and PGF₂α) from the myometrium (Wrobel et al. 2009), endometrium and endothelial cells from oviduct (Wrobel et al. 2010). Moreover, certain individual congeners impair the effect of FSH-stimulation on OT secretion from bovine granulosa cells (Młynarczuk et al. 2005) and suppress the effects of LH on progesterone (P₄) and OT secretion from bovine luteal cells (Młynarczuk and Kotwica 2006). The mechanism underlying this effect is not known, but it has been suggested that xenobiotics may disrupt the signals transmission, generated from membrane receptors (Nikula et al. 1999). Additionally, PCBs can disturb the intracellular concentration of Ca⁺² in human granulocytes (Voie and Fonnum 1998), decrease the concentrations of cAMP and phosphoinositols concentrations in cortical cells (Inglefield et al. 2001) and decrease intracellular calcium mobilization in bovine myometrial cells (Bae et al. 1999, Wrobel and Kotwica 2005) and mouse thymocytes (Yilmaz et al. 2006). Thus, PCBs may influence the affinity of some membrane receptors for their ligands or impair intracellular signaling pathways.

Both gonadotropins, FSH and LH, play a pivotal role in the regulation of bovine ovarian functions (Shemesh and Hannsel 1976, Baird 1992, Hunter et al. 2004). FSH is an essential factor for the recruitment and growth of antral follicles (Fortune 1993) and oocyte maturation (Armstrong et al. 1994, Adriaens et al. 2004). Additionally, LH is responsible for the survival of the dominant follicle to ovulation after FSH surge (Campbell et al. 1999), the alteration of follicular steroidogenesis towards P₄ synthesis and secretion (Brendtson 1995) and the initiation of luteinization in theca interna and granulosa cells (Fortune et al. 2001). The receptors for these hormones belong to a structurally unique subfamily of G-protein-coupled receptors (Ji et al. 1998) that are linked to the adenyl cyclase and phospholipase C signal transduction pathway (Zhu et al. 1994). After activation of these receptors by their ligands and signal

transduction initiation, Ca⁺² ions are released into the cytoplasm. Hence, intracellular fluctuations in Ca⁺² concentration are an indicator of these receptors activity (Zhu et al. 1994, Ryu et al. 1998).

In this study, ovarian steroidogenic cells were incubated with different individual congeners of PCBs or a mixture of them (Aroclor 1248) and treated with FSH or LH, after which intracellular calcium mobilization was measured. The estrogen-like congener PCB 153 (Katzenellenbogen et al. 1993), the dioxin-like congener PCB 126 (Lind et al. 1999), the bifunctional congener PCB 77 (Nesaretnam et al. 1996, Pang et al. 1999) and industrial mixture of PCBs – Aroclor 1248 were chosen for the experiments.

Materials and Methods

Collection of ovaries and cell preparation

Bovine ovaries were collected from cows or mature heifers at a defined stage of estrous cycle (Ireland et al. 1980, Fields and Fields 1996) at a commercial slaughterhouse within 15-20 min after killing the animals. The ovaries were placed in ice-cold 0.9% NaCl containing penicillin (10 IU/ml), streptomycin (100 µg/ml), amphotericin (2 µg/ml) and L-glutamine (100 µg/ml) and then transported to the laboratory in an ice bath within 1 h. Atretic follicles were eliminated on the basis of the procedure described by Henderson et al. (1987). All materials used were purchased from Sigma (Poznan, Poland) unless otherwise stated. The ovaries utilized for the collection of granulosa cells were from days 16-21 of the estrous cycle. For each experiment, cell mixtures from 10 follicles > 1 cm were collected via vigorous aspiration of follicular fluid (Voss and Fortune 1991). The cells were suspended in DMEM/HAM-12 medium supplemented with 10% NCS and transferred to 4-well plates (5 × 10⁴ cells/ml) (Nunclon NUNC, Denmark). After a 24-h preincubation, the medium was changed, and the cells were washed twice with M-199 medium containing 0.1% BSA. The DMEM/HAM-12 incubation medium (1 ml) was supplemented with 0.1% BSA, ascorbic acid (20 µg/ml), transferrin (5 µg/ml) and sodium selenite (5 ng/ml).

A suspension of luteal cells was obtained from the corpus luteum (CL) between days 5-10 of the estrous cycle (*n* = 3) by perfusion with a mixture of enzymes (1 mg/ml collagenase IA and 5 µg/ml DNase I) as described by Okuda et al. 1992. Cells showing a viability above 85% were used for further analyses. Luteal cells were suspended in DMEM/HAM-12 medium supplemented with 10% FCS and transferred to 4-well plates (5 × 10⁴ cells/ml) (Nunclon NUNC, Denmark). After a 24 hrs preincubation, the medium was

changed, and the cells were washed twice with M-199 medium containing 0.1% BSA. The incubation medium (1 ml) was the same as was used for granulosa cells.

Both, granulosa and luteal cells were then incubated with PCB: 126, 77, 153 and Aroclor 1248 at a dose of 100 ng/ml for 72 h. The cells were cultured in an atmosphere of air containing 5% CO₂ at 100% humidity at 38°C (Heraus BB-6060, Hanau, Germany). All of the media used were phenol red-free and contained 20 µg/ml gentamycin (ICN).

Preparation of cells for measuring Ca⁺² mobilization

After incubation, the cells were washed three times in DMEM/HAM-12 with 0.1% BSA, and calcium-free PBS was added to each well, supplemented with 0.1% BSA (500 µl) and Fura 2AM dye (5 µM/ml), as described by Gryniewicz et al. (1985). The cells were incubated for 60 min in a cells culture incubator and then washed three times again in DMEM/HAM-12 with 0.1% BSA. The wells were subsequently replenished with calcium-free PBS at half of the previous volume of medium (250 µl). The cells were cultured for 45 min in heater (ZALMED 32/250, Poland) at 30°C with a controlled atmosphere (95% air + 5% CO₂).

Determination of Ca⁺² mobilization

The fluctuations in intracellular Ca⁺² mobilization were determined using an inverted microscope (Olympus IX70) with a fluorescence objective (Olympus Fl, 40x) and a UV filter for Fura 2AM (340 nm wavelength). The total magnification was 400x. The Olympus VLS 100HG UV burner was used as the source of UV light. Images were collected (frame by frame) for 70 sec at 5 sec intervals using an Ikegami FL3 CCD color camera (at a VGA resolution of 640 × 480). The files were saved in the uncompressed TIFF format. The smallest scale of measurement was 3 × 3 pixel squares. The images were analyzed using graphics software (Olympus MicroImage 4.0, Japan). The intensity and area of fluorescence excited by the UV Fura 2AM dye were detected in the green spectrum of visible light (510-520 nm).

Preliminary studies

The spontaneous mobilization of intracellular Ca⁺² was visualized either with the shutter of the

microscope open during the entire time of the measurements (SPO) or with the shutter was open only during a specific time required to the acquire images by the camera (STO, 0.5-1 sec/picture). Luteal cells (from 3-4 CL and 5-10 days of the cycle) and granulosa cells from 10 follicles (> 1 cm in diameter) were used for imaging. The cells were prepared as previously described. Two luteal cells or 4 granulosa cells were placed in the photo frame of the microscope. The mean ± SEM of the first four points in each measurement was used as the control value. The means ± SEM of further points were compared with the control using ANOVA and Tukey's post-test. The obtained data (Fig. 1) showed that the constant exposure of cells treated with Fura to UV light (SPO) provoked a progressive, but statistically significant ($P < 0.05$) decline in dye luminescence after either 40 sec (luteal cells) or 45 sec (granulosa cells). However, when the shutter was opened only at the exposure time (STO), the luminescence intensity was not significantly changed ($P > 0.05$) during the time of measurement (70 sec); hence, this method was chosen for further investigations.

Detection of intracellular Ca⁺² mobilization

Intracellular Ca⁺² mobilization was induced by treating the cells with LH (luteal cells) or FSH (granulosa cells). The gonadotropins were microinjected (1 ng/5 µl of solvent) near the observed cells after the fourth image. The first four pictures were used to determine the control value of excitation. The obtained images were analyzed as described above.

Statistical analyses

For this study, luteal cells from 3 different CL were used, and detection was repeated twice for each of the groups of CL cells ($n = 6$). For granulosa cells, 3 cultures were used (each contained cells from 10 follicles > 1 cm in diameter), and detection was also repeated twice for each culture ($n = 6$). The means ± SEM of the green area and green intensity of the products in the next measurement points were obtained. Analogical points in the control and PCB treatment curves were compared with each other. The mean ± SEM areas under the curves were then analyzed. For these calculations, ANOVA and the Tukey post-test were used. The statistical procedures were performed with the software program PRISM 4.0 (GraphPad Software Inc., CA, USA).




Fig. 1. The effect of still and intermitted of UV stream light on the spontaneous luminescence of Fura 2AM in granulosa (a) and luteal (b) cells ($P < 0.05$). SPO – Shutter permanently open, STO – Shutter temporary open.

Results

Aroclor 1248 and PCBs 153 and 77 strongly decreased ($P < 0.01-0.001$) the effect of FSH on intracellular Ca^{+2} mobilization in granulosa cells (Fig. 2a). This effect was detected at every point where measurements were taken ($P < 0.01-0.001$), and it is visible as the decrease ($P < 0.01$) in the area under the curves compared with the control (Fig. 2b). The effect of PCB 126 was not as dramatic as that of the other congeners but was still statistically significant ($P < 0.05$).

In luteal cells, PCB 77 significantly decreased Ca^{+2} mobilization ($P < 0.05$) after LH surge. PCBs 126 and 153 showed a similar effect, however it was not significant ($P > 0.05$) at all points measured (for example at 45 sec of measurement, Fig. 3a). However, the area under the curve for all congeners was significantly smaller than area under the curve for the control (Fig. 3b). Although Aroclor 1248 decreased Ca^{+2} mobilization at individual points of measurement ($P < 0.05-0.001$), it did not significantly ($P > 0.05$) decrease the value of the area under the curve (Fig. 3b).




Fig. 2. The fluctuations of Ca^{+2} mobilization (a; a – $P < 0.001$, b – $P < 0.01$, c – $P < 0.05$) and changes in its total concentrations (b; as area under curve, $P < 0.05$) in granulosa cells treated by PCBs (100 ng/ml) after FSH surge.

Discussion

It was previously shown that individual congeners and Aroclor 1248 do not affect the viability of granulosa and luteal cells at doses between 1-100 ng/ml under 24-72 h of incubation (Mlynarczuk and Kotwica 2005). Hence, the results obtained here are not a result of decreasing cell viability.

Our results has shown that estrogen-like PCBs were more effective in disrupting the mobilization of intracellular Ca^{+2} than the dioxin-like PCB 126. However, it is difficult to identify which phase of signal transduction was impaired. It is well established that the interaction of LH with its receptor leads to the activation of adenylate cyclase through stimulatory guanine nucleotide regulatory protein (Gs protein),

leading to an increase in cyclic AMP levels. This reaction is related to intracellular Ca^{+2} mobilization (Mennon and Mennon 2012). A similar effect has been observed on E2 secretion in the mTLC-1 cell line when treated with the estrogen-like substance bisphenol A after an FSH surge (Nikula et al. 1999). Moreover, bisphenol A and another estrogen-like chloroorganic insecticide, DDT, have been shown to decrease the cAMP concentration in cells (Nikula et al. 1999, Pohland and Tiemann 2003, Wójtowicz et al. 2007). Hence, the influence of PCBs on intracellular Ca^{+2} mobilization may occur through a similar mechanism, which is dependent on the biological properties and chemical structure of the PCBs, although PCBs can affect different steps of the intracellular signal transduction pathway (Inglefield et al. 2001,




Fig. 3. The fluctuations of Ca²⁺ mobilization (*a*; *a* — $P < 0.001$, *b* — $P < 0.01$, *c* — $P < 0.05$, $n = 6$) and changes in its total concentrations (*b*; as area under curve, $P < 0.05$) in luteal cells treated by PCBs (100 ng/ml) after LH surge.

Llansola et al. 2009). The intermediate activity of the industrial PCBs mixture, Aroclor 1248, may have been the result of its composition. It contains both estrogen- and dioxin-like PCB congeners (Frame et al. 1996) which may partially antagonize each other effects (Suh et al. 2003).

This peculiar “desensitization” of steroidogenic ovarian cells to gonadotropins caused by PCBs can entail considerable consequences. It has been reported (Kawate 2004) that disturbances in the signals received and transduced from FSH and LH receptors may be one of the reasons for the formation of follicular cysts, which can further disturb or even disrupt the ovarian cycle. Estrogen-like chloroorganic substances, such as DDT and some PCB congeners, are also known to predispose both rats and humans to ovarian cyst formation (Heirichs et al. 1971, Jonsson et al.

1975, Gotz et al. 2001). Cystic ovarian disease is a significant problem for the reproduction of cattle and other domestic animals. This syndrome has been identified in 10-13% of dairy cows (Bartlett et al. 1986), and in some herds, ovarian cysts have been detected in 30-40% of cows (Archbald and Thatcher 1992). Disruption of the normal functioning of the hypothalamic-hypophyseal-ovarian axis is the basic cause of this illness, but its exact etiology is still poorly understood. Many factors may lead to a predisposition to this disease, including chloroorganic substances (Vanholder et al. 1997). Hence, a decrease in the reactivity of granulosa and luteal cells to gonadotropins caused by PCBs may constitute part of the mechanism responsible for the onset of cystic ovarian syndrome in cattle.

In conclusion, the PCB congeners tested here, de-

creased intracellular Ca^{2+} mobilization in granulosa and luteal cells after an FSH or LH surge, and these PCBs may disturb reproductive processes by this mechanism. The estrogen-like congeners PCBs 77 and 153 were the most effective in this process.

References

- Adriaens I, Cortvriendt R, Smitz J (2004) Differential FSH exposure in preantral follicle culture has marked effects on folliculogenesis and oocyte developmental competence. *Hum Reprod* 19: 398-408.
- Archbald LF, Thatcher WW (1992) Ovarian follicular dynamics and management of ovarian cysts. In: van Horn HH, Wilcox CJ (eds), *Large Dairy Herd Management*. 1 ed., American Dairy Science Association, Champaign, pp 199-208.
- Armstrong DT, Irvine BJ, Earl CR, McLean D, Seamark RF (1994) Gonadotropin stimulation regimens for follicular aspiration and in vitro embryo production from calf oocytes. *Theriogenology* 42: 1227-1236.
- Bae J, Stuenkel EL, Loch-Carusio R (1999) Stimulation of oscillatory uterine contraction by the PCB mixture Aroclor 1242 may involve increased $[\text{Ca}^{2+}]_i$ through voltage-operated calcium channels. *Toxicol Appl Pharmacol* 155: 261-272.
- Baird DT (1992) Luteotrophic control of the corpus luteum. *Anim Reprod Sci* 28: 95-102.
- Bartlett PC, Ngategize PK, Kaneene JB, Kirk JH, Anderson SM, Mather EC (1986) Cystic follicular disease in Michigan Holstein-Friesian cattle: incidence, descriptive epidemiology and economic impact. *Prev Vet Med* 4: 15-33.
- Brendtson AK, Vincent SE, Fortune JE (1995) Low and high concentrations of gonadotropins differentially regulate hormone production by theca interna and granulosa cells from bovine preovulatory follicles. *Biol Reprod* 52: 1334-1342.
- Campbell BK, Dobson H, Baird DT, Scaramuzzi RJ (1999) Examination of the relative role of FSH and LH in the mechanism of ovulatory follicle selection in sheep. *J Reprod Fertil* 117: 355-367.
- Fields MJ, Fields PA (1996) Morphological characteristics of the bovine corpus luteum during the estrous cycle and pregnancy. *Theriogenology* 45: 1295-1325.
- Fortune JE (1993) Follicular dynamics during the bovine estrous cycle: a limiting factor in improvement of fertility? *Anim Reprod Sci* 33: 111-125.
- Fortune JE, Rivera GM, Evans AC, Turzillo AM (2001) Differentiation of dominant versus subordinate follicles in cattle. *Biol Reprod* 65: 648-654.
- Frame GM, Cochran JW, Bowadt SS (1996) Complete PCB congener distributions for 17 Aroclor mixtures determined by 3 HRCG systems optimized for comprehensive, quantitative, congener-specific analysis. *J High Res Chromatog* 19: 657-668.
- Glynn AW, Wernroth L, Atuma S, Linder C-E, Aune M, Nilsson I, Darnerud PO (2000) PCB and chlorinated pesticide concentrations in swine and bovine adipose tissue in Sweden 1991 – 1997: spatial and temporal trends. *Sci Total Environ* 246: 195-206.
- Götz F, Thieme S, Dörner G (2001) Female infertility – effect of perinatal xenoestrogen exposure on reproductive functions in animals and humans. *Folia Histochem Cytobiol* 39; (Suppl 2): 40-43.
- Gryniewicz G, Poenie M, Tsien RY (1985) A new generation of Ca^{2+} indicators with greatly improved fluorescence properties. *J Biol Chem* 260: 3440-3450.
- Henderson KM, McNatty KP, Smith P, Gibb M, O'Keeffe LE, Lun S, Heath DA, Prisk MD (1987) Influence of follicular health on the steroidogenic and morphological characteristics of bovine granulosa cells *in vitro*. *J Reprod Fertil* 79: 185-193.
- Heirichs WL, Gellert RJ, Bakke JL, Lawrence NL (1971) DDT administered to neonatal rats induces persistent estrus syndrome. *Science* 173: 642-643.
- Hunter MG, Robinson RS, Mann GE, Webb R (2004) Endocrine and paracrine control of follicular development and ovulation rate in farm species. *Anim Reprod Sci* 82-83: 461-477.
- Inglefield JR, Mundy WR, Shafer TJ (2001) Inositol 1,4,5-triphosphate receptor-sensitive Ca^{2+} release, store-operated Ca^{2+} entry, and cAMP responsive element binding protein phosphorylation in developing cortical cells following exposure to polychlorinated biphenyls. *J Pharmacol Exp Ther* 297: 762-773.
- Ireland JJ, Murphee RL, Coulson PB (1980) Accuracy of predicting stages of bovine estrous cycle by gross appearance of the corpus luteum. *J Dairy Sci* 63: 155-160.
- Ji TH, Grossmann M, Ji I (1998) G protein-coupled receptors. I. Diversity of receptor-ligand interactions. *J Biol Chem* 273: 17299-17302.
- Jonsson HT Jr, Keil JE, Gaddy RG, Loadholt CB, Hennigar GR, Walker EM Jr (1975-1976) Prolonged ingestion of commercial DDT and PCB; effects on progesterone levels and reproduction in the mature female rat. *Arch Environ Contam Toxicol* 3: 479-490.
- Kawate N (2004) Studies on the regulation of expression of luteinizing hormone receptor in the ovary and the mechanism of follicular cyst formation in ruminants. *J Reprod Dev* 50: 1-8.
- Kamarianos A, Karamanlis X, Goulas P, Theodosiadou E, Smokovitis A (2003) The presence of environmental pollutants in the follicular fluid of farm animals (cattle, sheep, goats and pigs). *Reprod Toxicol* 17: 185-195.
- Katzenellenbogen BS, Bhardwaj B, Fang H, Ince BA, Pakdel F, Reese JC, Schodin D, Wrenn CK (1993) Hormone binding and transcription activation by estrogen receptors: analyses using mammalian and yeast systems. *J Steroid Biochem Mol Biol* 47: 39-48.
- Kotwica J, Wróbel M, Młynarczuk J (2006) The influence of polychlorinated biphenyls (PCBs) and phytoestrogens in vitro on functioning of reproductive tract in cow. *Reprod Biol* 6 (Suppl 1): 189-194.
- Lind PM, Eriksen EF, Sahlin L, Edlund M, Orberg J (1999) Effects of the antiestrogenic environmental pollutant 3,3',4,4',5-pentachlorobiphenyl (PCB-126) in rat bone and uterus: diverging effects in ovariectomized and intact animals. *Toxicol Appl Pharmacol* 154: 236-244.
- Llansola M, Piedrafita B, Rodrigo R, Montoliu C, Felipo V (2009) Polychlorinated biphenyls PCB 153 and PCB 126 impair the glutamate-nitric oxide-cGMP pathway in cerebellar neurons in culture by different mechanisms. *Neurotox Res* 16: 97-105.
- Meeker JD, Hauser R (2010) Exposure to polychlorinated biphenyls (PCBs) and male reproduction. *Syst Biol Reprod Med* 56: 122-131.

- Menon KM, Menon B (2012) Structure, function and regulation of gonadotropin receptors – a perspective. *Mol Cell Endocrinol* 356: 88-97.
- Młynarczuk J, Kotwica J (2005) Influence of polychlorinated biphenyls on the secretion of oxytocin from bovine luteal cells and from granulosa cells obtained from the follicles of different size. *Pol J Vet Sci* 8: 261-267.
- Młynarczuk JJ, Niewiadowska A, Kotwica J (2005) Polychlorinated biphenyls impair FSH-stimulated effect on the secretion of steroids and oxytocin from cultured bovine granulosa cells. *Bull Vet I Pulawy* 49: 411-417.
- Młynarczuk J, Kotwica J (2006) Influence of polychlorinated biphenyls on LH-stimulated secretion of progesterone and oxytocin from bovine luteal cells. *Pol J Vet Sci* 9: 101-108.
- Nesaretnam K, Corcoran D, Dils RR, Darbre P (1996) 3,4,3',4'-Tetrachlorobiphenyl acts as an estrogen *in vitro* and *in vivo* model. *Mol Endocrinol* 10: 923-936.
- Nikula H, Talonpoika T, Kaleva M, Toppari J (1999) Inhibition of hCG-stimulated steroidogenesis in cultured mouse Leydig tumor cells by bisphenol A and octylphenols. *Toxicol Appl Pharmacol* 157: 166-173.
- Okuda K, Miyamoto A, Sauerwein H, Schweigert FJ, Schams D (1992) Evidence for oxytocin receptors in cultured bovine luteal cells. *Biol Reprod* 46: 1001-1006.
- Pang S, Cao JQ, Katz BH, Hayes CL, Sutter TR, Spink DC (1999) Inductive and inhibitory effects of non-ortho-substituted polychlorinated biphenyls on estrogen metabolism and human cytochromes P450 1A1 and 1B1. *Biochem Pharmacol* 58: 29-38.
- Pöhlend R, Tiemann U (2003) Forskolin-induced cyclic AMP signaling in single adherent bovine oviductal cells: effect of dichlorodiphenyltrichloroethane (DDT) and tris(4-chlorophenyl)methanol (TCPM). *Toxicol In Vitro* 17: 375-383.
- Roselli M, Reinhart K, Imthurn B, Keller PJ, Dubey RK (2000) Cellular and biochemical mechanisms by which environmental oestrogens influence reproductive function. *Hum Reprod Update* 6: 332-350.
- Ryu KS, Gilchrist RL, Koo YB, Ji I, Ji TH (1998) Gene, interaction, signal generation, signal divergence and signal transduction of the LH/GC receptor. *Int J Gynecol Obstet* 60: S9-S20.
- Shemesh M, Hansel W (1976) Steroid synthesis by ovarian follicles in response to LH and PGF₂α. *Proc Soc Exp Biol Med* 152: 86-89.
- Suh J, Kang JS, Yang KH, Kaminski NE (2003) Antagonism of aryl hydrocarbon receptor-dependent induction of CYP1A1 and inhibition of IgM expression by di-ortho-substituted polychlorinated biphenyls. *Toxicol Appl Pharmacol* 187: 11-21.
- Thomas GO, Sweetman AJ, Jones KC (1999) Metabolism and body-burden of PCBs in lactating dairy cows. *Chemosphere* 39: 1533-1544.
- Ulbrich B, Stahlmann R (2004) Developmental toxicity of polychlorinated biphenyls (PCBs): a systematic review of experimental data. *Arch Toxicol* 78: 252-268.
- Vanholder T, Opsomer G, de Kruijff, A (1997) Aetiology and pathogenesis of cystic ovarian follicles in dairy cattle: a review. *Reprod Nutr Dev* 46: 105-119.
- Voie OA, Fonnum F (1998) Ortho substituted polychlorinated biphenyls elevate intracellular [Ca(2+)] in human granulocytes. *Environ Toxicol Pharmacol* 5: 105-112.
- Voss AK, Fortune JE (1991) Oxytocin stimulates progesterone production by bovine granulosa cells isolated before, but not after, the luteinizing hormone surge. *Mol Cell Endocrinol* 78: 17-24.
- Wójtowicz AK, Kajta M, Gregoraszczyk EŁ (2007) DDT- and DDE-induced disruption of ovarian steroidogenesis in prepubertal porcine ovarian follicles: a possible interaction with the main steroidogenic enzymes and estrogen receptor beta. *J Physiol Pharmacol* 58: 873-885.
- Wrobel M, Kotwica J (2005) Effect of polychlorinated biphenyls (PCBs) on basal and OT-stimulated calcium concentrations in myometrial cells in cow. *Reprod Biol* 5: 321-330.
- Wrobel MH, Rekawiecki R, Kotwica J (2009) Involvement of prostaglandin F₂α in the adverse effect of PCB 77 on the force of contractions of bovine myometrium. *Toxicology* 262: 224-229.
- Wrobel MH, Młynarczuk J, Kotwica J (2010) Influence of polychlorinated biphenyls and their hydroxylated metabolites on prostaglandins secretion from epithelial cells of bovine oviduct, *in vitro*. *Toxicology* 270: 85-91.
- Yilmaz B, Sandal S, Chen CH, Carpenter DO (2006) Effects of PCB 52 and PCB 77 on cell viability, [Ca(2+)](i) levels and membrane fluidity in mouse thymocytes. *Toxicology* 217: 184-193.
- Zhu X, Gilbert S, Birnbaumer R, Birnbaumer L (1994) Dual signaling potential is common among Gs-coupled receptors and dependent on receptor density. *Mol Pharmacol* 46: 460-469.