

SELENIUM IN THE DIET OF RUMINANTS

Barna Kruzhel¹, Małgorzata Bąkowska², Stakh Vovk³,
Ewa Nowakowska², Pavkovych Sergei¹

¹Lviv State Agrarian University, Ukraine

²West Pomeranian University of Technology, Szczecin, Poland

³Institute of Agriculture in the Carpathian Region, National Academy
of Agricultural Sciences of Ukraine

Abstract. The results of the research show that there is a deficiency in Se in the environment in most regions. It is crucial to supplement selenium deficiency in ruminants from deficient regions as long-lasting lack of this element may lead to many diseases. Selenium serum or blood level can be an indicator of the supply of this element in ruminants (intravittally) as well as its content in the liver and kidneys. The most commonly used criterion for assessing the selenium status in livestock is its serum or blood content. At a low concentration of this element in the serum/blood of ruminants, its supplementation is necessary.

Key words: diseases, ruminants, selenium

INTRODUCTION

Studies on the role of selenium in animals are carried out for many years. This element belongs to the so-called trace elements, which are necessary for keeping the proper functions of an organism.

Selenium is not evenly distributed in the environment, there are areas with soils either poor or rich in this element. Most soils in the world have low content of this element [Kabata-Pendias and Pendias 1999, Borowska 2002, Cartes et al.

2005, Broadley et al. 2006]. According to Kabat-Pendias and Pendias [1999], the average content of selenium in soils on a global scale is equal to $0.33 \text{ mg} \cdot \text{kg}^{-1}$. Sandy soils generally have typically low quantities of this element (from 0.06 to $0.4 \text{ mg} \cdot \text{kg}^{-1}$), while the soils that originate from the formations rich in selenium (tertiary loams) may contain significantly higher quantities of this element ($2.3 \div 4.2 \text{ mg} \cdot \text{kg}^{-1}$). Deficit of this element occurs mainly in certain areas of China, Canada, Denmark, New Zealand and the Nordic countries [Yläranta 1990, Mäkelä-Kurtto and Sippola 2002, Hartikainen 2005]. Borowska [2010] claims that the so-called selenium provinces can be unique in the world. The content of selenium in these regions can even reach $1200 \text{ mg} \cdot \text{kg}^{-1}$ (e.g., Japan, Canada, USA).

This element as a component of selenoproteins plays an important structural and enzymatic function. According to Flohe et al. [2000] and Rayman [2000], the main selenoproteins include glutathione peroxidase, iodothyronine deiodinase, selenoproteins P and W and thioredoxin reductase. Selenium is an inactivator of toxic heavy metals [Rayman 2000], neutralizes the effect of aflatoxins [Navarro-Alarcon et al. 1998, Maehira et al. 2002], causes apoptosis in tumor cells [Griffin 1982, Behne et al. 1996], prevents or delays the cell aging by protecting the mitochondrial membranes [Wesołowski and Ulewicz 2000], protects the cell lipids from damaging effects of reactive oxygen species [Musik et al. 2003].

The consequences of selenium deficiency in ruminants

Many authors draw attention to the fact that the concentration of selenium in the body of animals depends primarily on the feed consumed, which enrichment largely depends on the content of this element in the soil [Pilarczyk et al. 2004, 2008 a, b, 2009, 2010, 2011 a, b, c, Sablik et al. 2011, Seremak et al. 2011, Flueck et al. 2012, Kawecka et al. 2013, Nowakowska 2013] (Table 1). According to Mayland [1994], and Kabat-Pendias and Pendias [1999], the content of selenium in the feed below $0.010 \text{ mg} \cdot \text{kg}^{-1}$ d.w. can cause deficiency symptoms in animals.

Selenium deficiency in cattle and sheep can cause nutritional muscular dystrophy (white muscle disease), which manifests in the stiffness of the limbs, difficulty in grasping the food and swallowing, aspiration pneumonia and myoglobinuria [Rederstorff et al. 2006, McCann and Ames 2011]. Selenium deficiency in cattle may also cause diarrhea in young animals and the increase in cirrhosis cases in beef cattle. In cows, placenta retention may occur after delivery, inflammation of the endometrium, fetal death due to their impaired development, extension of the calving and parturitional intervals, and lower breeding performance of bulls [Hulland 1985, Bostedt and Schramel 1990, Zust et al. 1996, Radostits et al. 2000, Suttle 2010].

Table 1. Selected Se content in ruminants from different regions

Tabela 1. Wybrane stężenia Se u przeżuwaczy z różnych regionów

Location Rejon	Tissue Tkanka	Se content Stężenie Se	Reference Piśmiennictwo
Cows – Krowy			
Ukraine – Ukraina	serum – surowica	0.085 µg · ml ⁻¹ 0.034 µg · ml ⁻¹	Pilarczyk et al. [2005] Pilarczyk et al. [2005]
Poland – Polska	serum – surowica	0.074 µg · ml ⁻¹ 0.020 µg · ml ⁻¹ 233.7–237.6 nmol · dm ⁻³	Sablik et al. [2011] Pilarczyk et al. [2013 c] Bombik et al. [2010]
Norway – Norwegia	blood – krew	0.090 µg · ml ⁻¹	Kommisrud et al. [2005]
Czech Republic – Czechy	plasma – osocze	0.031 µg · ml ⁻¹	Slavík et al. [2007]
Czech Republic – Czechy	blood – krew	0.038–0.095 µg · ml ⁻¹	Pavlata et al. [2002]
USA	serum – surowica	0.038–0.077 µg · ml ⁻¹	Segerson et al. [1981]
Slovenia – Słowenia	serum – surowica	0.014–0.017 µg · ml ⁻¹	Zust et al. [1996]
Romania – Rumunia	blood – krew	0.026 µg · ml ⁻¹	Pop et al. [2011]
Romania – Rumunia	serum – surowica	0.011 µg · ml ⁻¹	Pârvu et al. [2003]
Romania – Rumunia	hair – włosy	0.140 µg · g ⁻¹	
Columbia – Kolumbia	serum – surowica	0.052–0.097 µg · ml ⁻¹	Fenimore et al. [1983]
Water buffalos – Bawoły rzeczne			
India – Indie		0.190 µg · ml ⁻¹	Prasad and Arora [1991]
Poland – Polska	serum – surowica	0.021 µg · ml ⁻¹	Sablik et al. [2011]
Pakistan – Pakistan		0.100 µg · ml ⁻¹	Akhtar et al. [2009]
Sheeps – Owce			
Ukraine – Ukraina	serum – surowica	0.485 µg · ml ⁻¹	Pilarczyk et al. [2005]
Poland – Polska	serum – surowica wool – wełna	3.28–0.23 mmol · l ⁻¹ 6.51–4.24 mmol · l ⁻¹	Pięta and Patkowski [2007]
Poland – Polska	serum – surowica	0.031 µg · ml ⁻¹	Pilarczyk et al. [2005]
Poland – Polska	liver – wątroba	3.62 µg · g ⁻¹ w.w.	
Poland – Polska	kidneys – nerki	1.59 µg · g ⁻¹ w.w.	Słupczyńska et al. [2009]
Poland – Polska	muscle – mięsień	0.04 µg · g ⁻¹ w.w.	
Iran – Iran	blood – krew	0.028 µg · l ⁻¹	Karimi-Poor et al. [2011]
USA	blood – krew	0.090 µg · ml ⁻¹	Coggins [2006]
USA	liver – wątroba	0.173 µg · g ⁻¹ w.w.	Cox [2006]
Turkey – Turcja	serum – surowica	0.304 µg · ml ⁻¹	Pamukcu et al. [2001]
German – Niemcy	serum – surowica	0.045 µg · ml ⁻¹	Humann-Ziehank et al. [2013]
Goats – Kozy			
Poland – Polska	liver – wątroba muscle – mięsień kidneys – nerki lungs – płuca heart – serce	0.047 µg · g ⁻¹ w.w. 0.027 µg · g ⁻¹ w.w. 0.537 µg · g ⁻¹ w.w. 0.056 µg · g ⁻¹ w.w. 0.035 µg · g ⁻¹ w.w.	Pilarczyk et al. [2013 b]
Republic of Burundi Republika Burundi	kidney – nerka liver – wątroba	1.194 µg · g ⁻¹ w.w. 0.427 µg · g ⁻¹ w.w.	Benemariya et al. [1993]
Czech Republic – Czechy	blood – krew	0.068 µg · ml ⁻¹	Pechowa et al. [2012]
Croatia – Chorwacja	blood – krew	0.019–0.036 µg · ml ⁻¹	Antunović et al. [2013]
Mexico – Meksyk	serum – surowica	0.020–0.021 µg · ml ⁻¹	Ramírez-Bribiesca et al. [2001]
USA	serum – surowica	0.010 µg · ml ⁻¹	Harris [1987]
USA	blood – krew	0.076–0.499 µg · ml ⁻¹	McComb et al. [2010]
France – Francja	serum – surowica	0.021 µg · ml ⁻¹	Morand-Fehr [1981]

A decrease in the efficiency of the antioxidant system is observed in cows at high milk yields, particularly when combined with selenium deficiency [Flohe et al. 2000, Gladyshev 2001, Jankowiak et al. 2015]. Therefore, it is important to adequately supply females in selenium, which determines the efficacy of the antioxidant system [Castillo et al. 2006, Pilarczyk et al. 2012]. Selenium deficiency during high lactation increases the risk of mastitis [Villar et al. 2002].

Myocardial damage may occur in calves as a result of selenium and vitamin E deficiency. Three-month-old animals and younger are affected most frequently. Symptoms of deficiency of this element include: arrhythmia, murmurs, shortness of breath, congestive heart failure, which in turn may lead to sudden death [Rederstorff et al. 2006, McCann and Ames 2011].

Selenium deficiencies in sheep in New Zealand and Australia were the cause of 20–50% of infertility in females and also an increased lamb loss was observed [Suttle 2010, Ren et al. 2011]. These authors showed that Se deficiency also negatively affected the rams (the number of sperm in the ejaculate, sperm concentration and motility). Fetal death may also occur in pregnant ewes as well as placenta retention after birth [Ren et al. 2011].

The consequences of excess selenium in ruminants

Excess of selenium causes selenosis, a disease which is characterized by muscle degeneration, heavy breathing, roughness and matt hair, cardiovascular disorders [Ramadan et al. 2001, Deore et al. 2002, Terzano et al. 2005]. Poisoning associated with the intake of excess selenium in ruminant farming is occasional. There were no such cases reported in Poland and Ukraine. Selenium lethal dose (LD_{50}) in adult cattle is $0.501 \text{ mg} \cdot \text{kg}^{-1}$ body weight, while in lambs it is $0.455 \text{ mg} \cdot \text{kg}^{-1}$ [Grace 1994, Tinggi 2005]. Blood selenium levels of $>2 \text{ mg} \cdot \text{l}^{-1}$ in cattle and $>0.6\text{--}0.7 \text{ mg} \cdot \text{l}^{-1}$ in sheep are associated with selenosis with borderline toxicity at $1\text{--}2 \text{ mg} \cdot \text{l}^{-1}$ in cattle [Levander 1986, WHO 1987].

Standards of selenium supply in ruminants

Selenium serum or blood level can be an indicator of the supply of this element in ruminants (intravital) as well as its content in the liver and kidneys. Puls [1994] points that the concentration of selenium in the liver is a better indicator of selenium level than the kidney test. Oh et al. [1976] demonstrated that lambs fed with fodder poor in selenium had always a higher concentration of this element in the kidney compared to liver. The opposite situation was observed when a feed rich in this element was supplied to lambs (higher concentration of selenium in

the liver than in the kidneys). Mahan and Kim [1996] believe that the assessment of selenium status of the organism should involve blood or serum tests.

The most commonly used criterion for assessing the selenium status in livestock is the serum or blood content. According to Grace [1997], biochemical standards used in the diagnosis of serum selenium deficit in ruminants are as follows: less than $0.041 \text{ g} \cdot \text{ml}^{-1}$ – deficiency; $0.041\text{--}0.079 \text{ g} \cdot \text{ml}^{-1}$ – marginal level; above $0.079 \text{ g} \cdot \text{ml}^{-1}$ – physiological level. According to Smith et al. [1998], the following biochemical standards should be used in the diagnosis of whole blood selenium deficit in ruminants: less than $0.14 \text{ g} \cdot \text{ml}^{-1}$ – deficiency; $0.14\text{--}0.20 \text{ g} \cdot \text{ml}^{-1}$ – marginal level; above $0.20 \text{ g} \cdot \text{ml}^{-1}$ – physiological level. According to Puls [1994], biochemical standards used in the diagnosis of selenium deficit in the liver of ruminants should be as follows: less than $0.1 \text{ g} \cdot \text{ml}^{-1}$ of the above – deficiency; $0.1\text{--}0.24 \text{ g} \cdot \text{ml}^{-1}$ – marginal level; $0.25\text{--}1.2 \text{ g} \cdot \text{ml}^{-1}$ – optimal level.

Selenium supplementation in ruminants

The results shown in Table 1 confirm that in some geographical areas there is a deficiency of selenium in ruminants. At a low concentration of this element in the serum/blood of ruminants, its supplementation is necessary. Selenium may be supplemented in the inorganic or organic form. European Union permits supplementation of ruminant feed additives with sodium selenite, sodium selenate and selenium yeast *Saccharomyces cerevisiae*. Sodium selenite is most frequently applied in the feed industry. Selenium yeast are a source of organic selenium (mainly selenomethionine).

The bioavailability of selenium depends on the species of animals. For pigs, Se is available in both: organic and inorganic compounds. It was found that in pigs 77% of Se in the form of selenite was resorbed in the gastrointestinal tract, while in sheep it was only about 29% [Grela and Sembratowicz 1997]. Mineral selenium compounds in ruminants are strongly reduced by rumen bacteria to non-assimilable forms. According to Gerloff [1992], the factor that reduces the absorption is low rumen pH, typical for the high-yielding cows that receive high doses of energy feed.

Pilarczyk et al. [2004] found a significant improvement in reproductive indices (fertility – 96%, and fecundity – 137.5%) after application of sodium selenite in ewes. According to the authors, lamb losses in the experimental group were lower (9.2%) than in the control group (12.1%). Similar effects were obtained using selenium yeast [Pilarczyk et al. 2013 a]. Deaths in the group of lambs derived from the ewes fed selenium yeast were significantly lower, body weight of lambs on the day of birth was higher, the number of live-born lambs was also higher compared to the control group. Lambs derived from the mothers that were admi-

nistered selenium yeast had significantly higher body weight at 33 and 90 days of age compared to lambs of the mothers not receiving Se. Reproductive indices (fertility and fecundity) were also improved as a result of the application of selenium yeast in ewes. Segerson et al. [1986] established that the number of lambs born alive from control and Se-treated ewes was 1.61 and 1.81, respectively, while the number of lambs weaned per ewe was 1.1 and 1.5, respectively.

REFERENCES

- Akhtar M.S., Farooq A.A., Mushta Q.M., 2009. Serum concentrations of copper, iron, zinc and selenium in cyclic and anoestrus nili-ravi buffaloes kept under farm conditions. *Pakistan Vet. J.* 29 (1), 47–48.
- Antunović Z., Klapac T., Čavar S., Šperanda M., Pavić M., Novoselec J., Klir Ž., 2013. Status of selenium and correlation with blood GSH-Px in goats and their kids in organic breeding fed with different levels of organic selenium. *Archiv Tierzucht* 56 (16), 169–177.
- Behne D., Kyriakopoulos A., Weiss-Nowak C., Kalckloesch M., Westphal C., Gessner H., 1996. Newly found selenium-containing proteins in the tissues of the rat. *Biol. Trace Elem. Res.* 55, 99–100.
- Benemariya H., Robberecht H., Deelstra H., 1993. Zinc, copper, and selenium in milk and organs of cow and goat from Burundi, Africa. *Sci. Total Environ.* 128, 83–98.
- Bombik T., Bombik E., Górski K., Rymuz A.K., Saba L., 2010. Selenium content in feed and cows' blood serum in the central-eastern Poland. *Bull. Vet. Inst. Pulawy* 54, 273–276.
- Borowska K., 2002. Wpływ nawożenia organicznego i zróżnicowanych dawek azotu na zawartość selenu w bulwach ziemniaka i glebie na tle jej niektórych właściwości [Influence of organic fertilization and diversified nitrogen doses on selenium content in potatoes and soil, according to its selected features]. *Acta Agrophys.* 70, 55–63 [in Polish].
- Borowska K., 2010. Selen w glebie i roślinach w warunkach zróżnicowanego nawożenia organicznego i mineralnego [Selenium in soil and plants under conditions of different organic and mineral fertilization]. Wydaw. Univ. Technologiczno-Przyrodniczego, Bydgoszcz, Rozprawa nr 140. [in Polish].
- Bostedt H., Schramel P., 1990. The importance of selenium in the prenatal and postnatal development of calves and lambs. *Biol. Trace Elem. Res.* 24, 163–71.
- Broadley M.R., White P.J., Bryson R.J., Meacham M.C., Bowen H.C., Johnson S.E., Hawkesford M.J., Mc Grath S.P., Zhao F.J., Breward N., Harriman M., Tucker M., 2006. Biofortification of UK food crops with selenium. *Proc. Nutr. Soc.* 65, 169–181.
- Cartes P., Gianfreda L., Mora M.L., 2005. Uptake of selenium and its antioxidant activity on ryegrass when applied as selenate and selenite forms. *Plant Soil.* 276, 359–367.
- Castillo C., Hernández J., Valverde I., Pereira V., Sotillo J., López Alonso M., Benedito J.L., 2006. Plasma malonaldehyde (MDA) and total antioxidant status (TAS) during lactation in dairy cows. *Res. Vet. Sci.* 80, 133–139.

- Coggins V.L., 2006. Selenium supplementation, parasite treatment, and management of bighorn sheep at Lostine River, Oregon. Bienn. Symp. North Wild Sheep Goat Counc. 15, 98–106.
- Cox M.K., 2006. Effects of mineral supplements on California bighorn sheep in northern Nevada. Bienn. Symp. North Wild Sheep and Goat Counc. 15, 107–120.
- Deore M.D., Srivastava K.A., Sharma S.K., 2002. Blood selenium levels during different stages of selenosis in buffaloes and its evaluation as a diagnostic tool. Vet. Hum. Toxicol. 44 (5), 260–263.
- Fenimore R.L., Adams D.S., Puls R., 1983. Selenium Levels of Beef Cattle in Southeastern British Columbia Relative to Supplementation and Type of Pasture. Can Vet J. 24 (2), 41–45.
- Flohe L., Andreesen J.R., Brigelius-Flohe R., Maiorino M., Ursini F., 2000. Selenium, the element of the moon, in life on earth. IUBMB Life, 49, 411–420.
- Flueck W.T., Smith-Flueck J.M., Mionczynski J., Mincher B.J., 2012. The implications of selenium deficiency for wild herbivore conservation: a review. Eur. J. Wildl. Res. 58 (5), 761–780.
- Gerloff B., 1992. Effect of selenium supplementation on dairy cattle. J. Anim. Sci. 12, 3934–3940.
- Gladyshev V.N., 2001. Selenium in biology and human health: controversies and perspectives. In: D.L. Hatfield (ed.). Selenium: Its Molecular Biology and Role in Human Health, Boston, Kluwer, 313–317.
- Grace N.D., 1994. Selenium. In: Managing trace element deficiencies. N.D. Grace (ed.). New Zealand
- Grace N.D., 1997. Use of biochemical criteria to diagnose trace element deficiencies in sheep and cattle [in: Proceeding of 9th International Conference on Animal Production] Berlin 11–14 September 1997. Verlag Sttudgard.
- Grela E.R., Sembratowicz I., 1997. Organiczne związki selenu w żywieniu zwierząt [Organic selenium compounds in animal nutrition]. Med. Weter. 53 (7), 385–386 [in Polish].
- Griffin A.C., 1982. The chemopreventive role of selenium in carcinogenesis. In: Molecular interrelations of nutrition and cancer. Arnott MS, van Eys J, Wang YM (eds.). New York, NY: Raven Press, 401–408.
- Harris B.Jr., 1987. Feeding and management of dairy goats. FCES Publication No. 761, University of Florida, Gainesville, FL, USA, 1–10.
- Hartikainen H., 2005. Biogeochemistry of selenium and its impact on food chain quality and human health. J. Trace Elem. Med. Bio. 18, 309–318.
- Hulland T.J., 1985. Muscles and Tendons. In Pathology of Domestic Animals, Third Edn, (eds) Jubb K.V.F., P.C. Kennedy, Palmer 1, Academic Press Inc., London, 140–195.
- Humann-Ziehank E., Tegtmeyer P.C., Seelig B., Roehrig P., Ganter M., 2013. Variation of serum selenium concentrations in German sheep flocks and implications for herd health management consultancy. Acta Veterinaria Scandinavica 55, 82–83.
- Jankowiak D., Pilarczyk R., Drozd R., Pilarczyk B., Tomza-Marciniak A., Wysocka G., Rząd I., Drozd A., Kuba J., 2015. Activity of antioxidant enzymes in the liver of wild boars (*Sus scrofa*) from a selenium-deficient area depending on sex, age, and season of the year. Turk. J. Biol. 39, 129–138.
- Kabata-Pendias A., Pendias H., 1999. Biochemia pierwiastków śladowych [Biogeochemistry of trace elements]. PWN, wyd. II, Warszawa. [in Polish].

- Karimi-Poor M., Sayed-Noroden T., Zamani F., Pirestani A., Bahrami Y., 2011. Investigation of Selenium concentration of sheep's diet, blood and milk in different regions from a central state of Iran. *Ann. Biol. Res.* 2 (3), 51–61.
- Kawecka, M., Jacyno, E., Matysiak, B., Kołodziej-Skalska, A., Pietruszka, A., 2013. Effects of selenium and vitamin E supplementation on selenium distribution and meat quality of pigs. *Acta Agriculturae Scandinavica A: Animal Sciences* 63 (4), 194–200.
- Kommisrud E., Østerås O., Vatn T., 2005. Blood Selenium Associated with Health and Fertility in Norwegian Dairy Herds. *Acta Vet. Scand.* 46, 229–240.
- Levander O. A., 1986. Selenium. In *Trace Elements in Human and Animal Nutrition* (Mertz M., Ed.), Academic Press, London, 139–197.
- Maehira G., Luyo G.A., Oshiro M., 2002. Alterations of serum selenium concentrations in the acute phase of pathological conditions. *Clin. Chim. Acta*, 316, 137–146.
- Mahan D.C., Kim Y.Y., 1996. Effect of inorganic or organic selenium at two dietary levels on reproductive performance and tissue selenium concentrations in first-parity gilts and their progeny. *J. Anim. Sci.* 74, 2711–2718.
- Mäkelä-Kurtto R., Sippola J., 2002. Monitoring of Finnish arable land: changes in soil quality between 1987 and 1998. *Agr. Food Sci. Finland*, 11, 273–284.
- Mayland H.F., 1994. Selenium in plants and animal nutrition. [In] *Selenium in the Environment*, W.T. Frankenberg Jr., S. Benson (red.) Marcel Dekker Inc., New York, 29–45.
- McCann J.C., Ames B.N., 2011. Adaptive dysfunction of selenoproteins from the perspective of the triage theory: why modest selenium deficiency may increase risk of diseases of aging. *FASEB J.* 25, 1793–1814.
- McComb T., Bischoff K., Thompson B., Smith M.C., Mohammed H.O., Ebel J., Hillebrandt J., 2010. An investigation of blood selenium concentrations of goats in New York State. *J. Vet. Diagn. Invest.* 22(5), 696–701.
- Morand-Fehr P., 1981. Nutrition and feeding of goat – application to temperature climatic condition. In: Gall. C. (Ed.), *Goat Production*. Academic Press, London, UK, 193–232.
- Musik I., Starosławska E., Pasternak K., 2003. Effect of organic selenium compounds on the activity of glutathione peroxidase and superoxide dismutase in selected mouse tissues. *Bull. Vet. Inst. Pulawy*, 47, 567–573.
- Navarro-Alarcon L., Lopez de la Serrana H., Perez-Valero V., López-Martínez C., 1998. Serum selenium levels in indicators of body status in cancer patients and their relationship with other nutritional and biochemical markers. *Sci. Total Environ.* 212, 195–202.
- Nowakowska E., 2013. Zawartość selenu w wybranych narządach (wątroba, nerki) zwierząt wolno żyjących (sarna, jeleń, dzik) jako kryterium oceny zasobności środowiska w ten pierwiastek na terenie Polski [Selenium level (in liver and kidneys) of free-living animals (roe deer, red deer and wild boar) as a criterion for assessment of this element content in the environment in Poland]. Praca doktorska [Doctoral thesis]. WBiHZ w Szczecinie [in Polish].
- Oh S., Sunde R., Pope A., Hoekstra W., 1976. Glutathione peroxidase response to selenium intake in lambs fed a torulabased, artificial milk. *J. Anim. Sci.* 42, 977–983.
- Pamukcu T., Sel T., Yarim G., 2001. Blood Serum Concentrations of Selenium and Glutathione Peroxidase Activity in Akkaraman Sheep. *Turk. J. Vet. Anim. Sci.* 25, 731–734.

- Pârvu Gh., Costea M., Pârvu M., Nicolae B., 2003. Tratat de nutriție a animalelor, Editura Coral Sanivet, București. Pastoral Agricultural Research Institute, Palmerston North, New Zealand 9–23.
- Pavlata L., Illek J., Pechová A., Matújíáek M., 2002. Selenium Status of Cattle in the Czech Republic. *Acta Vet. Brno* 71, 3–8
- Pechowa A., Sevcikova L., Pavlata L., Dworak R., 2012. The effect of various forms of selenium supplied to pregnant goats on selected blood parameters and on the concentration of Se in urine and blood of kids at the time of weaning. *Veterinarni Medicina* 57(8), 394–403.
- Pięta M., Patkowski K., 2007. Skład mineralny wełny i surowicy krwi owiec rasy świńiarka jako wskaźnik oceny środowiska [Mineral composition of wool and blood serum of Świniańska sheep breed as an indice of environment evaluation]. *Annales Universitatis Marie Curie-Skłodowska Lublin – Polonia 2, Sectio EE*, 71–77 [in Polish].
- Pilarczyk B., Udała J., Stankiewicz T., Hendzel D., Tylkowska A., Vovk S., 2011 a. Beurteilung der Selengehalte im Blutserum von Pferden in Polen. *Tierärztl. Umschau* 66, 163–165.
- Pilarczyk B., Balicka-Ramisz A., Ramisz A., Adamowicz E., Bujak T., Tomza-Marciniak A., Bąkowska M., Dąbrowska-Wieczorek M., 2008 a. Selenium concentration in roe deer from the Western Pomerania, Poland. *Bull. Vet. Inst. Pulawy* 52, 631–633.
- Pilarczyk B., Balicka-Ramisz A., Ramisz A., Adamowicz E., Pilarczyk R., Tomza-Marciniak A., Bąkowska M., 2009. Selenium concentration in liver and kidney of free living animals (roe and red deer) from West Pomerania (Poland). *Eur. J. Wildl. Res.* 55, 279–283.
- Pilarczyk B., Balicka-Ramisz A., Ramisz A., Pilarczyk R., Sablik P., Wójcik J., 2008 b. Selengehalte im Blutserum von aus Deutschland, Tschechien, Frankreich und Holland nach Polen importierten Rindern. *Tierärztl. Umschau* 63, 243–246.
- Pilarczyk B., Balicka-Ramisz A., Ramisz A., Vovk S., Tomza A., Wieczorek M., 2005. Badania nad poziomem selenu w surowicy zwierząt gospodarskich pochodzących z terenu Pomorza Zachodniego i zachodniej Ukrainy [The investigation of serum selenium levels in farm animals from Western Pomerania and Western Ukraine]. *Folia Univ. Stein. Zootechnica* 243 (47), 137–142 [in Polish].
- Pilarczyk B., Balicka-Ramisz A., Ramisz A., Vovk S., Major D., Jastrzębski G., Cisek A., 2004. Wpływ podawania selenu na jego poziom w surowicy krwi oraz wybrane wskaźniki użytkowości krów, świń i owiec [Effect on selenium supplementation on serum Se levels and selected of performance parameters in cows, pigs and sheep]. *Folia Univ. Agric. Stetin., Zootechnica* 235 (46), 53–58. [in Polish].
- Pilarczyk B., Balicka-Ramisz A., Ramisz A., Vovk S., Vantukh A., Bąkowska M., Holowacz I., Templin E., 2010. Selengehalt im Serum von Hunden und Katzen in Westpommern und der Westukraine. *Tierärztl. Prax Kleintiere* 38 (6), 374–378.
- Pilarczyk B., Jankowiak D., Tomza-Marciniak A., Pilarczyk R., Sablik P., Drozd R., Tylkowska A., Skólmowska M., 2012. Selenium concentration and glutathione peroxidase (GSH-Px) activity in serum of cows at different stages of lactation. *Biol. Trace Elem. Res.* 147, 91–96.

- Pilarczyk B., Majewska D., Tomza-Marciniak A., Tylkowska A., Ligocki M., Rząd I., Majewski J., 2011 b. Beurteilung der Selenkonzentration bei Straußen in Polen. Tierärztl. Umschau 66 (11), 461–466.
- Pilarczyk B., Pilarczyk R., Tomza-Marciniak A., Kowieska A., Wójcik J., Sablik P., Tylkowska A., Hendzel D., 2011 c. Selenkonzentration im Blutserum und in der Milch von Kühen aus einem ökologischen Betrieb und aus konventionellen Betrieben in einem Selenmangelgebiet. Tierärztl. Umschau 66(6), 248–252.
- Pilarczyk B., Tomza-Marciniak A., Dobrzański Z., Szewczuk M., Stankiewicz T., Gączarzewicz D., Lachowski W., 2013 a. The effect of selenized yeast supplementation on some performance parameters in sheep. Turk. J. Vet. Anim. Sci. 37, 61–67.
- Pilarczyk B., Tomza-Marciniak A., Pilarczyk R., Kuba J., Mituniewicz-Małek A., Dmytrow I., Stankiewicz T., Udała J., 2013 b. Selengehalt im Fleisch und in ausgewählten Organen von Ziegenlämmern sowie in der Ziegenmilch und ihren Produkten aus dem Gebiet Westpommern. Züchtungskunde 85 (6), 451–461.
- Pilarczyk R., Wójcik J., Czerniak P., Sablik P., Pilarczyk B., Tomza-Marciniak A., 2013 c. Concentrations of toxic heavy metals and trace elements in raw milk of Simmental and Holstein–Friesian cows from organic farm. Environmental Monitoring and Assessment 185, 8383–8392.
- Pop I.M., Radu-Rusu C., Simeanu D., Pop C., Fleancu N., 2011. Effects induced by a source of organic selenium and e vitamin usage in dairy cattle feeding. Lucrări Științifice 55, Zootehnie 160–164.
- Prasad T., Arora S.P., 1991. Influence of different sources of injected selenium on certain enzymem, glutatione and adenosylmethionine concentration In Buffalo (*Bubalus bubalis*) calves. British Journal of Nutrition 66, 261–267.
- Puls R., 1994. Mineral levels in animal health: diagnostic data, 2nd edn. Sherpa International, Clearbrook, 356.
- Radostits O.M., Gay C.C., Blood D.C., Hinchcliff K.W., 2000. Veterinary Medicine: A textbook of the diseases of cattle, sheep, pigs, goats and horses. 9 ed. London, Harcourt Th Publishers Ltd.1877.
- Ramadan A.A., Ghoniem A.A., Hassan H.M., Youssef A.E., 2001. Effects of beta-carotene, selenium and vitamin A on in vitro polymorphonuclear leukocytic activity in peripartal buffalo (*Bubalus bubalis*). Theriogenology 55 (3), 693–704.
- Ramírez-Bribiesca J.E., Tortora J.L., Huerta M., Aguirre A., Hernandez L.M., 2001. Diagnosis of selenium status in grazing dairy goats on the Mexican plateau. Small Ruminant Research 41, 81–85.
- Rayman M.P., 2000. The importance of selenium to human health. Lancet, 356, 233–241.
- Rederstorff M., Krol A., Lescure A., 2006. Understanding the importance of selenium and selenoproteins in muscle function. Cell Mol Life Sci. 63, 52–59.
- Ren Y., Wang Q., Shi L., Yue W., Zhang C., Lei F., 2011. Effects of maternal and dietary selenium (Seenriched yeast) on the expression of p34cdc2 and CyclinB1 of germ cells of their offspring in goats. Anim. Reprod. Sci. 123, 187–191.
- Sablik P., Kobak P., Pilarczyk B., Szarkowski K., Syczewski A., 2011. Serum selenium concentrations in water buffaloes (*Bubalus bubalis*) and Black-and-White German Holstein-Friesian cows raised in the Notec River region of the Wielkopolska province. Bull. Vet. Inst. Pulawy 55, 141–144.

- Segerson E.C., Riviere G.J., Dalton H.L., Whitacre M.D., 1981. Retained Placenta of Holstein Cows Treated with Selenium and Vitamin E. *J. Dairy Sci.* 64, 1833–1836.
- Segerson E., Gunsett F., Getz W., 1986. Selenium-vitamin E supplementation and production efficiency in ewes marginally deficient in selenium. *Livest. Prod. Sci.* 14, 149–159.
- Seremak B., Pilarczyk B., Tomza-Marciniak A., Pilarczyk R., Jakubczak A., Dziadosz M., Pławski K., Hendzel D., 2011. Assessment of selenium concentration in selected organs of farmer raccoon dogs (*Nyctereutes procyonoides*). *Ann. Anim. Sci.* 11 (4), 519–527.
- Slavík P., Illek J., Zelený T., 2007. Selenium Status in Heifers, Late Pregnancy Cows and Their Calves in the Šumava Region, Czech Republic. *Acta Vet. Brno* 76, 519–524.
- Słupczyńska M., Kinal S., Hadryś M., Król B., 2009. Utilization of selenium compounds in nutrition of lambs. *J. Elementol.* 14 (1), 157–164.
- Smith K.L., Weiss W.P., Hogan J.S., 1998. Influence of vitamin e and selenium on mastitis and milk quality in dairy cows. Texas Animal Nutrition Council (<http://txanc.org/wp-content/uploads/2011/08/vitamine.pdf>), date of access: 31.12.2014.
- Stuttle N.F., 2010. Selenium. In: Mineral nutrition of livestock, 4rd Ed. CABI Publ, Oxon 2010, UK. 377–425.
- Terzano G. M., Allegrini S., Borghese A., Roncoroni C. Alfieri L., 2005. Metabolic and hormonal parameters In buffaloes. Buffalo production and research. Roma 219–249.
- Tinggi U., 2005. Selenium toxicity and its adverse health effects. Wyd: Taylor & Francis, Boca Raton, Preedy R. and Watson R.R. Reviews in Food and Nutrition Toxicity. FL, 29–55.
- Villar D., Arthur J.R., Gonzalez J.M., Pallares F.J. Carson T.L., 2002. Selenium status in cattle: interpretation of laboratory results. *Bov. Pract.* 36, 73–80.
- Wesołowski M., Ulewicz B., 2000. Selen – pierwiastek śladowy niezbędny dla człowieka, występowanie, znaczenie biologiczne i toksyczność [Selenium – a trace element essential for human, occurrence, biological importance and toxicity]. *Farm. Polska* 56, 1004–1019 [in Polish].
- World Health Organization, 1987. Environmental Health Criterion 58–Selenium, World Health Organization, Geneva.
- Yläränta T., 1990. The selenium content of some agricultural crops and soils before and after the addition of selenium to fertilizers in Finland. *Ann. Agric. Fenn.* 29, 131–139.
- Zust J., Hrovatin B., Simundic B., 1996. Assessment of selenium and vitamin E deficiencies in dairy herds and clinical disease in calves. *Vet. Rec.* 16, 391–394.

SELEN W ŻYWIEŃIU PRZEŻUWACZY

Streszczenie. Liczne wyniki badań wskazują, że na większości obszarów istnieją niedobory selenu w środowisku. Wskaźnikiem zaopatrzenia organizmu w selen może być jego stężenie w surowicy lub krwi (przyżyciowo), jak i również jego zawartość w wątrobie oraz nerkach (pośmiertnie). Najczęściej stosowanym kryterium oceny stanu selenowego u zwierząt gospodarskich jest jego zawartość w surowicy lub krwi. Ważne jest, aby uzupełniać niedobór selenu u zwierząt bytujących w regionach niedoborowych, zwłaszcza u przeżuwaczy, gdyż długotrwały deficyt tego pierwiastka może doprowadzić do licznych schorzeń.

Słowa kluczowe: choroby, przeżuwacze, selen

Accepted for print – Zaakceptowano do druku: 23.12.2014