

Article ID: 174414
DOI: 10.5586/am/174414

Publication History
Received: 2023-09-21
Accepted: 2023-10-23
Published: 2023-11-23

Handling Editor
Katarzyna Patejuk; Wrocław
University of Environmental and
Life Sciences Wrocław, Poland;
<https://orcid.org/0000-0001-7236-8005>

Authors' Contributions
US: research concept and
design, data analysis and
interpretation, writing the entire
manuscript, description of the
specimen, photographic
documentation, and figure; RR:
a collection of material in the
field, manuscript revising; AW:
identification of the specimen.



Funding
The study was supported by the
Institute of Biology and
Biotechnology, Maria
Curie-Skłodowska University in
Lublin through its statutory
funds.

Competing Interests
No competing interests have
been declared.

Copyright Notice
© The Author(s) 2023. This is an
open access article distributed
under the terms of the [Creative
Commons Attribution License](https://creativecommons.org/licenses/by/4.0/),
which permits redistribution,
commercial and
noncommercial, provided that
the article is properly cited.

ORIGINAL RESEARCH PAPER

Puccinia scirpi – First finding of the fungus in the uredinial stage and new notes of the aecial and telial stages in Poland

Urszula Świdarska ^{1*}, Robert Rozwałka ², Agata Wołczańska^{1†}

¹Department of Botany, Mycology and Ecology, Maria Curie Skłodowska University in Lublin, Akademicka 19, 20-033 Lublin, Poland

²Department of Zoology and Animal Ecology, University of Life Sciences in Lublin, Akademicka 13, 20-033 Lublin, Poland

* To whom correspondence should be addressed. Email: urszula.swidarska@mail.umcs.pl

† Deceased.

Abstract

Puccinia scirpi causing the flowering rust disease on *Nymphoides peltata* and *Schoenoplectus lacustris* was recorded in 2018 from Bolestraszyce Arboretum near Przemyśl (Subcarpathian Voivodeship, Poland). The occurrence of the heteroecious rust species in the aecial (stage I), uredinial (stage II), and telial (stage III) stages are reported, and morphological descriptions, illustrations, and microphotographs are shown in the current study. This is the second report on the fungus in the aecial stage since the end of the 19th century, the first report on the uredinial stage, and the ninth description of the telial stage in Poland.

Keywords

rust fungus; Pucciniales; *Nymphoides*; *Schoenoplectus*; phytopathogen; microfungus

1. Introduction

Rust fungi are pathogens causing diseases in plants cultivated and used by humans. As biotrophs, rusts are specially adapted to obtain nutrients from living plant cells. The name of these fungi is derived from the rusty color of the so-called summer spores (urediniospores). They are highly specialized pathogens with the ability to complete the entire morphologically and cytologically complicated life cycle on one (autoecious) or two (heteroecious) hosts. As pleomorphic fungi, they have the ability to form up to five different successive morphological stages of sporulation (basidiospores, spermatia, aeciospores, urediniospores, and teliospores) in a specific order (Majewski, 1979; Zhao et al., 2023). The Pucciniales order comprises one of the largest groups of phytopathogens within both the Fungi kingdom and rusts, which occur on mosses, ferns, and advanced monocots and dicots (Aime & McTaggart, 2021; Kirk et al., 2008; Zhao et al., 2023). Over half of the known rust species represent the *Puccinia* Pers. genus. The genus comprises approx. 4,000

species common in the world, occurring almost exclusively on angiosperms (Kirk et al., 2008).

Puccinia scirpi DC. (common club rush rust) was described in 1805 by Lamarck and de Candolle under this name and as *Aecidium nymphoides* DC., now considered a synonym of the previous name. Currently, the fungus *Dicaeoma scirpi* (DC.) Gray, introduced in 1821, and *Aecidium nymphaeae* Wallr. described as a new species in 1833, are regarded as *P. scirpi* (Index Fungorum, 2023). The fungus is macrocyclic and heteroecious rust noted on representatives of the Menyanthaceae and Cyperaceae families (Gäumann, 1959).

P. scirpi is a rare species in Poland. It was recorded on the first host, *N. peltate*, from only one locality over 100 years ago and has not been reported since then. In the telial stage, it was recorded at the same time from six localities (Majewski, 1979), one locality in 2012 (Czerniawska et al., 2012), and one in 2017 (locality from 2009) (Mazurkiewicz-Zapałowicz et al., 2017). To date, it has never been found in Poland in the uredinial stage; therefore, this is the first report of the fungus.

The *Nymphoides* genus (Menyanthaceae) comprises species of aquatic plants. The geographical range of *Nymphoides peltata* (S.G. Gmel.) Kuntze covers Europe from the Baltic States to the Iberian Peninsula and Asia from the Middle East through northern India, Siberia, Mongolia to China, Japan, and the Korean Peninsula. The species was also adventive to North America, where it spread (Lansdown, 2014; Ławicki & Marchowski, 2019). Throughout Europe, the plant is quite common in the valleys of large southern European rivers, but the number of populations is unknown. In some countries or their regions, the species is rare or very rare, while in several countries, it is classified as threatened on several national red lists (Lansdown, 2014). In Poland, the fringed water lily is a scarce species under legal protection and has the status of an endangered species. The plant occurs along rivers and estuaries. In 2001, the localities were not very numerous; at that time, there were about 45 existing, 40 declining, and several unconfirmed localities, but many of these no longer exist (Kłosowski, 2014; Zając & Zając, 2001). It inhabits mainly oxbow lakes, river bends, eutrophic lakes, and fish ponds (Ławicki & Marchowski, 2019).

The genus *Schoenoplectus* (Rchb.) Palla (Cyperaceae) is represented by over 70 species and subspecies of hydrophytes and helophytes (Mazurkiewicz-Zapałowicz et al., 2017; Smith, 2002). In Poland, four species of *Schoenoplectus* occur naturally, the most common of which is *S. lacustris* (L.) Palla (Mirek et al., 2002). *S. lacustris* is a common species throughout Poland (Zając & Zając, 2001).

Summarising the occurrence of both hosts of *P. scirpi* in Poland, it can be expected that, due to the occurrence of both plants close to each other, rust should occur slightly more frequently in the country; however, the sporadic occurrence of *N. peltata* reduces the incidence of rust. The fungus can complete its entire life cycle only in the presence of both hosts.

2. Material and methods

Infected specimens of *N. peltata* and *S. lacustris* were collected in 2018 in Bolestraszyce Arboretum near Przemyśl in Poland. Microscope slides were prepared from air-dried specimens. Aeciospores, teliospores, and urediniospores were stained with cotton blue in lactic acid, heated, and observed under a light microscope. Microphotographs of fungal diagnostic structures of the species were taken with an Olympus digital camera SC180 and an Olympus BX53 light microscope.

Monographs by Majewski (1979) and Gäumann (1959) were used for rust identification. The nomenclature of the host plants follows the World Flora Online (WFO, 2023). The collected specimens are deposited in the herbarium of Maria Curie-Skłodowska University in Lublin (LBL).

3. Results

Puccinia scirpi, the heteroecious rust species in the aecial (stage I) on *N. peltata* and uredinial (stage II), and telial (stage III) stages on *S. lacustris* were collected in 2018 in Poland from Bolestraszyce Arboretum near Przemyśl.

3.1. Taxonomy

Puccinia scirpi DC., in Lamarck & de Candolle, Fl. franç., Edn 3 (Paris) 2: 223 (1805) (Figure 1)

Synonyms:

Aecidium nymphoidis DC., in Lamarck & de Candolle, Fl. franç., Edn 3 (Paris) 2: 597 (1805)

Dicaeoma scirpii (DC.) Gray, Nat. Arr. Brit. Pl. (London) 1: 542 (1821)

Aecidium nymphaeae Wallr., Fl. crypt. Germ. (Norimbergae) 2: 255 (1833)

Description:—*Aecia* epiphyllous, cup-shaped, up to 0.5 µm in diameter. *Peridium* cup-shaped, with a folded margin; peridium cells in regular rows, external wall up to 7 µm thick, surface densely and coarsely verrucose, inner wall up to 4 µm thick, surface densely and finely verruculose, 22–34 × 17–23 µm. *Aeciospores* oblate, catenulate, 15–20 × 19–23 µm, wall up to 1.4 µm thick, usually verruculose. *Uredinia* elongated, up to 0.5 mm long, long covered by a vesicularly raised epidermis. *Urediniospores* irregularly ovate or elliptical, 20–25(–28) × 14–21 µm; yellowish, 1.5–2 µm thick, covered by fine spikes. Germ pores 2, ± equatorial. *Telia* elongated, up to 2 mm long, covered by a persistent brown epidermis, after tearing, they are black, dust-free. *Teliospores* (sub-)clavate, usually rounded at the apex, less often irregularly flattened or pointed, tapering at the base, constricted at the septum, 33–58 × 13–23 µm. Wall brown-yellow, smooth, thickened in the distal cell up to 7–10 µm. Germ pore indistinct. Pedicel/stalk up to 62 µm long, persistent, separated from the base of the teliospores by a diagonal wall.

Examined collections:—On *Nymphoides peltata* (S.G. Gmel.) Kuntze (Menyanthaceae), stage I. POLAND. SUBCARPATHIAN VOIVODESHIP: Bolestraszyce Arboretum n. Przemyśl, 2 June 2018, leg. R. Rozwałka, det. A. Wołczańska (LBL M–33129). On *Schoenoplectus lacustris* (L.) Palla (Cyperaceae), stage II, III. POLAND. SUBCARPATHIAN VOIVODESHIP: Bolestraszyce Arboretum n. Przemyśl, 2 June 2018 (last year's shoots), leg. R. Rozwałka, det. A. Wołczańska (LBL M–33127); 10 August 2018, leg. R. Rozwałka, det. A. Wołczańska (LBL M–33128); 10 October 2018, leg. R. Rozwałka, det. A. Wołczańska (LBL M–33126).

Literature data:—On *N. peltata*, *Limnanthemum nymphoides* (hosts need revision). POLAND. Motława River in Gdańsk (Majewski, 1979). On *S. lacustris*. POLAND. Wolin Island, Kołobrzeg, near Legnica and Ścinawa, vicinity of Cracow and Międzyrzec Podlaski (Majewski, 1979); Wasosze Lake (West Pomerian province) (Czerniawska et al., 2012); Płociczno Lake (Mazurkiewicz-Zapałowicz et al., 2017).

General distribution:— Europe, North Africa; Asia, North America, New Zealand.

4. Discussion

P. scirpi is a dioecious and heteroecious rust, usually with members of *Nymphoides* as aecial hosts and representatives of *Schoenoplectus* (= *Scripus*) as uredinial and telial hosts. The fungus has also been noted in all stages of sporulation on representatives from a few other host genera (Farr & Rossman, 2021).




Figure 1 *Puccinia scirpi* (A) and (C) aecia with aeciospores on the leaf of *N. peltata* (LBL M–33129); (B) stem of *S. lacustris* with telia (LBL M–33126); (D) aeciospore (LBL M–33129); (E) urediniospores (LBL M–33128); (F) teliospores (LBL M–33127). Scale bars: (A) 0.2 mm; (B) 0.5 mm; (C) 100 μ m; (D,F) 20 μ m; (E) 10 μ m. Photographs: (A), (C–F) U. Świdwerska; (B) M. Szewczyk.

On *N. peltata*, the rust fungus has been reported from Bulgaria, Lithuania, Norway, and the United Kingdom, in addition to Poland. On the members of the *Nymphoides* genus, the fungus has also been reported worldwide on *N. forbesii* (probably *N. forbesiana*) and *N. indica* subsp. *occidentalis* (both from Uganda), *N. grayana* (from Cuba), and *Nymphoides* sp. (from the USA – Florida) (Farr & Rossman, 2021; Soares et al., 2006). The fungus has been noted on *S. lacustris* in Bulgaria, Denmark, France, Germany, Greece, Lithuania, Norway, Poland, Romania, Spain, Sweden, the United Kingdom, Ukraine, Puerto Rico, China, and Iran (Farr & Rossman, 2021). It is known to infect a few other *Schoenoplectus* species, i.e., *S. litoralis* (India, Iran, and Pakistan), *S. tabernaemontani* (the Dominican Republic, Japan, Lithuania, New Zealand, Puerto Rico, Thailand, the Virgin Islands, and the West Indies), *S. triquetter* (China, Japan, Taiwan), and *S. validus*

(New Zealand and Thailand) (Farr & Rossman, 2021; Soares et al., 2006; Wu et al., 1982).

The fungus has been reported from eight localities in Poland (Wolin Island, Kołobrzeg, Jakuszów near Legnica, Orzeszków near Ścinawa, surroundings of Cracow and Międzyrzec Podlaski, Wasosze Lake, and Płociczno Lake) on *S. lacustris* in the telial stage (stage III) (Czerniawska et al., 2012; Majewski, 1979; Mazurkiewicz-Zapałowicz et al., 2017). In the uredinial stage (stage II), the fungus has not been published from Poland to date. In the aecial stage (stage I), the species has been noted only in Gdańsk (on the Motława river) (Majewski, 1979), and this is the second report in this stage since the end of the 19th century in Poland. The rare occurrence of *N. peltata* may be associated with the sporadic, although the dispersed occurrence of this host in its natural habitats (Gdaniec, 2010; Mazurkiewicz-Zapałowicz et al., 2017).

Acknowledgments

The authors thank the management and staff of the Arboretum in Bolestraszyce for their cooperation and assistance in collecting the material. The authors would also like to thank Dr. Marian Szewczyk for taking the photo of *Schoenoplectus lacustris* with visible telia.

References

- Aime, M. C., & McTaggart, A. R. (2021). A higher-rank classification for rust fungi, with notes on genera. *Fungal Systematics and Evolution*, 7, 21–47. <https://doi.org/10.3114/fuse.2021.07.02>
- Czerniawska, B., Adamska, I., & Dziegielewska, M. (2012). Fungal diseases on the reed-bed vegetation of the eutrophic Wasosze lake. *Ecological Chemistry and Engineering A*, 19(12), 1547–1553. [https://doi.org/10.2428/ecea.2012.19\(12\)148](https://doi.org/10.2428/ecea.2012.19(12)148)
- Farr, D. F., & Rossman, A. Y. (2021). *Fungal databases, systematic mycology and microbiology Laboratory, ARS, USDA*. Retrieved May 6, 2021 from <http://nt.ars-grin.gov/fungaldatabases/>
- Gäumann, E. (1959). *Die Rostpilze Mitteleuropas*. Beiträge zur Kryptogamenflora der Schweiz. Vol. 12. Buchdruckerei Buehler & Co.
- Gdaniec, M. (2010). Nowe stanowisko *Nymphoides peltata* (S. G. Gmel.) Kuntze w Borach Tucholskich [New locality of *Nymphoides peltata* (S. G. Gmel.) Kuntze in the Tuchola Forest]. *Acta Botanica Cassubica*, 7(9), 243–245.
- Index Fungorum. (2023). Retrieved April 12, 2023 from <http://www.indexfungorum.org/>
- Kirk, P. M., Cannon, P., Stalpers, J., & Minter, D. W. (2008). *Dictionary of the fungi* (10th ed.). CABI Publishing.
- Kłosowski, S. (2014). *Nymphoides peltata* (S. G. Gmel.) Kuntze. Grzybieńczyk wodny. In R. Kaźmierczakowa, K. Zarzycki, & Z. Mirek (Eds.), *Polish red data book of plants: Pteridophytes and flowering plants* (3rd edition revised and expanded, pp. 402–404). Institute of Nature Conservation PAS.
- Lansdown, R. V. (2014). The IUCN Red List of Threatened Species 2014: e.T164309A42398483. *Nymphoides peltata*. <https://doi.org/10.2305/IUCN.UK.2014-1.RLTS.T164309A42398483.en>
- Ławicki, Ł., & Marchowski, D. (2019). Distribution and prospects for protection of the fringed waterlily *Nymphoides peltata* in the Lower Odra river valley. *Przegląd Przyrodniczy*, 30(2), 33–42.
- Majewski, T. (1979). Podstawczaki (Basidiomycetes), rdzawnikowe (Uredinales) II. In J. Kochman & A. Skirgiełło (Eds.), *Flora polska, Grzyby (Mycota)* (Vol. 11). Państwowe Wydawnictwo Naukowe.
- Mazurkiewicz-Zapałowicz, K., Rybińska, A., & Łopusiewicz, Ł. (2017). Microscopic fungi on *Schoenoplectus lacustris* in Płociczno and Płociowe lakes in Drawa National Park (northwest Poland). *Acta Mycologica*, 52(2), Article 1099. <https://doi.org/10.5586/am.1099>
- Mirek, Z., Piękoś-Mirkowa, H., Zając, A., & Zając, M. (2002). *Flowering plants and pteridophytes of Poland. A checklist*. W. Szafer Institute of Botany, Polish Academy of Sciences.
- Smith, S. G. (2002). *Schoenoplectus*. In Flora of North America Editorial Committee (Eds.), *Flora of North America: North of Mexico. Vol. 23: Magnoliophyta: Commelinidae (in part): Cyperaceae* (pp. 44–60). Oxford University Press.
- Soares, D. J., Ferreira, F. A., & Barreto, R. W. (2006). First report of the aecial stage of *Puccinia scirpi* on *Nymphoides indica* in Brazil, with comments on its worldwide distribution. *Australasian Plant Pathology*, 35, 81–84. <https://doi.org/10.1071/AP05091>
- WFO. (2023). World Flora Online. Published on the Internet. Retrieved May 25, 2023 from <http://www.worldfloraonline.org>
- Wu, C. G., Tseng, H. Y., & Chen, Z. C. (1982). Fungi inhabiting on *Schoenoplectus triqueter* (L.) Palla (I). *Taiwania*, 27, 35–38.
- Zając, A., & Zając, M. (Eds.). (2001). *Distribution atlas of vascular plants in Poland*. Laboratory of Computer Chorology, Institute of Botany, Jagiellonian University.
- Zhao, P., Li, Y., Li, Y., Liu, F., Liang, J., Zhou, X., & Cai, L. (2023). Applying early divergent characters in higher rank taxonomy of *Melampsorineae* (Basidiomycota, *Pucciniales*). *Mycology*, 14(1), 11–36. <https://doi.org/10.1080/21501203.2022.2089262>