



World News of Natural Sciences

An International Scientific Journal

WNOFNS 24 (2019) 366-372

EISSN 2543-5426

Antibiotic Susceptibility Test of Bacteria Isolated From Fruit Juices Sold in Cafes and Restaurants of Debre-Markos Town, North Western Ethiopia

Kindu Geta^{1,*}, Ameha Kebede², Meseret Chemedissa²

¹Department of Biology, Faculty of Natural and Computational Sciences, Debre Tabor University, Debre Tabor, Po.box 272, Ethiopia

²Department of Biology, Faculty of Natural and Computational Sciences, Haramaya University, Harar, Po.box-138, Ethiopia

*E-mail address: kindu2012@gmail.com

ABSTRACT

The prevalence of antimicrobial resistance among food pathogens has increased during recent decades. In this work, pathogenic bacteria such as *Staphylococcus aureus*, *Salmonella* spp., *Shigella* spp., and *E. coli* were isolated following standard methods. The bacterial isolates were then tested for their sensitivity to common antibiotics using the disc diffusion method on Mueller-Hinton Agar. All of the pathogenic bacteria were found to be resistant to erythromycin and almost all were sensitive to penicillin.

Keywords: Antibiotics, Antimicrobial resistance, Debre Markos, Fruit juices, Pathogens, *Salmonella*, *Shigella*, *E. coli*, *Staphylococcus aureus*, *B. cereus*, *Klebsiella*, *Enterobacter*, *Ps. aeruginosa*

1. INTRODUCTION

The antimicrobial resistance of bacteria isolated from food and other sources, against commonly used antibiotics has increased from time to time (Vicas and Singh, 2010). Not only their presence, but also their resistance to the commonly used antibiotics has become a concern for consumers.

Some reports have revealed that antibiotic resistance levels are becoming elevated among food-borne pathogens such as *Salmonella* and *Shigella* (Mache, 2002). Although, it is difficult

to prove a direct role of drug resistance in bacteria contaminating food items with increased clinical cases of resistant infections, the presence of such bacteria in food items could play a role in the spread of antimicrobial resistance amongst food-borne pathogens (Farzana *et al.*, 2009). The incidence of resistant bacteria in foodstuff is a worldwide phenomenon. It is a major public health threat (Khan and Malik, 2001) as these organisms have been isolated from wide range of foodstuffs consumed by human.

The prevalence of antimicrobial resistance among food pathogens has increased during recent decades (Davis *et al.*, 1999; Garau *et al.*, 1999; Threfall *et al.*, 2000 and Chui *et al.*, 2002). Despite the extensive studies, the antibiotic sensitivity of these bacteria has not been well studied especially in a developing country. The aim of this research is therefore to test the antibiotic susceptibility of bacterial pathogens to some of the commonly prescribed antibiotics in the study area.

2. MATERIAL AND METHODS

Thirty-six samples of avocado and mango of locally prepared unpasteurized fruit juices were collected from six cafe or restaurant in Debre-Markos town from February 2014 to May 2014. 250 ml of juice samples were collected from cafes, in sterile beakers aseptically labeled, and immediately transported to Debre-Markos University Laboratory in an icebox where they were processed immediately.

Pathogenic bacteria such as *Salmonella*, *Shigella*, *E. coli*, *S. aureus*, *Klebsiella* spp and *B. cerues* were isolated according to the procedures outlined by Food and Drug Administration (FDA) (2001). For detecting, the presence of *Salmonella* and *Shigella*, 25 ml of juice sample was added into 225 ml of sterile peptone water and homogenized by shaking. The resulting dilution was then serially diluted up to 10^{-5} dilution. From the 10^{-1} and 10^{-2} dilutions, 1 ml was taken and inoculated into tubes of Lactose broth and incubated at 37 °C for 48 hours. A loopful of sample from each culture was then transferred to Rappaport Vassiliadis (RV) broth and incubated at 42 ± 0.2 °C for 24 hrs in water bath. Positive samples were confirmed by streaking on Hektoen Enteric Agar and then biochemically tested for the presence of *Salmonella* and *Shigella*. In all cases, for confirmation of the pathogens, typical colonies were identified based on cultural, microscopic and biochemical characteristics (Buchanan and Gibbons NE., 1974).

All isolates of pathogenic bacteria were tested for their sensitivity to antibiotics by means of the disc diffusion method on Mueller-Hinton Agar (Difco, Detroit, MI) as described previously by Bauer *et al.* (1966) using *E. coli* ATCC 25922 and *S. aureus* ATCC 25923 as standard obtained from Amhara Regional State Central Laboratory. All disks used in the disk diffusion test were obtained from BECTON, USA, in the following concentrations: Ampicillin (Amp 10 µg), Amoxicillin (Amx 10 µg), Gentamycin (Gm 10 µg), Chloramphenicol (C 30 µg), Streptomycin (S 10 µg), Penicillin G (P 5 Units), Tetracycline (Te 30 µg) and Erythromycin (Er 2 µg). Briefly, five colonies of each isolate were introduced into 5 ml of Nutrient broth, incubated for 4 hours, and the culture turbidity was adjusted to a 0.5 McFarland standard. Sterile cotton swab was dipped into the suspension and spread evenly over the entire Mueller-Hinton Agar surface.

The antibiotics impregnated discs were then placed onto the surface of the inoculated plates and incubated at 37 °C for 16-18 hrs. After incubation, diameters of the zones of inhibition were measured in mm and interpreted as susceptible, intermediate and resistant (CLSI, 2007).

3. RESULT AND DISCUSSION

According to current study, all the juice samples tested were devoid of *Salmonella* and *Shigella*. The prevalence of *Staphylococcus aureus* and *E. coli* was 6 (16.67%) and 4 (11.1%) out of 36 fruit juices, respectively (Table 1).

Table 1. Bacterial pathogens detected from avocado and mango fruits juices collected from Debere Markos town.

Isolates	Juice	N	Positive samples	Total	Percent	χ^2	P
<i>Salmonella</i>	Mango	18	0	0	0		
	Avocado	18	0				
<i>Shigella</i>	Mango	18	0	0	0		
	Avocado	18	0				
<i>E. coli</i>	Mango	18	1	4	11.1	1.125	.289
	Avocado	18	3				
<i>S. aureus</i>	Mango	18	1	6	16.7	3.200	.074
	Avocado	18	5				

Previous study in Bangladesh showed that *B. cereus*, *Salmonella* species and *Staphylococcus aureus* were found in 64.91%, 7.89% and 6.14%, of the tested samples respectively and (99%) the tested samples showed the presence of coliform and *E. coli* (Shakir *et al*, 2009). Another study conducted in India documented that 27.7%, 16.6%, 38.8% of fruit juices were positive for *E. coli*, *Shigella*, and *Salmonella* spp. positive respectively (Lewis *et al*, 2006).

According to the study conducted in Amravati city, India the incidence of bacterial pathogen recorded was *E.coli* (40%), followed by *Ps. aeruginosa* (25%), *Salmonella* spp. (16%), *Proteus* spp. (9%), *S. aureus* (6%), *Klebsiella* spp. (3%) and *Enterobacter* spp. (1%) in street vended fruit juices samples (Tambekar *et al*, 2009). Incidence of current study was differing from study conducted in Asian countries. The probable reason for difference may be attributed to fruit type, geographical variation, seasonal variation, sanitation habit and variation in methods of detection.

This study also tried to address antimicrobial susceptibility testing by means of a disc diffusion method on Mueller-Hinton Agar. The results of the antibiotic sensitivity test were interpreted and are presented as the resistant of bacterial isolates to the antibiotics (Table 2).

Most isolates were susceptible to Penicillin, Ampicilin, Gentamicine, and Chloramphenicol. All isolates were resistance to Erythromycin, and most isolates were resistance to Streptomycin, Amoxicillin, and Tetracycline. The antimicrobial resistance of

bacteria isolated from food and other sources, against commonly used antibiotics has increased from time to time (Vicas, 2010). According to the finding erythromycin was not active against all bacterial isolates. All isolates of *S.aureus* were resistance to erythromycin and amoxicillin. 16.7% of isolates were resistance to tetracycline and streptomycin. 33.3% and 66.7% to gentamicin and chloramphenicol respectively. All isolates were sensitive to penicillin and ampicillin. High rates of drug resistance were observed for *Staphylococcus* spp. against ampicillin (93%) and amoxicillin (92%) (Rashed, *et al.* 2013).

Table 2. Antimicrobial susceptibility patterns of pathogenic bacterial isolates from avocado and mango juice samples (in %).

Antibiotics	Bacterial isolates											
	<i>S.aureus</i> N = 6			<i>Klebsiella spp</i> N = 3			<i>E. coli</i> N = 4			<i>B. cereus</i> N = 2		
	R	I	S	R	I	S	R	I	S	R	I	S
Er	100	0	0	100	0	0	100	0	0	100	0	0
Amx	100	0	0	0	33.3	66.7	25	25	50	100	0	0
S	33.3	16.7	50	33.3	66.7	0	50	25	25	100	0	0
Te	16.7	0	83.3	66.7	33.3	0	75	25	0	50	50	0
C	66.7	16.7	0	0	0	100	25	0	75	50	0	50
Gm	33.3	33.3	33.3	0	0	100	25	0	75	50	0	50
P	0	0	100	66.7	33.3	0	0	0	100	0	0	100
Amp	0	16.7	83.3	66.7	0	33.3	50	25	25	50	0	50

Key: N = number of isolates, Er = Erythromycin, Amx = Amoxicillin, S = Streptomycin, Te = Tetracycline, C = Chloramphenicol, Gm = Gentamicin, P = Penicillin, Amp = Ampicillin.

Lateef (2004) reported that *S.aureus* were sensitive to erythromycin, gentamicin, and chloramphenicol was disagree with this finding. The works of Daniyan and Ajibo (2011) who reported that *S. aureus* was susceptible to streptomycin support this finding. Srinu *et al.*, (2012) also reported that *S. aureus* was sensitive to streptomycin.

66.7% of *Klebsiella spp.* isolates were resistant to tetracycline, penicillin, and ampicillin. 33.1% of isolates were resistance to streptomycin and all were sensitive to chloramphenicol and gentamicin. Rashed *et al.* (2013) reported that *Klebsiella spp.* showed higher resistance against ampicillin (74%) and amoxicillin (72%) resistance of *Klebsiella spp* against amoxicillin was disagree with this result. According to Stock and Wiedemann (2001), *Klebsiella spp.* were naturally sensitive or intermediate to several penicillins, all tested aminoglycosides, quinolones, tetracyclines, trimethoprim, cotrimoxazole, chloramphenicol and nitrofurantoin.

Some *E. coli* isolates were resistant to amoxicillin, chloramphenicol, and gentamicin. 50% were resistant to streptomycin and ampicillin; all isolates were sensitive to penicillin and 75% were resistance to tetracycline. A high level of resistance was obtained among the five *E. coli* strains. Amoxicillin were not active against the strains of *E. coli* (Lateef, A., 2004). Srinu *et al.* (2012) also reported that *E. coli* was sensitive to streptomycin. Osterbald *et al.* (1999) reported that *Escherichia* spp, *Klebsiella* sp shows high sensitivity to amoxicillin was in line with this study. Marwa *et al.* (2012) reported that most *E. coli* isolates from food were sensitive to amoxicillin was disagree with this result. All *B. cerues* isolates were resistance to streptomycin, erythromycin, and amoxicillin while penicillin was active against all isolates. Agwa *et al.* (2012) found that *B. cereus* was susceptible to erythromycin and streptomycin was disagree with this result. This result was agree with the work of Adesetan *et al.*, (2013) who reported that *B. cereus* was resistant to erythromycin.

4. CONCLUSION

The intensive and incorrect use of antimicrobial agents leads to the emergency of antimicrobial-resistant bacteria. Physicians should prescribe the correct antibiotics for the target bacteria and inform the correct use of antibiotics for patients.

Acknowledgements

I am thankful to Haramaya University School of Graduate Studies for providing me with financial support to conduct this project. I would like to thank Debre Markos University for their willingness to allow me to use the laboratory for conducting my experiments. My heartfelt thanks go to anonymous reviewers for their critical and constructive comment.

References

- [1] Adesetan, T.O., Egberongbe, H.O., Ilusanya, O.A.F., and Bello, O.O., 2013. Antimicrobial sensitivity of bacterial isolates from street vended fruits in Ijebu area of Ogun state, Nigeria. *Int. Res. J.Microbiol.* 4(9): 220-225
- [2] Agwa OK, Uzoigwe CI, Wokoma EC., 2012. Incidence and antibiotic sensitivity of *B. cereus* isolated from ready to eat foods sold in some markets in Port Harcourt, River State. *Asian. Jr. Microbiol. Biotech. Env. Sc.* 14(1): 13-18
- [3] Bauer, A.W., Kirby, W. M. M., Sherris, J. C. and Turck, M., 1966. Antibiotic susceptibility testing by a standard single disc diffusion method. *Am. J Clinic. Pathol.* 45, 493–496
- [4] Buchanan RE, Gibbons NE., 1974. *Bergey's Manual of Determinative Bacteriology*, Baltimore. Williams and Wilkins Co. 8th Edn. pp. 34-89.
- [5] Chui CH, Wu TL, Su LH, Chu C, Chia JH, Kuo AJ, Chien MS, Llin TY (2002). The emergence in Taiwan of Fluoroquinolone resistance in *Salmonella enterica* serotype cholerasuis. *N. Engl J. Med* 346: 416-419

- [6] CLSI. Performance Standards for Antimicrobial Susceptibility Testing. 27th ed. CLSI supplement M100. Wayne, PA: Clinical and Laboratory Standards Institute; 2017. Wayne, PA 19087 US. Print ISBN 1-56238-804-5
- [7] Daniyan SY, Ajibo CQ. 2011. Microbiological Examination of sliced fruits sold in Minna Metropolis. *Int. Res. J. Pharmacy* 2 (7): 124-129
- [8] Davis MA, Hancock DD, Besser TE, Rice DH, Gay JM, Gay C, Gearhart L, Difiacomo R (1999). Changes in antimicrobial resistance among *Salmonella enterica* serovar. *Infect. Dis.* 5: 802-806
- [9] Farzana, K., S. AK htar, and F. Jabeen. 2009. Prevalence and antibiotic resistance of some bacteria in two ethnic milk based products. *Pak. J. Bot.* 41: 935-943
- [10] FDA, 2001. In Bacteriological Analytical Manual Online, pp. 1–6. USA
- [11] Garau J, Xercavins M, Podriguez-Carballeira M, Gomez-vera JR, Coll I, Vidal D, Wovet T, Ruiz-Breman A (1999). Emergence and dissemination of quinolone resistant *Escherichia coli* in the community. *Antimicrob. Agents Chemother.* 43: 2736-2741
- [12] Khan MK, Malik A., 2001. Antibiotic resistance and detection of β -lactamase in bacterial strains of *Staphylococci* and *Escherichia coli* isolated from foodstuffs. *World J. Microbiol. Biotechnol.* 17: 863- 868
- [13] Lateef A., 2004. The Microbiology of a Pharmaceutical effluent and its Public health implications. *World J. Microbiol. Biotechnol.* 20: 167-171
- [14] Lewis Joy E, Thompson P, Rao B, Kalavati, C, Rajanna B., 2006. Human Bacteria in Street scale fruit juice processing. A Case Study of Visakhapatnam City, India. *Int. J. Food Safety*, 8, 35-38
- [15] Mache A., 2002. *Salmonella* serogroups and their antibiotic resistance patterns isolated from diarrhea stools of pediatric outpatients in Jimma Hospital and Jimma Health center, southwest Ethiopia. *Ethiop J Health Sci.* 12: 37-45
- [16] Marwa EAA, Tamer ME, Magdy AM., 2012. Antibiotic resistance profile of *E. coli* strains isolated from clinical specimens and food samples in Egypt. *Int. J. Microbiol. Res.* 3(3): 176-182
- [17] Osterbald M, Pensala O, Peterzens M, Heleniusc H, Huovinen P., 1999. Antimicrobial susceptibility of Enterobacteriaceae isolated from vegetable. *J. Antimicrobial Chemotherapy.* 43: 503-509
- [18] Rashed, N., Md. Aftab, U., Md. Azizul, H., Saurab, K. M., Mrityunjoy, A., and M. Majibur, R. 2013. Microbiological study of vendor and packed fruit juices locally available in Dhaka city, Bangladesh. *Int. Food Res. J.* 20(2): 1011-1015
- [19] Shakir M, Ahmed U, Nasreen T, Feroza B and Parveen S., 2009. Microbiological Quality of Local Market Vended Freshly Squeezed Fruit Juices in Dhaka City, Bangladesh. *Bangladesh J. Sci. Ind. Res.* 44, 421-424
- [20] Srinu B, Vijaya Kumar A, Kumar E, Madhava Rao T. (2012). Antimicrobial resistance of bacterial foodborne pathogens. *J. Chemical and Pharmaceutical Res.* 4(7): 3734-3736

- [21] Stock I, Wiedemann B (2001). Natural antibiotic susceptibility of *Klebsiella pneumoniae*, *K. oxytoca*, *K. planticola*, *K. ornithinolytica* and *K. terrigena* strains. *J. Med. Microbiol.* 50(5): 396-406
- [22] Tambekar, D.H., S.D. Shirsat, S.B.Suradkar, P.N. Rajankar, and Y.S. Banginwar, 2007. Prevention of transmission of infectious disease: Studies on hand hygiene in health-care among students. *Continental J. Biomedical Sciences* 1: 6-10
- [23] Threfall EJ, Ward LR, Frost JA and Willshaw GA (2000). The emergence and spread of antibiotic resistance in foodborne bacteria. *Int. J. Food Microbiol.* 62: 1-5.
- [24] Vicas M. Sanchaita S, Singh NP. 2010. Multidrug Resistant *Acinetobacter*. *J Glob Infect Dis.* 2: 291-304