



DIPLOID CHROMOSOME NUMBERS IN *HIERACIUM* AND *PILOSELLA* (ASTERACEAE) FROM MACEDONIA AND MONTENEGRO

ZBIGNIEW SZELĄG^{1*} AND TOMASZ ILNICKI²

¹*Institute of Botany, Jagiellonian University,
Kopernika 31, 31-501 Cracow, Poland*

²*Department of Plant Cytology and Embryology, Jagiellonian University,
Grodzka 52, 31-044 Cracow, Poland*

Received August 24, 2011; revision accepted November 13, 2011

The authors report the first discovery of diploid populations of *Hieracium naegelianum* Panč. subsp. *naegelianum* and *H. naegelianum* subsp. *lyubotenicum* Behr & Zahn., and give the first chromosome counts for *H. cernuum* Friv., *H. gymnocephalum* Griseb. ex Pant., *H. sparsum* Friv., *Pilosella pavichii* (Heuff.) Holub and *P. serbica* (F. W. Schultz & Schultz-Bip.) Szeląg from Macedonia and/or Montenegro. A diploid chromosome count for *Hieracium renatae* Szeląg is confirmed based on material from the whole distribution range of the species. An emasculation experiment showed that all the analyzed diploid *Hieracium* taxa reproduce sexually.

Key words: Asteraceae, Balkan Peninsula, chromosome number, *Hieracium*, karyotype, mode of reproduction, *Pilosella*.

In Europe the genera *Hieracium* L. and *Pilosella* Vaill. (*Hieracium* sensu Zahn 1921–1923) are dominated by polyploids (triploids to pentaploids in *Hieracium* and triploids to octoploids in *Pilosella*) with a basic chromosome number of $x = 9$. Sexual diploids are very rare and restricted mainly to refugial areas of Southern Europe such as the Balkan Peninsula and Iberian Peninsula (Merxmüller, 1975; Schuhwerk and Lippert, 1998; Vladimirov, 2003; Vladimirov and Szeląg, 2006; Chrtek et al., 2007, 2009). Looking for diploids, the first author carried out field research in the region of the former Yugoslavia, visiting sites including the highest massifs of Montenegro and Macedonia, the most refugial areas of mountain flora on the Balkan Peninsula (Horvat et al., 1974). During these studies many living plants of both genera were collected and analyzed karyologically. This paper presents the most significant results of this work focused on Balkan endemics.

Living plants were transplanted to an experimental garden of the first author, their chromosomes were counted at mitotic metaphase, and the mode of reproduction was tested with an emasculation experiment. Detailed information on the karyological procedure and experimental methods fol-

lowed here is given in Joachimiak et al. (1999) and Szeląg (2006).

Vouchers are stored in the herbarium of the first author (Herb. Hierac. Z. Szeląg) and the plants analyzed karyologically remain in cultivation in his garden.


Hieracium cernuum Friv. $2n = 2x = 18$, sexual
Macedonia, Nidže Mts., along road to summit of Mt. Kajmakčalan, 2300 m a.s.l., alpine grassland on siliceous bedrock.

This is the first chromosome number for this species from Macedonia; it confirms the count recently published from Bulgaria (Ilnicki and Szeląg, 2011).

Hieracium gymnocephalum Griseb. ex Pant. $2n = 2x = 18$, sexual
Montenegro, Prokletije Mts., Mt. Maja Rosave, 1650 m a.s.l., calcareous grassland on western slope above Zastan Koliba military checkpoint.

This is the first diploid chromosome number for the species from Montenegro; it confirms the recently published count from Albania (Chrtek et al., 2009). A triploid chromosome number was reported from the Durmitor Mts. in Montenegro (Niketić et al., 2006).

* e-mail: azszelag@wp.pl



Figs. 1–4. Metaphase plates of: **Fig. 1.** *Hieracium naegelianum* Panč. subsp. *naegelianum*, $2n = 18$. **Fig. 2.** *Hieracium naegelianum* subsp. *ljubotenicum* Behr & Zahn, $2n = 18$. **Fig. 3.** *Hieracium renatae* Szelag, $2n = 18$. **Fig. 4.** *Pilosella serbica* (F. W. Schultz & Schultz-Bip.) Szelag, $2n = 18$. Bar = 10 μm .

Hieracium naegelianum Panč. subsp. *naegelianum*
 $2n = 2x = 18$, sexual (Fig. 1)

1. Macedonia: Šarplanina Mts., Mt. Ljuboten, 2350 m a.s.l., calcareous rock crevices.
2. Macedonia, Korab Mts., between Mt. Šila Vort and Mt. Ništrovski Korab, 2050 m a.s.l., calcareous scree.

This is the first diploid chromosome number for this species, previously known only from triploid populations on the Balkan Peninsula (Merxmüller, 1975; Grau and Erben, 1988; Buttler, 1991; Vladimirov and Szelag, 2001a; Niketić et al., 2003, 2006; Chrtek et al., 2007), including plants from the *locus classicus* on Mt. Kom Vasojevički in Montenegro (Ilnicki and Szelag, 2011). Until now the only known diploid in *Hieracium* sect. *Naegeliania* Zahn ex Szelag was *H. renatae* Szelag described from Macedonia (see below).

Hieracium naegelianum subsp. *ljubotenicum* Behr & Zahn $2n = 2x = 18$, sexual (Fig. 2)
Macedonia, Šarplanina Mts., Mt. Ljuboten 2300 m a.s.l., calcareous rock crevices.

This is the first chromosome number for this taxon known only from Mt. Ljuboten.

Hieracium renatae Szelag $2n = 2x = 18$, sexual (Fig. 3)
Macedonia, Jakupica Mts.: 1. Mt. Solunska Glava NW slope, 2400 m a.s.l. (*locus classicus*); 2. above Gorno Begovo plateau, rocky slope along road to summit of Mt. Solunska Glava, 2200 m a.s.l.; 3. on ridge between Mt. Solunska Glava and Grob pass, 2250 m a.s.l.; 4. on ridge between Mt. Solunska Glava and Mt. Pržal, 2370 m a.s.l.; 5. summit region of Mt. Solunska Glava above Nežilovska Stena cliff, 2520 m a.s.l.

The present count is based on abundant material from the whole known range of *H. renatae* and confirms the diploid chromosome number published in the protologue of the species (Szelag, 2010).

Hieracium sparsum Friv. $2n = 2x = 18$, sexual
Macedonia, Nidže Mts., Redir ridge along road to summit of Mt. Kajmakčalan, 1650 m a.s.l., *Picea abies* forest margin on siliceous bedrock.

This is the first karyological data for this species from Macedonia; it confirms the chromosome number for the species previously published from Bulgaria (Vladimirov and Szeląg, 2001) and Serbia (Szeląg et al., 2007).

Pilosella pavichii (Heuff.) Holub $2n = 2x = 18$

1. Macedonia, Jakupica Mts., siliceous rocks along road from Karadžica chalet to Mt. Krivul, 1720 m a.s.l.
2. Montenegro, Prokletije Mts., Čakor pass, 1840 m a.s.l., secondary rocky grassland.

This is the first karyological data for this species from Macedonia and Montenegro. Diploids were reported earlier from Bulgaria, Greece, Romania and Serbia (Christov and Popov, 1933; Strid and Frazén, 1981; Vladimirov, 2000; Mráz and Szeląg, 2004; Szeląg et al., 2007).

Pilosella serbica (F. W. Schultz & Schultz-Bip.) Szeląg $2n = 2x = 18$ (Fig. 4)

1. Macedonia, Korab Mts., Mt. Šila Vort, 2040 m a.s.l., rocky places on granite.
2. Montenegro, Prokletije Mts., Mt. Starac, 2100 m a.s.l., grassland with *Juncus trifidus* on granite.

This is the first chromosome number report for the species from Macedonia and Montenegro; it confirms the previously published count from Serbia (Szeląg et al., 2007; Szeląg, 2008).

REFERENCES

- BUTTLER KP. 1991. *Hieracium* L. In: Strid A, Tan K [eds.], *Mountain Flora of Greece* 2, 595–642. Edinburgh University Press, Edinburgh.
- CHRTEK J, MRÁZ P, ZAHRADNÍČEK J, MATEO G, and SZELĄG Z. 2007. Chromosome numbers and DNA ploidy levels of selected species of *Hieracium* s.str. (Asteraceae). *Folia Geobotanica* 42: 411–430.
- CHRTEK J, ZAHRADNÍČEK J, KRAK K, and FEHRER J. 2009. Genome size in *Hieracium* subgenus *Hieracium* (Asteraceae) is strongly correlated with major phylogenetic groups. *Annals of Botany* 104: 161–178.
- CHRISTOV M, and POPOV A. 1933. Cytologische Studien über die Gattung *Hieracium*. *Planta* 20: 440–447.
- GRAU J, and ERBEN M. 1988. Chromosomenzahlen griechischer Hieracien. *Mitteilungen der Botanischen Staatssammlung München* 27: 97–98.
- HORVAT I, GLAVAČ V, and ELLENBERG H. 1974. *Vegetation Südosteuropas*. Geobotanica selecta 4. Gustav Fischer Verlag, Stuttgart.
- ILNICKI T, and SZELĄG Z. 2011. Chromosome numbers in *Hieracium* and *Pilosella* (Asteraceae) from Central and Southeastern Europe. *Acta Biologica Cracoviensia Series Botanica* 53: 102–110.
- JOACHIMIĄK A, ILNICKI T, and MITKA J. 1999. Karyological studies on *Aconitum lasiocarpum* (Rchb.) Gayer (Ranunculaceae). *Acta Biologica Cracoviensia Series Botanica* 41: 205–211.
- MERXMÜLLER H. 1975. Diploide Hieracien. *Anales del Instituto Botánico A. J. Cavanilles* 32: 189–196.
- MRÁZ P, and SZELĄG Z. 2004. Chromosome numbers and reproductive systems in selected species of *Hieracium* and *Pilosella* (Asteraceae) from Romania. *Annales Botanici Fennici* 41: 405–414.
- NIKETIĆ M, BAREKA P, and KAMARI G. 2003. Karyosystematic study of selected *Hieracium* taxa (Compositae) from Mt Durmitor (Montenegro). *Botanica Chronica* 16: 23–45.
- NIKETIĆ M, VLADIMIROV V, and MRÁZ P. 2006. Chromosome numbers and taxonomic-chorological notes on selected species of *Hieracium* s.str. (Asteraceae) from Montenegro. *Phytologia Balcanica* 12: 85–97.
- SCHUHWERK F, and LIPPERT W. 1998. Chromosomenzahlen von *Hieracium* (Compositae, Lactuceae) Teil 2. *Sendtnera* 5: 269–286.
- STRID A, and FRAZÉN R. 1981. [Report on *Hieracium pavichii*] In: Löve Å [ed.], *Chromosome number reports LXXIII*. *Taxon* 30: 829–861.
- SZELĄG Z. 2006. Taxonomic revision of *Hieracium* sect. *Cernua* (Asteraceae) in the Carpathians, Sudetes and Alps. *Polish Botanical Journal* 51: 97–153.
- SZELĄG Z. 2008. Taxonomic and nomenclatural notes on *Pilosella alpicola* agg. (Asteraceae) in the Balkans and Carpathians. *Annales Botanici Fennici* 45: 301–306.
- SZELĄG Z. 2010. *Hieracia balcanica* V. A new diploid species in *Hieracium* sect. *Naegeliania* (Asteraceae) from Macedonia. *Annales Botanici Fennici* 47: 315–319.
- SZELĄG Z, ILNICKI T, NIKETIĆ M, and TOMOVIĆ G. 2007. Diploid chromosome numbers in five *Hieracium* species from Serbia and Montenegro. *Acta Biologica Cracoviensia Series Botanica* 49: 119–121.
- VLADIMIROV V. 2000. Diploid species of the genus *Hieracium* s.l. in Bulgaria. *Abhandlungen und Berichte des Naturkundemuseums Görlitz* 72 (Suppl.): 16.
- VLADIMIROV V. 2003. A new diploid *Hieracium* (Asteraceae: Lactuceae) from Bulgaria. *Botanical Journal of the Linnean Society* 143: 213–218.
- VLADIMIROV V, and SZELĄG Z. 2001a. Reports (1271–1277). In: Kamari G, Blanché C, Garbari F [eds.] *Mediterranean chromosome number reports 11*. *Flora Mediterranea* 11: 478–483.
- VLADIMIROV V, and SZELĄG Z. 2001b. Chromosome numbers of six species of the genus *Hieracium*, subgenus *Pilosella* (Asteraceae) from Bulgaria. *Polish Botanical Journal* 46: 269–273.
- VLADIMIROV V, and SZELĄG Z. 2006. A new diploid species of *Hieracium* sect. *Pannosa* (Asteraceae) from Bulgaria. *Botanical Journal of the Linnean Society* 150: 261–265.
- ZAHN KH. 1921–1923. *Hieracium* L. In: Engler A [ed.], *Das Pflanzenreich Regni Vegetabilis Conspectus* IV/280, 1–1705. Wilhelm Engelmann, Leipzig.