



Marcin Zarek

Seasonal fluctuations of photosynthetic pigments content in *Taxus baccata* needles

Received: 29 September 2015; Accepted: 5 April 2016

Abstract: Yew is a coniferous, evergreen, dioecious species. The objective of the study was to present a comprehensive characteristic of changes occurring throughout the year in terms of the content of photosynthetic pigments and related compounds in the needles of yew, depending on the sex of individuals and age of needles. Eight compounds, particularly chlorophyll a (Chl *a*), chlorophyll b (Chl *b*), carotenoids (Car), protoporphyrin IX (PPIX), magnesium protoporphyrin IX (MgPPIX), protochlorophyllide (Pchlde), chlorophyllide a (Chlide *a*), and chlorophyllide b (Chlide *b*), were subjected to quantitative analysis. Based on the several parameters under study, significant differences between male and female individuals were observed, while most commonly, the largest differences were reported in the autumn and winter period. They were related to the content of Chl *a*, Chl *b* and Chl/Car ratio. The remaining compounds showed no significant differences according to the sex and were slightly different only in single periods. For all the studied parameters except for Chl/Car ratio, interaction between sampling dates and sex was not statistically significant. Significant differences between the needles of different age were observed only in terms of the content of Chl *b*, MgPPIX, Pchlde, Chlide *b*, and Chl *a/b* ratio, and these differences were always caused by the current-year needles.

Keywords: chlorophyll, chlorophyllide, needles age, pigments, sex

Address: M. Zarek, Department of Forest Pathology, Mycology and Tree Physiology, Institute of Forest Ecosystem Protection, Faculty of Forestry, University of Agriculture in Krakow, Al. 29 Listopada 46, 31-425 Kraków, e-mail: rlzarek@cyf-kr.edu.pl

Introduction

Yew is a coniferous, good shade-tolerant, dioecious, long-living, and relatively late maturing species (Garcia et al., 2000; Thomas & Polwart, 2003). As long as the generative organs of reproduction do not appear, it is not possible to determine the sex of a particular individuals; studies on sex differences in yew thus face many difficulties. A detailed description of the biochemical and physiological differences occurring between male and female individuals constitutes

the first step in the determination of sex markers. Many studies show that dioecious species are more prone to extinction, which is caused by two reasons. Firstly, the necessity of the occurrence of an appropriate number of male and female individuals and secondly the uneven reproductive effort made by both sexes, which directly restricts the vegetative growth in females (De Jong & Van Der Meijden, 2004; Vamosi & Vamosi, 2005; Iszkuło et al., 2009, 2011). As a result of global climate changes, the risk of dioecious species also increases, especially when a given sex ex-

hibits greater habitat requirements (De Jong & Van Der Majden, 2004; Iszkuło et al., 2009). Additionally, in terms of European yew both, excessive acquisition of raw materials and biological characteristics of the species have caused that in many European countries, its range was limited to few and fragmented population. It has led to recognize this species as endangered and to subject it for protection (Peterken, 1996; Seidling, 1999; Svenning & Magård, 1999; Boratyński et al., 2001; Iszkuło & Boratyński, 2005).

Upon entering into generative reproduction stage, needs of male and female organisms change. Both male and female individuals when reaching sexual maturity and entering into the stage of generative reproduction begin to exhibit physiological differences caused by their reproductive effort. Male flowers of *Taxus baccata* begin to develop in the summer during the year preceding pollination. In the autumn or late winter, production of pollen is observed, which is spread by the wind in February and March, followed by pollination. From the pollination to fertilization, 6–12 weeks elapse. Fertilization occurs between May and June, and from that moment, growth and development of the embryo occurs, maturation of arils and seeds is noticed, which ends in September. At the same time, from May to June, continuous development of female “flowers” happen, which will be pollinated the next year (Hejnowicz, 1978) (Fig. 1). All changes in the physiology of male and female organisms may be observed by the content of biochemical compounds (e.g., pigments, proteins, carbohydrates, enzymatic activity) (Nicotra, 1999; Obeso & Retuerto, 2002; Leigh et al., 2006; Montesinos et al., 2006; Iszkuło et al., 2011).

Previous studies determining the sex of yew were relatively few and were conducted in a limited range. It has been observed that in *T. baccata*, male individuals tend to be larger and have higher annual growth in comparison to female individuals (Lloyd & Webb, 1977). Whereas, needles derived from female indi-

viduals are characterized by its higher number and higher density of stomata (Iszkuło et al., 2009). There are also differences in the production of adventitious roots from cuttings collected from male and female *T. baccata* individuals depending on the type of auxin used; however, the mechanism of this reaction is not entirely clear (Nandi et al., 1996). Studies on carbon and nitrogen content have demonstrated the presence of comparable concentrations in both sexes in *T. baccata* (Iszkuło et al., 2009), similarly the taxol content depending on the age and sex of *T. wallichiana* from the Himalayan did not differ significantly between both the sexes (Nadeem et al., 2002). In contrast, Iszkuło et al. (2013) found that female individuals of *T. baccata* exhibited a significantly higher level of taxanes. Female individuals of *T. baccata* are also more demanding in terms of fertility and humidity of the soil (Cedro & Iszkuło, 2011).

The occurrence of biochemical differences occurring in the needles of different life span, related to the processes of development and aging of needles is a different study. In deciduous trees, the process of both synthesis and degradation of pigments is restricted to a single vegetative season. In evergreen trees, the process is extended in time, so most commonly, needles of different age occur on the tree. As for yews, the life span of needles is dependent on the environmental conditions, from 4 to 8 years (Bartkowiak et al., 1978). The progressive aging of the needles eventually leads to an imbalance between biosynthesis and degradation of chlorophyll and may be observed, among others, through changes in the concentrations of pigments and their degradation products (Kutík, 1998; La Porta et al., 2006). The amount of photosynthetic pigments in trees is subjected to changes during growth and development; it also depends, among others, on their age, position around the crown, light conditions, and access to minerals and water (Merzlyak et al., 1999; Silkina & Vinokurova, 2009; Sarala & Saravana, 2011). As the amount of




Fig. 1. Diagram of yew generative and vegetative organs annual development




Fig. 2. The partial pathways of chlorophyll biosynthesis and degradation based on compounds used in this research

light absorbed by the plant is based on the amount of photosynthetic pigments, it can be understood that chlorophyll content has a direct impact on the photosynthetic potential and primary production.

Chlorophyll biosynthesis is a very complex process which involves over 50 biochemical reactions during the synthesis from precursors through, among others, protoporphyrin IX (PPIX), magnesium protoporphyrin IX (MgPPIX), protochlorophyllide (Pchlde), chlorophyllide a (Chlide *a*) and chlorophyllide b (Chlide *b*), and ending with chlorophyll degradation to pheophytin and through pheophorbide to low-molecular weight degradation products (Fig. 2) (Von Wettstein et al., 1995; Yang et al., 1998).

The main objective of the study was to determine differences in the content of photosynthetic pigments and their selected precursors and degradation products between female and male individuals of yew and to determine the course of changes over one year. Additionally, the effect of needles age on the selected parameters of photosynthetic apparatus was investigated.

Materials and methods

Plant material

The study had been conducted for 12 months from March 2013 to February 2014. In the Botanical Garden of the Jagiellonian University in Krakow, 18 European yew individuals (including 12 female and 6 male individuals) grown under similar light conditions were selected. Each of the individual selected was precisely identified in terms of sex by repeated observations of

the occurrence of visible generative reproduction organs: male “flowers” and seeds in arils or the remaining arils in female individuals. Collecting of biological material in the form of shoots with needles was conducted regularly every month, separately for each individual. To minimize the effect of changeable light conditions on the results, shoots were evenly collected from the entire perimeter of the crown of each individual, from its outer part at a height between 1 and 2 m, and then placed in the container on ice. After transportation to the laboratory, shoots were categorized based on its annual increment and classified into one of four groups as: 0 annual – current-year-old needles, 1 annual – one-year-old needles, 2 annual – two-year-olds needles and 3 annual – three-year-olds needles. Then, needles were collected followed by their further homogenization in liquid nitrogen and freeze-drying (Labconco FreeZone 2.5 Liter Freeze Dry System, Kansas City, USA). Material protected in such a way was stored until analysis.

Extraction and quantification

Absorbance analysis was performed using Helios Gamma spectrophotometer (Thermo Spectronic, Cambridge, UK). Quantitative analysis was conducted for eight compounds: Chl *a* and Chl *b* according to PORRA et al. (1989), Car according to Lichtenthaler (1987), Chlide *a* and Chlide *b* according to McFeeters et al. (1971), Pchlde, PPIX, and MgPPIX according to Kahn et al. (1976). The extraction was performed by using 50 mg of freeze-dried powder with 4 ml of 80% buffered acetone. In order to determine Chl *a*, Chl *b*, and Car, the measure of absorbance was performed directly from the buffered acetone after appropriate dilution of the samples. Other compounds were determined after purification of samples with hexane (1:1 v/v). Based on the measurements obtained, the ratio of Chl *a/b* and Chl/Car were calculated.

Statistical analysis

In order to determine the significant differences in the content of the compounds determined, in each of the tested periods, repeated measures of ANOVA (RM-ANOVA) was used. Tukey’s test was used as a post-hoc test for different *N*. In terms of evaluation of the differences between sexes, *N* for each term was established at *N* = 24 for male and *N* = 48 for female individuals. Assessing differences between sexes, samples from different year-old needles were combined in groups for particular individuals. In studies regarding differences between needles age in particular sampling dates, *N* was established at 18 for each year investigated (no division into male and female individuals was made). Homogeneity of variance was tested with Bartlett’s

test (Sokal & Rohlf, 1995). For the data which did not meet the ANOVA assumptions, data transformation or at least ANOVA Kruskal–Wallis non-parametric test was performed. All analyzes were performed using two grouping variables: sex (male and female – without dividing needle's age) and age of needles (current-year-old, one-year-old, two-years-old, and three-years-old – without dividing sex). Annuals of needles were compared from the moment of current year's increment, that is, within the period from May to February. All statistical analyzes were conducted using Statistica software, version

10.0 (StatSoft Inc., Tulsa, OK, USA). Average values are presented as the mean \pm one standard error of the mean (SE). Statistical significance was established as $p \leq 0.05$.

Results

Sex differences

For several investigated parameters, significant differences between male and female individuals




Fig. 3. Seasonal changes in the content of: (A) Chl a, (B) Chl b, (C) Car, (D) Chl a/b ratio and (E) Chl/Car ratio in extracts from the needles of male and female individuals of *Taxus baccata*. The means with the same letters are not significantly different ($P > 0.05$) within a given sampling dates. The months without letters are not significantly different between sexes. Each point is the mean of 24 measurements (\pm SE) for male and 48 measurements (\pm SE) for female. Asterisks (single * for female and double ** for male) indicates a significant difference between means of adjacent sampling dates ($P < 0.05$)

could be observed. They related to the content of Chl *a* (Fig. 3A), Chl *b* (Fig. 3B), and Chl/Car ratio (Fig. 3E). Other parameters: Chl *a/b* ratio (Fig. 3D), Car (Fig. 3C), PPIX (Fig. 4A), MgPPIX (Fig. 4B), Pchlde (Fig. 4C), Chlide *a* (Fig. 4D), and Chlide *b* (Fig. 4E) showed no significant differences between sexes. For all the studied parameters, except for Chl/Car ratio, interaction between sampling dates and sex was not statistically significant. Higher concentrations of Chl *a*, Chl *b*, and Car compounds were observed in female individuals; these differences were statistically significant in the autumn and winter period (Chl *a* from October to December, Chl *b*

from September to December, Car from November to December).

Chl *a* level (Fig. 3A) in female individuals increased slightly from March ($1.95 \text{ mg g}^{-1} \text{ DW}$) to April ($1.99 \text{ mg g}^{-1} \text{ DW}$) and then intensively till June when it reached its maximum value ($3.28 \text{ mg g}^{-1} \text{ DW}$). From that moment onwards, it was slowly decreasing until November ($2.75 \text{ mg g}^{-1} \text{ DW}$) with a sudden decrease ending in January ($2.03 \text{ mg g}^{-1} \text{ DW}$) was observed. In male individuals, a slight decrease in the value of Chl *a* (from $1.99 \text{ mg g}^{-1} \text{ DW}$ in March to $1.82 \text{ mg g}^{-1} \text{ DW}$ in April) was initially observed and the increase in concentrations from April to May was slightly slower




Fig. 4. Seasonal changes in the content of some chlorophyll intermediants and derivatives: (A) PPIX, (B) MgPPIX, (C) Pchlde, (D) Chlide *a*, and (E) Chlide *b*, in extracts from the needles of male and female individuals of *Taxus baccata*. Each point is the mean of 24 measurements ($\pm \text{SE}$) for male and 48 measurements ($\pm \text{SE}$) for female. Asterisks (single * for female and double ** for male) indicates a significant difference between means of adjacent sampling dates ($P < 0.05$)

in comparison to female individuals and still was not statistically significant. From May to June, an intense production of Chl *a* was already observed, although maximum is reached only in July (2.99 mg g⁻¹ DW). High concentrations were established in the period from June to August followed by a fairly rapid decline of this value, ending in December (1.89 mg g⁻¹ DW), therefore earlier than in female individuals.

A similar pattern was observed for Chl *b* (Fig. 3B). The level of this pigment was gradually increasing from March (0.61 mg g⁻¹ DW for females and 0.63 mg g⁻¹ DW for males) until May (0.75 mg g⁻¹ DW for females and 0.70 mg g⁻¹ DW for males), followed by a sudden increase to the level of 1.14 mg g⁻¹ DW for females and 1.02 mg g⁻¹ DW for males, reported in June. The high level was maintained for about 3 months and then from August, steady decline was reported to a level of 0.53 mg g⁻¹ DW for males in December and a little later for females (0.60 mg g⁻¹ DW in January).

Similar course of changes was observed in terms of Car (Fig. 3C) content from March to August, while slightly lower concentrations were observed in male individuals. Initially, the content of Car significantly decreased from March (0.69 mg g⁻¹ DW) to April (0.62 mg g⁻¹ DW for females and 0.57 mg g⁻¹ DW for males), followed by a sudden increase in May (0.64 mg g⁻¹ DW for females and 0.62 mg g⁻¹ DW for males) till June (0.84 mg g⁻¹ DW and 0.79 mg g⁻¹ DW, respectively). Starting from August, in female individuals gradual increase of Car concentration was observed, which ends in November (0.86 mg g⁻¹ DW), followed by an initially fast and subsequent slow decline in the content ending in February (0.71 mg g⁻¹ DW). At the same time, the concentration of Car in male individuals decreases slightly from August to October, while in November it increases to a level of 0.76 mg g⁻¹ DW followed by a rapid decrease to the value of 0.66 mg g⁻¹ DW in December.

The change in the course of Chl *a/b* ratio curve (Fig. 3D) was similar in both sexes and reached its minimum in April (2.89 for females and 2.82 for males), while the maximum value was obtained in December, when it was slightly higher in male individuals (3.56 for males and 3.45 for females). In terms of Chl *a/b* ratio in the annual course, clear fluctuations were observed in the values from March to June and then there was a steady increase from August to December.

The value of Chl/Car ratio (Fig. 3E) in turn increased regardless of the sex in March and April. In the following months, in female individuals, there was a steady increase in the value with the slight slowdown in July and culmination in August established to the level of 5.57. In male individuals, from April to May and then from June to July, growth inhibition was observed followed by the occurrence of maximum similarity as in female individuals in Au-

gust, and it was slightly lower (5.36). From that moment, an intense decline in the value of Chl/Car ratio in both sexes was observed lasting until January.

Early precursors of chlorophyll (PPIX and MgPPIX) reach the highest values in the period between May and June (Figs. 4A,B), then initially their levels decrease rapidly followed by a slow decrease until the initial level. Despite the lack of significant differences between male and female individuals in terms of the content of PPIX and MgPPIX, a shift in the concentration maximum for the analyzed compounds was observed, which for females occurs in May (198.5 nmol g⁻¹ DW and 59.1 nmol g⁻¹ DW, respectively) and for males in June (227.4 nmol g⁻¹ DW and 72.4 nmol g⁻¹ DW, respectively).

Different levels of concentrations are observed for more direct chlorophyll precursors: Pchlide, Chlide *a*, and Chlide *b* (Figs. 4C, D, E). The highest values are reached in March and April, being slightly higher in females in comparison to males. Then, their level decreases rapidly, starting to increase slowly in the autumn and winter period.

Age of needles differences

Significant differences between needles of different ages were observed only in the case of Chl *b* (Fig. 5B), MgPPIX (Fig. 6B), Pchlide (Fig. 6C), Chlide *b* (Fig. 6E) content, and Chl *a/b* ratio (Fig. 5D), and these differences were always caused by the needles classified as current-year needles. Considering only the annuals from the 1st to 3rd year (excluding the current increment), none of the investigated parameters showed statistical significance. Although in needles, a slight increase in the content of Chl *a* (Fig. 5A), Chl *b* (Fig. 5B), Car (Fig. 5C) and Chl/Car ratio (Fig. 5E) with age could be observed, and simultaneous decrease in the concentration of Pchlide (Fig. 6C), Chlide *a* (Fig. 6D), Chlide *b* (Fig. 6E), and Chl *a/b* ratio (Fig. 5D) with age was also reported.

In terms of Chl *a/b* ratio (Fig. 5D), high similarity in the course of curves for 1–3 annuals and different course for 0 annuals was observed, especially in the period from May to June, when the value for 0 annuals increases with simultaneous decrease reported in others. In the subsequent seasons, plots already have a similar course, while at the same time much higher value for 0 annuals is depicted. At the initial stage, current-year needles also showed high levels of concentrations of intermediates (PPIX and MgPPIX in May and Pchlide, Chlide *a*, and Chlide *b* in June) (Figs. 6A,B,C,D,E), which then suddenly decreased, and then in the subsequent part of the year showed much greater instability compared to 1–3 annuals.

Discussion

Differences observed by our group in terms of the occurrence of assimilation pigments (higher concentrations in female and lower in male individuals) are consistent with other studies conducted on dioecious plants. Lower content of assimilation pigments in male individuals was observed, i.e. in *Piper betle* L. (Kumar et al., 2006), *Populus deltoides* (Yang et al., 2011), *Ilex aquifolium* (Obeso et al., 1998), and *Ginkgo biloba* (Mori et al., 2000). Similarly, the leaf photosynthetic rates in *Acer negundo* (Dawson & Ehleringer, 1993) and *Salix arctica* (Dawson & Bliss, 1993) was higher in female individuals. On the con-

trary, in previous studies, Mitchell (1998) observed that male and female individuals from *T. brevifolia* and *T. baccata* growing under similar light conditions did not differ significantly among themselves.

Different results in relation to those obtained by our group may be caused by sampling dates because we observed in our study that the occurrence of differences increases or decreases depending on the month of the year. Differences in the content of assimilation pigments and the rate of photosynthesis also depend on the species. Unlike the European yew, leaves of male individuals in *Salix* and *Populus* contained more chlorophyll in comparison to female individuals (Bourdeau, 1958), while for *Populus trem-*




Fig. 5. Seasonal changes in the content of: (A) Chl a, (B) Chl b, (C) Car, (D) Chl a/b ratio and (E) Chl/Car ratio in extracts from the different age needles of *Taxus baccata*. The means with the same letters are not significantly different ($P > 0.05$) within a given sampling dates. The months without letters are not significantly different between age needles. Each point is the mean of 18 measurements (\pm SE)




Fig. 6. Seasonal changes in the content of some chlorophyll intermediants and derivatives: (A) PPIX, (B) MgPPIX, (C) Pchl a, (D) Chl a, and (E) Chl b in extracts from the different age needles of *Taxus baccata*. The means with the same letters are not significantly different ($P > 0.05$) within a given sampling dates. The months without letters are not significantly different between age needles. Each point is the mean of 18 measurements (\pm SE)

uloides (Bourdeau, 1958) and *Silene latifolia* (Gehring & Monson, 1994), no difference in the rate of photosynthesis was observed.

Differences in the content of assimilation pigments could result from different reproductive effort borne by both sexes. On the one hand, the main effort borne by male individuals is associated with the need to produce as much pollen as possible, and with vegetative growth, which provides increased distribution and greater pollen coverage (Obeso, 2002; Montesinos et al., 2006; Cedro & Iszkuło, 2011). On the other hand, female individuals bear much higher costs related to the production of seeds and fruits during the year, which occurs

from June, i.e., since the end of intensive vegetative growth. The effort borne by female individuals is not only higher, but also more evenly distributed over the year. It begins in the spring with intensive vegetative growth and is further replaced by the effort associated with the production and maturation of fruits and seeds.

To meet this challenge, female individuals not only maintain higher level of assimilation pigments almost throughout the year, but also begin their production (especially Chl a) a little earlier in the spring and retain it longer in autumn. The need for earlier initiation of intensive spring growth in female individuals aimed at compensating a greater effort borne

throughout the year may simultaneously cause an increased risk of late spring frost, over time leading to (as one of the factors) a change in the demographic structure in favor of male individuals in populations, which was, i.e., observed by Iszkuło et al. (2009).

Changes in the content of assimilation pigments observed in our study throughout the year have a similar pattern to that observed by other authors in deciduous trees and among trees such as *Fagus sylvatica* or *Morus alba* (García-Plazaola & Becerril, 2001; Sývacý & Sokmen, 2004) as well as coniferous trees, e.g., *T. cuspidata* (Swanberg & Verhoewen, 2002), *Picea abies*, and *Abies sibirica* (Silkina & Vinokurova, 2009). An increase observed in the amount of assimilation pigments in spring season is the result of chlorophyll biosynthesis prevalence over degradation processes. With the advent of summer, gradual increase in the prevalence of photodegradation processes of chlorophyll is observed, which is no longer compensated by biosynthesis, leading to a gradual decrease in the concentration of chlorophyll.

Because the process of production and decay of chlorophyll is controlled by light, it is important to maintain similar light conditions as possible during the sample collection. With high intensity of light, degradation processes outweigh the synthesis processes so that in plants growing in full sun, less chlorophyll content in comparison to plants growing in the shade was observed (García-Plazaola & Becerril, 2001; Swanberg & Verhoewen, 2002). In the shade, the content of Chl *b*, which allows for the absorption of light in a wider wavelength range, was observed. Therefore, Chl *a/b* ratio is often used as an indicator of plants' response to shading (Goncalves et al., 2001). Similar course of changes in Chl *a/b* ratio to that observed by our group in *T. baccata* was also present in *Piper betle* L. (Kumar et al., 2006) for which it reached the highest value in November (that is a month earlier than *T. baccata* observed by our group), and these values were slightly higher for male in comparison to female individuals. Higher value of Chl *a/b* ratio in male individuals may be caused by a higher degree of conversion of Chl *b* into Chl *a* by specific enzymes (Folly & Engel, 1999; Scheumann et al., 1999) or higher membrane sensitivity (Zu et al., 2010).

Two peaks of Car concentration observed by our group are probably related to the intensive spring production of assimilation pigments (first peak) and to a period of intensive chlorophyll degradation, which leads to an increased photooxidation risks (second peak). Simultaneously, differences in Car content occurring in the vicinity of the second maximum between male and female individuals may be associated with maturation and seed dispersal occurring at that time. The lower content of Car in male individuals was also observed in *Piper betle* L. (Kumar et al., 2006), *Populus deltoides* (Yang et al., 2011),

and *Ginkgo biloba* (Mori et al., 2000). Female individuals exhibit higher levels of Car throughout the year so that they seem to be protected better against photooxidative damages (Gonçalves et al., 2001; Robakowski & Wyka, 2009).

Maximum content of early chlorophyll precursors (PPIX and MgPPIX) delayed by a month in male individuals is associated with later initiation of chlorophyll production and may indicate their delayed spring start, which at the same time constitutes a protection against late frost. Delayed start in the spring and earlier degradation of pigments in autumn may result from natural selection of male individuals under the influence of low temperatures. More sensitivity of male individuals against low temperatures in other species is also highlighted by other researchers (Mori et al., 2000; Kumar et al., 2006; Yang et al., 2011).

Because of assimilation pigments synthesis is highly affected by light conditions it is generally difficult to divide the effect of light from the needles' age. In the case of studies conducted by our group, samples were collected from the outer part of the crown, which along with morphological comb-like structure of the needles' arrangement on the shoots in *T. baccata* should minimize the influence of self-shading effect. Comparing the content of assimilation pigments in needles of different age tested, it can be observed that in terms of Chl *a*, this content does not depend on age throughout the year, while in Chl *b* significantly lower concentrations throughout the year can be found in the youngest needles. In turn, Jach and Ceulemans (2000) observed higher concentration of chlorophyll in this year's needles of *Pinus sylvestris* in comparison to last year's, i.e., in June and September. In the first year, we observed a gradual increase in the content of chlorophyll, while during the second year, a decrease in the concentration of chlorophyll was reported. Differences observed by our group in *T. baccata* in relation to *Pinus sylvestris* may be caused by, i.e., lifetime of the needles in particular species, which in terms of European yew are estimated at 2–6 and even eight years as reported by other authors (Szaniawski, 1978). According to Szaniawski (1978), the ability of European yew needles to perform photosynthesis decreases with age and drops to 50% in seven-year-old needles compared to young needles.

Differences observed by our group in terms of Car content between different year-old needles exhibit a slight increase in concentrations, which probably results from their gradual accumulation along with needles' age. Increased instability in three-year-old needles throughout the year may indicate early symptoms of aging. The value of Chl/Car ratio also slightly increases with needles' age, which is caused by a faster increase in the content of chlorophyll and a slightly lower increase in Car content. Increase in Car content along with age of needles was also observed

in, i.e., Scots pine (Miazek & Ledakowicz, 2013). The value of Chl/Car ratio was lower in younger tissues of Scots pine's and also increased with age. For young, susceptible tissues, higher content of Car in relation to chlorophyll can serve as a protective function against oxidative stress or excessive radiation. During vegetative season, chlorophyll is decomposed earlier than Car, which leads to a decrease in the Chl/Car ratio. In studies conducted by our group, we did not observe any decrease in this value with needles' age, which may indicate that three-year-old needles have not initialized the process of aging on a large scale.

High value of Chl *a/b* ratio observed in young *T. baccata* needles reported by our group is mainly caused by a relatively low content of Chl *b*. In turn, decreasing Chl *a/b* ratio in progressively older *T. baccata* needles is similar to changes observed in *Cupressus sempervirens* aging needles (La Porta et al., 2006). In case of *Cupressus sempervirens*, authors explained the changes by decreasing amount of Chl *a* resulting from aging processes of needles. In our case, decrease in Chl *a/b* ratio was mainly caused by the increased content of Chl *b* in relation to Chl *a*. Decreased values of this coefficient observed by our group in one-, two-, and three-year-old *T. baccata* needles in relation to this year's needles may indicate better acclimatization of older needles' year to changing light conditions related to, i.e., increased self-shading from year to year.

Moreover, in terms of Scots pine (Miazek & Ledakowicz, 2013), individuals exposed to intense light were characterized by higher Chl *a/b* ratio in comparison to those growing under less intense light. According to Zu et al. (2010), increase in Chl *a/b* ratio may be caused by the increasing degree of decomposition of grana containing photosystem PSII.

The youngest needles during the most intense growth phase exhibit very high concentration of early chlorophyll precursors (PPIX and MgPPIX), which is most likely related to the need of rapid production of the appropriate level of pigments ensuring an adequate level of assimilation. More direct precursors such as Pchl*ide*, Chl*ide a*, and Chl*ide b* reach their maxima in the youngest needles a month later in relation to early precursors, which is caused by the increased production of assimilation pigments. In subsequent three years, the level of all precursors in needles remains very balanced with no signs of decline with increased age. Our studies do not indicate the existence of significant differences in the content of assimilation pigments and their precursors associated with the process of needles' aging during the first four years, but only small differences related to their development in the first year of life.

Conclusions

Differences in the content of assimilation pigments can constitute a kind of biochemical marker of sex under the maintenance of specific sampling conditions. The most suitable period of plant material harvesting ranges from November to December, when the greatest differences in the concentrations of both Chl *a*, Chl *b* and Car as well are observed.

Analyzing the impact of needles' age on the concentrations of compounds tested, it can be concluded that the youngest needles (current-year-old needles) are most different from others (both in terms of concentration range and dynamics of seasonal changes). In turn, among three-year-old needles, seasonal fluctuations slowly start to increase, which may constitute first symptoms of aging. When collecting biological material for research purposes, it is best to choose one-year-old or two-year-old needles, which, on the one hand, are already fully developed and stable and, on the other hand, do not demonstrate any symptoms of aging.

In addition, one should also take into account a great impact of lighting conditions on the concentrations of the compounds described. To obtain meaningful results, the influence of light intensity should be excluded as much as possible, which is often difficult to achieve in field conditions; however, it may allow for the identification of sex in case of artificial cultivation of propagation material carried out under controlled conditions. The studies conducted also indicate a quite high variability of the traits studied throughout the year, depending on the sex of individuals investigated and age of needles that indicates the necessity of a strict consideration of these parameters when comparing the results obtained. The difference observed between male and female individuals need to be confirmed by further analysis of biochemical compounds, and in the future, it may become an element of biochemical markers' system that identifies sex.

Acknowledgements

This Research was financed by the Ministry of Science and Higher Education of the Republic of Poland. The author gratefully wish to thank the Director of the UJ Botanical Garden in Krakow for providing yew collection.

References

- Bartkowiak S, Bugała W, Czartoryski A, Hejnowicz A, Król S, Środoń A & Szaniawski RK (1978) The Yew – *Taxus baccata*. Foreign Scientific Publications, Department of the National Center for Scientific and

- Technical, and Economic Information (for the Department of Agriculture and the National Science Foundation, Washington, DC), Warsaw.
- Boratyński A, Didukh Y & Lucak M (2001) Yew (*Taxus baccata* L.) population in Knyazhdvir Nature Reserve in the Carpathians (Ukraine). *Dendrobiology* 46: 3–8.
- Bourdeau PF (1958) Photosynthetic and respiratory rates in leaves of male and female Quaking Aspens. *Forest Science* 4: 331–334.
- Cedro A & Iszkuło G (2011) Do Females Differ from Males of European Yew (*Taxus baccata* L.) in Dendrochronological Analysis? *Tree-Ring Research* 67: 3–11.
- Dawson TE & Bliss LC (1993) Plants as Mosaics: Leaf-, Ramet-, and Gender-Level Variation in the Physiology of the Dwarf Willow, *Salix arctica*. *Functional Ecology* 7: 293–304.
- Dawson TE & Ehleringer JR (1993) Gender-specific physiology, carbon isotope discrimination, and habitat distribution in boxelder, *Acer negundo*. *Ecology* 74: 798–815.
- Folly P & Engel N (1999) Chlorophyll b to chlorophyll a conversion precedes chlorophyll degradation in *Hordeum vulgare* L. *Journal of Biological Chemistry* 274: 21811–21816.
- García D, Zamora R, Hódar JA, Gomez JM & Castro J (2000) Yew (*Taxus baccata* L.) regeneration is facilitated by fleshy-fruited shrubs in Mediterranean environments. *Biological Conservation* 95: 31–38.
- García-Plazaola J & Becerril JM (2001) Seasonal changes in photosynthetic pigments and antioxidants in beech (*Fagus sylvatica*) in a Mediterranean climate: implications for tree decline diagnosis. *Australian Journal of Plant Physiology* 28: 225–232.
- Gehring JL & Monson RK (1994) Sexual Differences in Gas-Exchange and Response to Environmental-Stress in Dioecious *Silene latifolia* (*Careophyllaceae*). *American Journal of Botany* 81: 166–174.
- Gonçalves JFDC, Marengo RA & Vieira G (2001) Concentration of photosynthetic pigments and chlorophyll fluorescence of mahogany and tonka bean under two light environments. *Revista Brasileira de Fisiologia Vegetal* 13: 149–157.
- Hejnowicz A (1978) Anatomy, embryology and karyology: The Yew – *Taxus baccata* L. (ed. by S Bartkowiak, W Bugala, A Czartoryski, A Hejnowicz, S Król, A Środon & RK Szaniawski) Foreign Scientific Publications, Department of the National Center for Scientific and Technical, and Economic Information (for the Department of Agriculture and the National Science Foundation, Washington, DC), Warsaw, Poland, pp. 33–54.
- Iszkuło G & Boratyński A (2005) Different age and spatial structure of two spontaneous subpopulations of *Taxus baccata* as a result of various intensity of colonization process. *Flora* 200: 195–206.
- Iszkuło G, Jasińska AK, Giertych MJ & Boratyński A (2009) Do secondary sexual dimorphism and female intolerance to drought influence the sex ratio and extinction risk of *Taxus baccata*? *Plant Ecology* 200: 229–240.
- Iszkuło G, Jasińska AK & Sobierajska K (2011) Dendroecological differences between *Taxus baccata* males and females in comparison with monoecious *Abies alba*. *Dendrobiology* 65: 55–61.
- Iszkuło G, Kosiński P & Hajnos M (2013) Sex influences the taxanes content in *Taxus baccata*. *Acta Physiologiae Plantarum* 35: 147–152.
- Jach ME & Ceulemans R (2000) Effects of season, needle age and elevated atmospheric CO₂ on photosynthesis in Scots pine (*Pinus sylvestris*). *Tree Physiology* 20: 145–157.
- De Jong TJ & Van Der Meijden E (2004) Sex ratio of some long-lived dioecious plants in a sand dune area. *Plant Biology* 6: 616–620.
- Kahn VM, Avivi-Bieise N & von Wettstein D (1976) Genetic regulation of chlorophyll synthesis analyzed with double mutants in barley: Genetics biogenesis of chloroplasts and mitochondria (ed. by T Bhuchler) Elsevier/North-Holland Biomedical Press, Amsterdam, pp. 119–131.
- Kumar N, Gupta S & Tripathi AN (2006) Gender-specific responses of *Piper betle* L. to low temperature stress: Changes in chlorophyllase activity. *Biologia Plantarum* 50: 705–708.
- Kutik J (1998) The development of chloroplast structure during leaf ontogeny. *Photosynthetica* 35: 481–505.
- Leigh A, Cosgrove MJ & Nicotra AB (2006) Reproductive allocation in a gender dimorphic shrub: Anomalous female investment in *Gynatrix pulchella*? *Journal of Ecology* 94: 1261–1271.
- Lichtenthaler HK (1987) Chlorophylls and carotenoids: Pigments of photosynthetic biomembranes. *Methods in Enzymology* 148: 350–382.
- Lloyd DG & Webb CJ (1977) Secondary sex characters in plants. *The botanical review* 43: 177–216.
- McFeeters RE, Chichester CO & Whitaker JR (1971) Purification and Properties of Chlorophyllase from *Ailanthus altissima* (Tree-of-Heaven). *Plant Physiology* 47: 609–618.
- Merzlyak MN, Gitelson AA, Chivkunova OB & Rakitin VYU (1999) Non-destructive optical detection of pigment changes during leaf senescence and fruit ripening. *Physiologia Plantarum* 106: 135–141.
- Miazek K & Ledakowicz S (2013) Chlorophyll extraction from leaves, needles and microalgae: A kinetic approach. *International Journal of Agricultural and Biological Engineering* 6: 107–115.
- Mitchell AK (1998) Acclimation of Pacific yew (*Taxus brevifolia*) foliage to sun and shade. *Tree Physiology* 18: 749–757.

- Montesinos D, De Luís M, Verdú M, Raventós J & García-Fayos P (2006) When, how and how much: Gender-specific resource-use strategies in the dioecious tree *Juniperus thurifera*. *Annals of Botany* 98: 885–889.
- Mori M, Suzuki K & Kohzaki R (2000) Variations in chlorophyll and carotenoid content in the growth process of the *Ginkgo* leaf. *Journal of the Japanese Society for Food Science and Technology-Nippon Shokuhin Kagaku Kogaku Kaishi* 47: 448–451.
- Nadeem M, Rikhari HC, Kumar A, Palni LMS & Nandi SK (2002) Taxol content in the bark of Himalayan Yew in relation to tree age and sex. *Phytochemistry* 60: 627–631.
- Nandi SK, Palni LMS & Rikkari HC (1996) Chemical induction of adventitious root formation in *Taxus baccata* cuttings. *Plant Growth Regulation* 19: 117–122.
- Nicotra AB (1999) Reproductive allocation and the long-term costs of reproduction in *Siparuna grandiflora*, a dioecious neotropical shrub. *Journal of Ecology* 87: 138–149.
- Obeso JR (2002) The costs of reproduction in plants. *New Phytologist* 155: 321–348.
- Obeso J, Alvarez-Santullano M & Retuerto R (1998) Sex ratios, size distributions, and sexual dimorphism in the dioecious tree *Ilex aquifolium* (Aquifoliaceae). *American Journal of Botany* 85: 1602–1608.
- Obeso JR & Retuerto R (2002) Dimorfismo sexual en el acebo, *Ilex aquifolium*: ¿Coste de la reproducción, selección sexual o diferenciación fisiológica? *Revista Chilena de Historia Natural* 75: 67–77.
- Peterken GF (1996) *Natural woodland: ecology and conservation in northern temperate regions*. Cambridge University Press.
- Porra RJ, Thompson WA & Kriedemann PE (1989) Determination of accurate extinction coefficients and simultaneous equations for assaying chlorophylls a and b extracted with four different solvents: verification of the concentration of chlorophyll standards by atomic absorption spectroscopy. *BBA – Bioenergetics* 975: 384–394.
- La Porta N, Bertamini M, Nedunchezian N & Muthuchelian K (2006) Photosynthetic changes that occur during aging of cypress (*Cupressus sempervirens* L.) needles. *Photosynthetica* 44: 555–560.
- Robakowski P & Wyka T (2009) Winter photoinhibition in needles of *Taxus baccata* seedlings acclimated to different light levels. *Photosynthetica* 47: 527–535.
- Sarala TD & Saravana KR (2011) The monthly changes of chloroplast pigments content in selected plant species exposed to cement dust pollution. *Journal of Research in Biology* 8: 660–666.
- Scheumann V, Schoch S & Rüdiger W (1999) Chlorophyll b reduction during senescence of barley seedlings. *Planta* 209: 364–370.
- Seidling W (1999) Raumlische struktur einer subspontanen population von *Taxus baccata*-Jungpflanzen. *Flora* 194: 439–451.
- Silkina OV & Vinokurova RI (2009) Seasonal dynamics of chlorophyll and microelement content in developing conifer needles of *Abies sibirica* and *Picea abies*. *Russian Journal of Plant Physiology* 56: 780–786.
- Sokal RR & Rohlf FJ (1995) *Biometry: The Principles and Practices of Statistics in Biological Research*. W.H. Freeman and Co.
- Swanberg A & Verhoeven A (2002) Seasonal acclimation of leaf antioxidant systems in the evergreen *Taxus cuspidata* growing in sun and shade. *The Journal of the Minnesota Academy of Science* 66: 30.
- Svenning JC & Magård E (1999) Population ecology and conservation status of the last natural population of English yew *Taxus baccata* in Denmark. *Biological Conservation* 88: 173–182.
- Sývácý A & Sökmen M (2004) Seasonal changes in antioxidant activity, total phenolic and anthocyanin constituent of the stems of two *Morus* species (*Morus alba* L. and *Morus nigra* L.). *Plant Growth Regulation* 44: 251–256.
- Szaniawski RK (1978) *An Outline of Yew Physiology: The Yew – Taxus baccata* L. (ed. by S Bartkowiak, W Bugała, A Czartoryski, A Hejnowicz, S Król, A Środoń & RK Szaniawski) Foreign Scientific Publications, Department of the National Center for Scientific and Technical, and Economic Information (for the Department of Agriculture and the National Science Foundation, Washington, DC), Warsaw, Poland, pp. 55–64.
- Thomas PA & Polwart A (2003) *Taxus baccata* L. *Journal of Ecology* 91: 489–524.
- Vamosi JC & Vamosi SM (2005) Present day risk of extinction may exacerbate the lower species richness of dioecious clades. *Diversity and Distributions* 11: 25–32.
- Von Wettstein D, Gough S & Kannangara CG (1995) Chlorophyll Biosynthesis. *The Plant Cell* 7: 1039–1057.
- Yang C-M, Chang K-W, Yin M-H & Huang H-M (1998) Methods for the Determination of the Chlorophylls and their Derivates. *Taiwania* 43: 116–122.
- Yang F, Wang Y, Wang J, Deng W, Liao L & Li M (2011) Different eco-physiological responses between male and female *Populus deltoides* clones to waterlogging stress. *Forest Ecology and Management* 262: 1963–1971.
- Zu Y, Pang H-H, Yu J-H, Li D-W, Wei X-X, Gao Y-X & Tong L (2010) Responses in the morphology, physiology and biochemistry of *Taxus chinensis* var. *mairei* grown under supplementary UV-B radiation. *Journal of Photochemistry and Photobiology. B, Biology* 98: 152–158.