



# World News of Natural Sciences

An International Scientific Journal

WNOFNS 24 (2019) 64-70

EISSN 2543-5426

---

---

## Antioxidant Enzyme Mechanism of Cluster Bean (*Cyamopsis tetragonaloba* (L.) Taub). Under Cobalt Stress

**Kaliyamoorthy Jayakumar**

Department of Botany, A.V.C. College (Autonomous), Mannampandal 609 305, Tamil Nadu, India

E-mail address: [jk81botany@yahoo.com](mailto:jk81botany@yahoo.com)

### ABSTRACT

The research work has been carried out to estimate the low level cobalt application enhance antioxidant enzyme activities of cluster bean plants. The experiments were conducted in earthen pots containing 3 kg of air dried soil. The inner surface of pots was lined with polythene sheet. Cluster bean plants were raised in soil amended with different concentration of cobalt (0, 50, 100, 150, 200 and 250 mg/kg). The antioxidant enzyme activities were analysed at 30, 60, and 90 days after sowing (DAS). There found a significant enhance antioxidant enzyme activities at 50 mg/kg cobalt application in the soil in all the sampling days when compared to control. Further increase in the cobalt level (100-250 mg/kg) in the soil decreased antioxidant enzyme activities of cluster bean plants in all the sampling days.

**Keywords:** Cobalt, *Cyamopsis tetragonaloba*, Catalase, Peroxidasa, Polyphenol oxidase

### 1. INTRODUCTION

The presence of heavy metals in the environment is of major concern because of their toxicity and threat to plant and animal life. Heavy metal availability in soil which depends on natural processes, particularly lithogenic and pedogenic, as well as anthropogenic factors such as industrial activity and mining, sewage disposal, traffic, etc., are mainly responsible for the increase in the concentration of heavy metals in air, water and soil. The presence of heavy metals in residues from city and sewage sludge as well as some inorganic fertilizers applied to agricultural lands is beginning to be the cause as a new source of pollution in soils and plants.

The heavy metal concentrations are increasing as environmental pollutants particularly in densely industrialized and heavy road traffic areas. The heavy metals can create a major ecological crisis since they are non-degradable and often accumulate through trophic level causing a deleterious biological effect. Some of the heavy metals to be considered as essential trace elements for the normal growth of plants.

The biological importance of cobalt was first recognized by the discovery that small amounts of the elements would certain deficiency symptoms in plants. Even though cobalt has been recognized as a micronutrient for animals and certain microorganisms, until recently there has been no conclusive evidence of its essentiality for higher plants. The cobalt as a heavy metal pollutant in plants has been studied previously.

In abiotic stress, metal response will result in the production of reactive oxygen species (ROS) which leads to the activation of defense mechanisms in terms of antioxidant enzymes. Generation of ROS such as superoxide, H<sub>2</sub>O<sub>2</sub> and hydroxyl molecules cause rapid cell damage by triggering off a chain reaction. Plants under stress produce some defence mechanisms to protect themselves from the harmful effect of oxidative stress. ROS scavenging is one among the common defense response against abiotic stresses. ROS scavenging depends on the detoxification mechanism provided by an integrated system of non-enzymatic reduced molecules like ascorbate and glutathione and enzymatic antioxidants. The major ROS scavenging activities include complex non-enzymatic (ascorbate, glutathione,  $\alpha$ -tocopherol) and enzymatic antioxidants like catalase (CAT), ascorbate peroxidase (APX), glutathione reductase (GR), superoxide dismutase (SOD), polyphenol oxidase (PPO), peroxidase (POX) etc. The pathways include the water-water cycle in chloroplasts and the ascorbate-glutathione cycle. Antioxidant mechanisms may provide a strategy to enhance metal tolerance in plants.

Cluster bean (*Cyamopsis tetragonaloba* (L.) Taub.) is one of the world's main sources of edible vegetable oils and high-protein livestock feed. There are many reports on the effect of heavy metals up on the morphological and biochemical characteristics of Cluster bean.

The effect of cobalt on nodulation, leg-haemoglobin content and antioxidant enzyme activities of crop plants attracted little attention. So, the present investigation was carried out to evaluate the effect of this heavy metal, cobalt on the root nodule formation, leg-haemoglobin content antioxidant enzyme activities of *Cyamopsis tetragonaloba*.

## **2. MATERIALS AND METHODS**

The seeds of Cluster bean (*Cyamopsis tetragonaloba*.) were obtained from Tamilnadu Agricultural University, Tamilnadu. Plants were grown in untreated soil (control) and in soil to which cobalt had been applied (50, 100, 150, 200 and 250 mg/kg soil). The cobalt as finely powdered cobalt chloride (CoCl<sub>2</sub>) was applied to the surface soil and thoroughly mixed with soil. Ten seeds were sown in each pot. All the pots were watered to field capacity daily. Plants were thinned to a maximum of five per pot, after a week of germination. The total number of root nodules and leg-haemoglobin content were analysed at 15 days intervals, namely 15, 30, 45, 60 and 75 days after sowing (DAS). The leg-haemoglobin contents of root nodules were analysed by following method. Catalase (CAT) (EC 1.11.1.6) activity was measured according to the method of Chandlee and Scandalios 1963 with small modification. 0.5 g of frozen plant material was homogenized in a prechilled pestle and mortar with 5 ml of ice cold 50 mM sodium phosphate buffer (pH 7.5) containing 1 mM phenylmethylsulfonyl fluoride (PMSF).


The extract was centrifuged at 4 °C for 20 min at 12,500 rpm. The supernatant was used for enzyme assay. The assay mixture contained 2.6 mL of 50 mM potassium phosphate buffer (pH 7.0), 400  $\mu$ L of 15 mM H<sub>2</sub>O<sub>2</sub> and 40  $\mu$ L of enzyme extract. The decomposition of H<sub>2</sub>O<sub>2</sub> was followed by the decline in absorbance at 240 nm.

Peroxidase (POX; EC 1.11.1.7) was assayed by the method of Kumar and Khan 1982. Assay mixture of POX contained 2 mL of 0.1 M phosphate buffer (pH 6.8), 1 mL of 0.01 M pyrogallol, 1 mL of 0.005 M H<sub>2</sub>O<sub>2</sub> and 0.5 mL of enzyme extract. The solution was incubated for 5 min at 25 °C after which the reaction was terminated by adding 1 mL of 2.5 N H<sub>2</sub>SO<sub>4</sub>. The amount of purpurogallin formed was determined by measuring the absorbance at 420 nm against a blank prepared by adding the extract after the addition of 2.5 N H<sub>2</sub>SO<sub>4</sub> at zero time. The activity was expressed in unit mg<sup>-1</sup> protein. One unit (U) is defined as the change in the absorbance by 0.1 min<sup>-1</sup> mg<sup>-1</sup> protein.

Polyphenol oxidase (PPO; EC 1.10.3.1) activity was assayed by the method of Kumar and Khan 1982. Assay mixture for PPO contained 2 mL of 0.1 M phosphate buffer (pH 6.0), 1 mL of 0.1 M catechol and 0.5 mL of enzyme extract. This was incubated for 5 min at 25 °C, after which the reaction was stopped by adding 1 mL of 2.5 N H<sub>2</sub>SO<sub>4</sub>. The absorbancy of the purpurogallin formed was read at 495 nm. To the blank 2.5 N H<sub>2</sub>SO<sub>4</sub> was added of the zero time of the same assay mixture. PPO activity is expressed in U mg<sup>-1</sup> protein (U = Change in 0.1 absorbance min<sup>-1</sup> mg<sup>-1</sup> protein). The enzyme protein was estimated by the method of Bradford 1976, for expressing all the enzyme activities.

Statistical analysis was performed using one way analysis of variance (ANOVA) followed by Duncan's Multiple Range Test (DMRT). The values are mean  $\pm$  SD for six samples in each group. *P* values  $\leq$  0.05 were considered as significant.

### 3. RESULTS



**Figure 1.** The effect of different concentrations of cobalt on catalase, peroxidase and polyphenol oxidase activity (units mg<sup>-1</sup> protein) of Cluster bean.

#### 4. DISCUSSION

The maximum number of nodule formation was occurred in 50 mg/kg on 75 DAS and minimum number of nodule formation was observed at 250 mg/kg on 15 DAS. The number of nodule formation of Cluster bean decreased (except 50 mg/kg) with an increase in the cobalt concentration of soil. Similar reduction in the number of nodule formation was observed under different heavy metal treatments in Cluster bean. The leg-haemoglobin contents of nodule of Cluster bean plants increased at 50 mg/kg soil level and decreased further with an increase in cobalt level in the soil. Minimum leg-haemoglobin content was recorded at 250 mg/kg cobalt level. This can be compared with earlier report. CAT activity decreased with increasing concentration of Co (100-250 mg kg<sup>-1</sup>) than the control and low level of Co (50 mg kg<sup>-1</sup>) treated Cluster bean plants. POX and PPO activities increased (except 50 mg kg<sup>-1</sup>) with an increase in Co level in the soil. To be able to endure oxidative damage under conditions which favours increased oxidative stress such as high/low temperatures, water deficit, salinity etc., plants must possess efficient antioxidant system. Plants possess antioxidant systems in the form of enzymes such as SOD, APX, CAT and metabolites viz., ascorbic acid, glutathione,  $\alpha$ -tocopherol, carotenoid, flavonoids, etc. These antioxidant enzymes and metabolites are reported to increase under various environmental stresses as well as comparatively higher activity has been reported in fungicide, triadimefon in medicinal plants, suggesting that higher antioxidant enzymes activity have a role in imparting tolerance against any type of environmental stresses. From the results of this investigation it can be concluded that, the application of cobalt at 50 mg/kg may be proved to be a useful tool in increasing the root nodule formation and leg haemoglobin contents in Cluster bean plants. This study is of high significance, because it is previously reported that using isotopic nitrogen, roots and nodules grown in the presence of added cobalt possess a greater capacity for nitrogen fixation than those without added cobalt.

#### 5. CONCLUSION

Cobalt treatment at all levels tested (except 50 mg/ kg) decreased the nodules, leg-haemoglobin contents of plants; antioxidant enzyme (CAT) activity of cluster bean plants. However the antioxidant enzymes (POX and PPO) increased with an increase in cobalt level in the soil. From the present investigation it can be concluded that the 50 mg /kg level of cobalt in the soil is beneficial for the growth of cluster bean plants.

#### References

- [1] Aery, N.C. and S. Sarkar, (1991). Studies on the effect of heavy metal stress on growth parameters of soybean. *J. Environ. Biol.* 12(1): 15-24
- [2] Bradford, M.M., (1976). A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein dye binding. *Anal. Biochem.* 72: 248-253
- [3] Chandlee, J.M., Scandalios, J.G., (1984). Analysis of variants affecting the catalase development program in maize scutellum. *Theor. Appl. Gen.* 69: 71-77

- [4] Chen, L.M., Linard, C.C., Kao, C.H., (2001). Copper toxicity in rice seedlings: changes in antioxidative enzyme activities, H<sub>2</sub>O<sub>2</sub> level and cell wall peroxidase activity in roots. *Bot. Bull. Acad. Sinica* 41: 99-103
- [5] Delwiche, C.C., C.M. Johnson, H.N. Reisenauer. (1960). Influence of Cobalt on nitrogen by *Medicago*. *Plant Physiol.* 73-78
- [6] El-Sheekh, M.M., A.H. El-Naggar, M.E.H. Osman and E. El-Mazaly, (2003). Effect of cobalt on growth, pigment and the photosynthetic electron transport in *Monoraphidium minutum* and *Nitzhia perminuta*. *Braz. J. Plant Physiol.* 15(3): 159-166
- [7] Imlay, J.A., (2003). Pathways of oxidative damage. *Annu. Rev. Microbiol.* 57: 395-418
- [8] Jaleel C.A., B. Sankar, P.V. Murali, M. Gomathinayagam, G.M.A. Lakshmanan, R. Panneerselvam. Water deficit stress effects on reactive oxygen metabolism in *Catharanthus roseus*; impacts on ajmalicine accumulation, *Colloids Surf. B: Biointerfaces*, 29 Sep 2007, 62(1): 105-111. doi:10.1016/j.colsurfb.2007.09.026
- [9] Jaleel C.A., G.M.A. Lakshmanan, M. Gomathinayagam, R. Panneerselvam, (2008). Triadimefon induced salt stress tolerance in *Withania somnifera* and its relationship to antioxidant defense system. *South African J. Bot.* 74:126-132
- [10] Jaleel C.A., P. Manivannan, A. Kishorekumar, B. Sankar, R. Gopi, R. Somasundaram, R. Panneerselvam, (2007). Alterations in osmoregulation, antioxidant enzymes and indole alkaloid levels in *Catharanthus roseus* exposed to water deficit, *Colloids Surf. B: Biointerfaces* 59: 150-157
- [11] Jaleel C.A., P. Manivannan, A. Kishorekumar, B. Sankar, R. Panneerselvam, (2007). Calcium chloride effects on salinity induced oxidative stress, proline metabolism and indole alkaloid accumulation in *Catharanthus roseus*. *C.R. Biologies* 330: 674-683
- [12] Jaleel C.A., P. Manivannan, B. Sankar, A. Kishorekumar, R. Gopi, R. Somasundaram, R. Panneerselvam, (2007). Induction of drought stress tolerance by ketoconazole in *Catharanthus roseus* is mediated by enhanced antioxidant potentials and secondary metabolite accumulation, *Colloids Surf. B: Biointerfaces* 60: 201-206.
- [13] Jaleel C.A., P. Manivannan, M. Gomathinayagam, R. Sridharan, R. Panneerselvam, (2007). Responses of antioxidant potentials in *Dioscorea rotundata* Poir. following paclobutrazol drenching, *C. R. Biologies* 330: 798-805
- [14] Jaleel C.A., R. Gopi, A. Kishorekumar, P. Manivannan, B. Sankar, R. Panneerselvam, (2007). Interactive effects of triadimefon and salt stress on antioxidative status and ajmalicine accumulation in *Catharanthus roseus*, *Acta Physiol. Plant.* DOI 10.1007/s11738-007-0119-1
- [15] Jaleel C.A., R. Gopi, B. Sankar, P. Manivannan, A. Kishorekumar, R. Sridharan, R. Panneerselvam, (2007). Studies on germination, seedling vigour, lipid peroxidation and proline metabolism in *Catharanthus roseus* seedlings under salt stress, *South Afr. J. Bot.* 73: 190-195
- [16] Jaleel C.A., R. Gopi, P. Manivannan, M. Gomathinayagam, P.V. Murali, R. Panneerselvam, (2007). Soil applied propiconazole alleviates the impact of salinity on

- Catharanthus roseus* by improving antioxidant status, *Pestic. Biochem. Physiol.* doi:10.1016/j.pestbp.2007.11.003
- [17] Jaleel C.A., R. Gopi, P. Manivannan, R. (2007). Panneerselvam, Antioxidative potentials as a protective mechanism in *Catharanthus roseus* (L.) G. Don. Plants under salinity stress, *Turkish J. Botany* 31: 245-251.
- [18] Jaleel C.A., R. Gopi, P. Manivannan, R. Panneerselvam, (2007). Responses of antioxidant defense system of *Catharanthus roseus* (L.) G. Don. to paclobutrazol treatment under salinity, *Acta Physiol. Plant.* 29: 205-209
- [19] Jaleel C.A., R. Gopi, P. Manivannan, R. Sridharan, M. Gomathinayagam, R. Panneerselvam, (2007). Antioxidant potential and indole alkaloid profile variations with water deficits along different parts of two varieties of *Catharanthus roseus*, *Colloids Surf. B: Biointerfaces*, doi:10.1016/j.colsurfb.2007.10.013
- [20] Jaleel C.A., R. Gopi, G.M. Alagu Lakshmanan, R. Panneerselvam, (2006). Triadimefon induced changes in the antioxidant metabolism and ajmalicine production in *Catharanthus roseus* (L.) G. Don., *Plant Sci.* 171, 271-276
- [21] Jaleel CA, Manivannan P, Lakshmanan GMA, Sridharan R, Panneerselvam R (2007). NaCl as a physiological modulator of proline metabolism and antioxidant potential in *Phyllanthus amarus*. *Comptes Rendus Biologies* 330: 806-813
- [22] Jayakumar K., C.A. Jaleel, P. Vijayarengan, (2007). Changes in growth, biochemical constituents and antioxidant potentials in radish (*Raphanus sativus* L.) under cobalt stress, *Turk. J. Biol.* 31: 112-117
- [23] Jayakumar, K. and P. Vijayarengan, (2006). Influence of cobalt on seed germination and seedling growth of *Vigna mungo* (L.) Hepper. *Plant. Arch.* 6 (2): 681-682
- [24] Jayakumar, K., P. Vijayarengan, M. Vijayaragavan, S. Rajasekaran, K. Sumathi, P. Sundaramoorthy, (2006). Toxicological aspects of cobalt induced growth and morphological changes in *Helianthus annuus* L. *Plant. Arch.* 6 (2): 829-830
- [25] Jayakumar, K., Vijayarengan, P., (2006). Alterations in the carbohydrate metabolism of *Vigna mungo* (L.) Hepper as affected by cobalt stress. *Indian J. Environ. Ecoplan.* 12: 693-696
- [26] Kumar, K.B., Khan, P.A., (1982). Peroxidase and polyphenol oxidase in excised ragi (*Eleusine coracana* cv. PR 202) leaves during senescence. *Indian J. Exp. Bot.* 20: 412-416
- [27] Moral R, Gomez J, Pedreno N and Mataix J. (1994). Effect of cadmium on nutrient distribution, yield and growth of tomato grown in soil less culture. *J. Plant Nutr.* 17: 953-962
- [28] Saravanan, S., A. Subramani, P. Sundaramoorthy, M. Selvaraju and A.S. Lakshmanachary, (2001). Influence of copper sulphate on germination, growth and yield of Soybean (*Glycine max* (L.) Merr.). *Ecol. Env. and Cons.* 7(2): 141-144
- [29] Savoure, A., Thorin, D., Davery, M., Xue, J.H., Mauro, S., Van Montagu, M., Inze, D., Verbruggen, N., (1999). NaCl and CuSO<sub>4</sub> treatments trigger distinct oxidative defense mechanisms in *Nicotiana plumbaginifolia*. *Plant Cell Environ.* 22: 387-396

- [30] Seliga, H., (1993). The role of copper in nitrogen fixation in *Lupinus luteus* L. *Plant Soil* 155: 349-352
- [31] Seliga. H., (1990). *Acta Physiol. Plant.* 12: 287-291
- [32] Vesper, S.J. and T.C. Weidensaul, (1978). Effects of cadmium, nickel, copper and Zinc on nitrogen fixation by Horse gram s. *Water, Air, Soil Pollut.* 9: 413-422
- [33] Wilson D.D. and Reisenaver, (1963). Determination of leg-haemoglobin in legume nodules. *Analytical Biochemistry*, 6: 21-30