

E. L. GREGORASZCZUK AND R. PIEKŁO

## THYROID HORMONE ACTION IN PORCINE LUTEAL CELLS. EFFECT OF TRIIODOTHYRONINE ON MITOCHONDRIAL CYTOCHROME P450-SCC ACTIVITY

Laboratory of Animal Endocrinology, Department of Animal Physiology, Institute of Zoology,  
Jagiellonian University, Krakow, Poland

In the present study hydroxylated cholesterol derivatives were used to monitor P450<sub>sc</sub> activity as an effect of triiodothyronine. Luteal cells isolated from mid-developing and mature corpora lutea were plated into 24-well plates by 18 h incubation with M199 supplemented with 5% of alf serum. After that time the plates were washed with fresh M199 and hydroxylated cholesterol derivatives (25-, 20-, and 22-hydroxylated cholesterol) were added to the control (not-T3-treated) and T3-treated cells. Two hours later, all cultures were terminated and the media were frozen until further progesterone analysis. Triiodothyronine added to the culture medium of cells isolated from mature corpora lutea increased both basal and hydroxylated cholesterol derivative-stimulated cells. Aminoglutethimide, a P450<sub>sc</sub> enzyme inhibitor, added to the culture medium in all doses used, had no effect on basal progesterone secretion while added to the T3-treated cells reduced progesterone production. This study strongly supports the hypothesis of a direct effect of thyroid hormone on mitochondrial cytochrome P450<sub>sc</sub> — catalysing side-chain cleavage of cholesterol enzyme in luteal cells.

*Key words: thyroid hormone, corpus luteum, cytochrome P450<sub>sc</sub>, progesterone production.*

### INTRODUCTION

The effects of thyroid hormones on mitochondria and energy metabolism have been well documented (1—4). It is, however, commonly assumed that hormonal effects on mitochondria are secondary to a primary action on nuclear gene expression, generating regulatory signals channelled to the mitochondria.

A possible basis for direct thyroid hormone action on mitochondria was provided by findings of a thyroid hormone-binding component in mitochondrial membranes (5—7). The saturable receptor was found in the

inner mitochondrial membranes of rat liver, kidney, myocardium and skeletal muscle (8). Previous research from our laboratory (9) showed the direct effect of T<sub>3</sub> on progesterone secretion by luteal cells. The effect of thyroid hormone on cyclic nucleotide release by ovarian cells was previously shown, suggesting the involvement of cAMP-dependent mechanism in the realization of T<sub>3</sub> action in ovarian cells (10). Surprisingly, there are no data concerning the presence of mitochondrial T<sub>3</sub> receptors in the ovarian tissue. The activity of the mitochondrial cholesterol side chain cleavage cytochrome (P450<sub>scc</sub>) has long been proposed as the rate-limiting step in the production of progesterone. It has been found that one cytochrome P-450 enzyme, is localized on the inner mitochondrial membrane (11). The study of Sugano *et al.* (12) provide direct evidence that cholesterol is metabolized to pregnenolone by way of 22R-hydroxycholesterol and 20R, 22R-dihydrocholesterol by P450<sub>scc</sub>.

The present experiment was therefore conducted to determine the effect of T<sub>3</sub> on the activity of P450<sub>scc</sub> — by measuring the conversion of hydroxylated cholesterol derivatives to progesterone.

## MATERIAL AND METHODS

### *Reagents*

Triiodothyronine ( $10^{-9}$  M), 25-, 20-, 22-R-hydroxycholesterol (10 µg/ml) and aminoglutethimide (1, 10 or 20 µg/ml) were purchased from Sigma Chemical Co, St. Louis, MO, USA.

Medium M199, penicillin, trypsin, and calf serum were from the Laboratory of Vaccines, Lublin, Poland.

### *Animals*

Porcine ovaries were collected in ice-cold physiological saline from a local abattoir immediately after slaughter and transported to the laboratory. The phase of the oestrous cycle was determined according to the established morphological criteria (13, 14). Ovarian tissues representing three different stages of the oestrus cycle were collected from: newly developed corpora lutea (CL1; 1—3 days after ovulation), mature corpora lutea (CL2; 7—10 days after ovulation), and regressing corpora lutea (CL3; 12—14 days after ovulation).

### *Cell culture*

Dissected corpora lutea of each type were enzymatically dissociated according to the technique of Gregoraszczyk (15). The cells were suspended in medium M199 supplemented with 5% of calf serum at a concentration of  $3.5 \times 10^5$  cells/ml medium.

Cells were grown in multiwell plates (Nunck) in a humidified atmosphere with 5% CO<sub>2</sub> in the air for 24 h. After 24 h incubation all cultures were terminated and the media were frozen until further steroid analysis. Cell viability using the trypan blue exclusion test was 85%.

## *Experimental procedure*

Experiment 1. To show the dose response to T3 with progesterone secretion dispersed cells were incubated without (control) and with three doses ( $10^{-7}$  M,  $10^{-9}$  M, and  $10^{-11}$  M) of T3. After 24 h incubation, all cultures were terminated and the media were frozen until further progesterone analysis.

Experiment 2. This experiment required the use of hydroxylated cholesterol derivatives (25-, 20-, and 22-hydroxylated cholesterol) therefore a preliminary study evaluated different doses of those cholesterol derivatives on progesterone production. It was found that 10  $\mu$ g/ml was a maximally stimulated dose for all cholesterol derivatives. This is in accordance with Wiltbank *et al.* (16). Cells were plated into 24-well plates by 18 h incubation with M199 supplemented with 5% of calf serum. After that time hydroxylated cholesterol derivatives (25-, 20-, and 22-hydroxylated cholesterol) was added to the control (not-T3-treated) and T3-treated cells. Six hours later, all cultures were terminated and the media were frozen until further progesterone analysis.

Experiment 3. Taking into consideration results obtained from Experiment 2, 20-hydroxylated cholesterol was used in this experiment. Luteal cells were cultured for 18 h with M199 supplemented with 5% of calf serum and 20-hydroxylated cholesterol. After that time aminoglutethimide, the inhibitor of P450<sub>scc</sub> was added to the control (not-T3-treated) and T3-treated cells. Six hours later, all cultures were terminated and the media were frozen for further progesterone analysis.

## *Progesterone analysis*


Progesterone was detected in the culture medium by a radioimmunoassay described elsewhere (13). A highly specific antibody raised in sheep against 11 $\alpha$ -hydroxy-progesterone hemisuccinate coupled to bovine serum albumin was used. The cross-reaction with pregnenolone was 2.9%. All other tested steroids showed less than 0.1% cross-reaction. (1,2,6,7-<sup>3</sup>H) progesterone (Radiochemical Centre, Amersham, England, sp. act. 80 Ci/mmol) was used as the tracer. The limit of sensitivity of the assay was 50 pg/ml. The coefficients of variation within and between assays were 15% and 2.5%, respectively.

All data points are expressed as means  $\pm$  SEM derived from at least three different replications, each in triplicate, resulting in at least nine observations. Significant differences between steroid levels in control and treated cells were compared by analysis of variance and by using Duncan's new multiple range test.

## RESULTS

### *The effect of exogenous triiodothyronine on progesterone secretion*

All the applied doses of T3 stimulated progesterone secretion by cells isolated from early developing corpora lutea (CL1), in a doses  $10^{-11}$  M and  $10^{-9}$  M increased progesterone production by cells isolated from mature corpora lutea (CL2) and had no effect on progesterone production by cells isolated from regressing corpora lutea (CL3) (*Fig. 1*).



*Fig. 1.* Effect of triiodothyronine ( $10^{-7}$ — $10^{-9}$  — and  $10^{-11}$  MT) on progesterone secretion measured following culture for 24 h of porcine luteal cells collected from: newly developed corpora lutea (CL1; 1—3 days after ovulation), mature corpora lutea (CL3; 7—10 days after ovulation) and regressing corpora lutea (CL4; 12—14 days after ovulation). Each bar represents the mean  $\pm$  SE of 3 replicate cultures each in triplicate. \* $p < 0.05$ ; \*\* $p < 0.01$ ; \*\*\* $p < 0.001$ .





Fig. 2. Progesterone production by luteal cells isolated from mature (CL3) corpora lutea. Cells were plated into 24-well plates for 18 h incubation with M199 supplemented with 5% of calf serum. The plates were then washed with fresh M199 and 25-hydroxylated cholesterol (25-OH); 20-hydroxylated cholesterol (20-OH), and 22-hydroxylated cholesterol (22-OH) was added to the control (not-T3-treated) and T3-treated cells. Two hours later, all cultures were terminated and the media were frozen until further progesterone analysis. Each bar represents the mean  $\pm$  SE of replicate cultures each in triplicate. \* $p < 0.05$ ; \*\* $p < 0.01$ ; \*\*\* $p < 0.001$ .



*Fig. 3.* Progesterone production by luteal cells mature (CL2) corpora lutea. Cells were cultured for 18h with M199 supplemented with 5% of calf serum. The plates were then washed with fresh medium and aminoglutethimide, the inhibitor P450<sub>scc</sub> was added to the control (not-T3-treated cells). Two hours later, all cultures were terminated and the media were frozen for further progesterone analysis. Each bar represents the mean  $\pm$  SE of 3 replicate cultures each in triplicate.

\* $p < 0.05$ ; \*\* $p < 0.01$ ; \*\*\* $p < 0.001$ .

*The effect of T3 on 25-, 20-, and 22-hydroxylated cholesterol-stimulated progesterone secretion by cells isolated mature corpora lutea.*

Triiodothyronine added to the culture medium of cells isolated from mature corpora lutea increased both basal ( $1546.6 \pm 46.1$  ng vs  $733.3 \pm 76.3$  ng/ $10^5$  cells in control culture),  $p < 0.01$ , and hydroxylated cholesterol derivative-stimulated cells ( $1220.6 \pm 39.0$  ng vs  $1071.6 \pm 59.2$  ng/ $10^5$  cells in 25-OH-stimulated cells;  $1195 \pm 14.2$  ng vs  $1022.6 \pm 10.3$  ng/ $10^5$  cells in 22-OH-stimulated cells and  $1210 \pm 53.7$  ng vs  $953.3$  ng/ $10^5$  cells in 20-OH-stimulated cells),  $p < 0.05$ . (Fig. 2).

*The effect of aminoglutethimide on basal and T-3 stimulated progesterone secretion*

Aminoglutethimide added to the control cultures in all the doses used, had no effect on progesterone secretion ( $845.5 \pm 60.3$  ng;  $764.9 \pm 20.0$  ng and  $756.3 \pm 23.1$  ng/ $10^5$  cells as the effect of 1, 10, and 20  $\mu$ g AG respectively vs  $857.0 \pm 52.0$  ng/ $10^5$  cells in the control culture) (Fig. 3).

Aminoglutethimide added to the T-3 treated cells in a dose dependent manner reduced T3-stimulated progesterone production ( $1412.5 \pm 36.1$  ng,  $1203.0 \pm 10.0$  ng and  $1054.0 \pm 17.2$  ng/ $10^5$  cells as the effect of 1, 10, and 20  $\mu$ g AG respectively as  $11485.3 \pm 34.1$  ng/ $10^3$  cells in the T3-stimulated cells). (Fig. 3).

## DISCUSSION

The effects of thyroid hormones on mitochondria and energy metabolism have been the subject of intensive research and are well documented (3, 8, 17—19). Demonacos *et al.*, (20) proposed that thyroid hormone effects on mitochondria gene(s) are direct, concomitant with the effects on nuclear genes and involving similar molecular mechanisms to those mediating steroid-thyroid hormone actions on nuclear gene transcription. In luteal cells the activity of P450<sub>sc</sub> has long been proposed as the rate-limiting step in the production of progesterone. It has been found that one catochrome P450 enzyme, localized on the inner mitochondrial membrane, is responsible for both the 20- and 22-hydroxylation of cholesterol and for the final oxidation step which causes cleavage of the side chain (21, 22). In the present study hydroxylated cholesterol derivatives were used to monitor P450<sub>sc</sub> activity as an effect of triiodothyronine. Triiodothyronine added to the culture medium increased both basal and hydroxylated cholesterol derivative-stimulated cells.

This study suggested that all the 20 and 22 hydroxylation steps and the side chain cleavage step of the  $450_{\text{scc}}$  are acutely regulated by triiodothyronine. The cholesterol content, phospholipid composition, and degree of fatty acid saturation all change in the mitochondrial inner membrane with the thyroid state (23). The possibility that T3 acted as an inhibitor of  $P450_{\text{scc}}$  was examined by using aminoglutethimide, the main inhibitor of  $P450$  cytochrome (11). Aminoglutethimide, added to the culture medium in all doses used, had no effect on basal progesterone secretion while added to the T3-treated cells reduced progesterone production. In conclusion, this study suggested that cytochrome  $P450_{\text{scc}}$  — catalysing side-chain cleavage of cholesterol may be a key target of T3 action on the mitochondria in luteal cells. Additional studies of the mitochondrial receptors will help clarify the role of thyroid hormones in the regulation of mitochondrial activity in the luteal cells.

*Acknowledgements:* The authors wish to thank to Professor S. Stoklosowa for her constant interest, A. Walocha MSc for performing RIA and Ms B. Snakowska for technical assistance. The work was supported by a grant from the Polish State Research Committee 5 PO6D 030 14.

#### REFERENCES

1. Gustafsson R, Tata JR, Lindberg O, Ernster L. The relationship between the structure and activity of rat skeletal muscle mitochondria after thyroidectomy and thyroid hormone treatment. *J Cell Biol* 1965; 26: 555—578.
2. Sterling K. Direct triiodothyronine (T3) action by a primary mitochondrial pathway *Endocr Res* 1989; 15: 683—715.
3. Kadenbach B. Effect of thyroid hormone on mitochondrial enzymes. In : Regulation of Metabolic Processes in Mitochondria. Targar JM, Papa S, Ouagliarello E, Slater EC (eds.) Amsterdam, Elsevier, 1966; pp. 508—517.
4. Mowbray J, Hardy DL. Direct thyroid hormone signalling via ADP-ribosylation controls mitochondrial nucleotide transport and membrane leakiness by changing the conformation of the adenine nucleotide transporter. *FEBS Letters* 1965; 394: 61—65.
5. Sterling K, Milch PO. Thyroid hormone binding by a component of mitochondrial membrane. *Proc Natl Acad Sci USA* 1975; 72: 3225—3229.
6. Gregory RB, Berry MN. The administration of triiodothyronine to rats resulting in a lowering of the mitochondrial membrane potential in isolated hepatocytes. *Biochem Biophys Acta* 1991; 1133: 89—94.
7. Hafner RP, Brown GC, Brand MD. Thyroid-hormone control of state 3 respiration in isolated rat liver mitochondria. *Biochem J* 1990; 265: 731—734.
8. Sterling K, Lazarus JH, Milch PO. Mitochondrial thyroid hormone receptors: localization and physiological significance. *Science* 1978; 201: 1126—1129.
9. Gregoraszczyk EL. The effect of triiodothyronine (T3) and TSH treatment in vitro on progesterone production by luteal cells isolated during different stages of the luteal phase. *J Physiol Pharmacol* 1996; 47: Suppl 1: 93—100.
10. Gregoraszczyk EL, Galas J. Triiodothyronine-dependent activation of cAMP accumulation and steroid secretion by follicular and luteal cells. *Endocr Regul* 1998; 32: 93—98.



11. Robel P. Steroidogenesis: The enzymes and regulation of their genomic expression. In: *Reproduction in Mammals and Man*. Thibault C, Levasseur MC, Hunter RHF (eds.), Paris, Elipses, 1993; pp. 136—145.
12. Sugano S, Miura R, Morishima N. Identification of intermediates in the conversion of cholesterol to pregnenolone with a reconstituted cytochrome P-450<sub>sc</sub> system: accumulation of the intermediate modulated by the adrenodoxin level. *J Biochem (Tokyo)* 1996; 120: 780—78.
13. Gregoraszczuk EL. Interrelationship between steroid hormone secretion and morphological changes of porcine corpora lutea at various periods of luteal phase. *Endocr Regul* 1992; 26: 189—194.
14. Schilling E. Stages of ovarian function in the sow. In: *Veterinary Medical Review*, Elwert EG (ed). Marburg/Lahn, University und Verlagsbuchlandlung, 1974; pp. 59—63.
15. Gregoraszczuk EL.: Steroid hormone release in cultures of pig corpus luteum and granulosa cells. Effect of LH, hCG, PRL and estradiol, *Endocrinol Exp* 1983; 17: 59—63.
16. Wiltbank MC, Belfiore CJ, Niswender GD. Steroidogenic enzyme activity after acute activation of protein kinase (PK) A and PKC in ovine small and large luteal cells. *Moll Cell Endocrinol* 1993; 97: 1—7.
17. Sterling K. Direct thyroid hormone activation of mitochondria: The role of adenine nucleotide translocase. *Endocrinology* 1986; 119: 292—295.
18. Sterling K, Brenner MA. Thyroid hormone action: Effect of triiodothyronine on mitochondrial adenine nucleotide translocase in vivo and in vitro. *Metabolism* 1995; 44: 193—199.
19. Wrutniak C, Cassar-Malek I, Marchal S *et al.* A 43-kDa protein related to c-Erb A  $\alpha 1$  is located in the mitochondrial matrix of rat liver. *J Biol Chem* 1995; 270: 16347—16354.
20. Demonacos CV, Karayanni N, Hatzoglou E, Tsiriyiotis CH, Spandidos DA, Sekeris CE. Mitochondrial genes as sites of primary action of steroid hormones. *Steroids* 1996; 61: 226—232.
21. Vickery LE. Cholesterol side chain cleavage cytochrome P450 (P450<sub>sc</sub>) In: *Handbook of Experimental Pharmacology*. Schenkman JB and Greim H (eds.) Berlin, Springer-Verlag. 1993; 105, pp. 651—665.
22. Miller WL. Molecular biology of steroid hormone synthesis. *Endocr Rev* 1988; 9: 295—318.
23. Horrum MA, Tobin RB, Ecklund RE. Effects of the novel thyroid hormone analogues. SKF L-94901, DIBIT, and 3'-AC-T<sub>2</sub> on mitochondrial function. *Biochem Mol Biol Int* 1996; 38: 61—72.

Received: April 9, 1998

Accepted: July 9, 1998

Author's address: Prof. Ewa Gregoraszczuk, Institute of Zoology, Jagiellonian University, Ingardena 6, 30-060 Cracow, Poland.