

THE CONTAMINATION OF THE ENVIRONMENT WITH *TOXOCARA* EGGS IN MAZOWIECKIE VOIVODSHIP AS A RISK OF TOXOCAROSIS IN CHILDREN

JAKUB GAWOR AND ANNA BORECKA

Laboratory of Domestic Animals, W. Stefański Institute of Parasitology of the Polish Academy of Sciences, Twarda 51/55, 00-818 Warszawa; E-mail: gaworj@twarda.pan.pl; ab@twarda.pan.pl

ABSTRACT. The home surroundings (109 places) of 120 children with diagnosed toxocarosis were examined to assess if the presence of *Toxocara* spp. eggs in the soil corresponds with the occurrence of clinical cases. 31 places (28.4%) were found to be contaminated. The highest level of soil contamination was revealed in rural and suburban regions (15.6% and 11.9%, respectively), while in urban areas it was 0.9%. These results confirmed that soil contamination might be recognized as a very important factor in the epidemiology of human toxocarosis.

Key words: children, contamination, soil, *Toxocara* spp, toxocarosis.

INTRODUCTION

Human toxocarosis is caused by an infection with larvae of canine or feline roundworm *Toxocara* spp. *T. canis* and *T. cati* are common parasites of young dogs and cats. Infection in Poland was recorded in 18.2-36.3% puppies in towns and 58.0-72.7% young dogs in shelters (Kozakiewicz 1983, Okulewicz 1996, Mizgajska and Luty 1998, Kornaś et al. 2001). *Toxocara cati* was noted in 40% cats in Łódź (Głuszkowska 1955), 68% in Wrocław (Ramisz and Martynowicz 1963), 10.5-62% in Warszawa (Żebrowska 1961, Borecka 1999) and 50% in Lublin and Puławy (Gundlach et al. 1991).

Humans are paratenic hosts of dog and cat roundworms. Men become infected through ingestion of infective eggs containing infective larvae. The major risk of infection is for children, because of their close contacts with puppies, poor hygiene, specific behaviour (geophagy) and lower level of resistance when compared with adults (Glickman et al. 1979, Marczyńska 1999, Ołdakowska et al. 1999).

In humans larvae of *Toxocara* spp. migrate to the liver or other internal organs, but can also cause serious ocular damage by migrating into the retina. According to localisation of larvae, toxocarosis can appear as the visceral form (*visceral larva migrans*), the ocular form (*ocular larva migrans*) or covert toxocarosis. Clinical

manifestation of toxocarosis depends on the number of larvae ingested and frequency of infection, distribution of larvae in the body and intensity of the host immunological response (Baever et al. 1952, Juszko et al. 1994, Ołdakowska et al. 1999).

The contamination of the environment with *Toxocara* eggs has been studied in towns in Poland, in Poznań – 6.3-10.0% soil samples positive (Mizgajska 1997, Maśnik 2000), in Wrocław – 6.0% (Mizgajska 1999), in Lublin and Puławy – 22.0-71.4% (Gundlach et al. 1996), in Warszawa – 5.6% (Borecka 2001), in Kraków – 17.0% (Mizgajska et al. 2001) and in Elbląg – 23.6% (Jarosz 2001).

The aim of the study was to assess if the presence of *Toxocara* spp. eggs in the soil corresponds with the occurrence of clinical cases in children.

MATERIAL AND METHODS

In 2002-2003 the home surroundings of children with diagnosed toxocarosis were examined. The entire group of patients comprise 120 cases with visceral, ocular and covert toxocarosis (52.5%, 25.8% and 21.7%, respectively) from rural areas (60%), suburban regions (30%) and towns (10%) in central of Poland (Table 1). Sand and soil samples were collected in total of 109 places, in backyards, gardens, sandboxes and playgrounds. In each place ten samples, each one about 100 grams were taken. *Toxocara* eggs were extracted using the Dada method.

Table 1. Clinical cases of toxocarosis in children in Mazowieckie voivodship

Form of toxocarosis	Number of cases	rural	Area suburban	urban
Visceral	63	38	19	6
Ocular	31	20	9	2
Covert	26	14	8	4

The dried soil samples were sieved through a mesh. 10 gram samples were transferred to volumetric flasks containing 50 ml of 0.1% Tween 80, and vortexed for 30 min. The suspensions were transferred to 60 ml centrifuge tubes, then centrifuged at 1500 rpm for 10 min, after which the supernatant free of ova was decanted. A saturated flotation solution – ZnSO₄ (specific gravity 1.52) was added and the tubes were vortexed again, then centrifuged at 1500 rpm for 10 minutes, after which the solution was added to form a meniscus and a coverslip was overlayed. The cover-slip was transferred to a glass slide after 15 min, and examined under x100 magnification and the number of *Toxocara* spp. ova were counted.

RESULTS AND DISCUSSION

The highest level of soil contamination was revealed in rural and suburban regions (15.6% and 11.9%, respectively), while in urban areas it was 0.9% (Table 2).

The results showed that in Poland, similarly as in other European countries there is a potential risk of toxocarosis for humans. This especially concerns rural areas, where the high number of toxocarosis cases had been noted (Table 1).

Comparable results of the soil contamination was recorded by Mizgajska and Luty (1998) and Jarosz et al. (2003) in villages near Poznań – Murowana Goślina (8% samples positive), Kołaczkowo (14.7%) and Lusowo (8%), near Kraków – Grodkowice and Łązkowice (16%) (Mizgajska 2000). A higher level of soil contamination with *Toxocara* eggs was found by Gundlach et al. (1996) in villages near Lublin and Puławy (35.5%).

Table 2. Contamination of soil with *Toxocara* eggs in Mazowieckie voivodship

Places examined/contaminated (%)			Total
rural	suburban	urban	
64/17	15.6%	37/13	11.9%
8/1	0.9%	109/31	28.4%

The present results revealed much smaller percentage of soil contamination with *Toxocara* eggs in urban areas (0.9%), when compare to village environments (15.6%) in Mazowieckie voivodship. The results of other authors in Poland have shown high level of soil contamination in towns – in Poznań 6.3% – 12.0%, in Wrocław 18.6%, in Elbląg 16.6%, in Kraków 16.0% and in Bytom 20.3% (Mizgajska and Luty 1998, Maśnik 2000, Mizgajska 2000, Mizgajska et al. 2001, Petryszak 2001, Petryszak et al. 2002, Petryszak and Nosal 2003).

The present study revealed that in the country children infect with toxocarosis in their home environment through contaminated soil. However, the direct contact with dogs and cats cannot be excluded as a source of infection. The examinations of Wolfe and Wright (2003) in the USA have shown that dogs hair can be a source of *Toxocara canis* infective eggs. Their results suggested that dogs infected with *T. canis* may infect people through direct contact.

ACKNOWLEDGMENTS

Work supported by funding from the State Committee for Scientific Research project No 3PO5D 074 23 – „Dependence of clinical cases of toxocarosis in children upon the prevalence of *Toxocara* spp. in dogs and cats and contamination of environment with *Toxocara* eggs”.

REFERENCES

- Beaver P.C., Snyder C.H., Carrera G.M., Dent J.H., Lafferty J.W. 1952. Chronic eosinophilia due to visceral larva migrans. *Pediatrics* 9: 7-19.
- Borecka A. 1999. Ekstensywność zarażenia kotów *Toxocara cati*. *Nowa Weterynaria* 13: 2-3.
- Borecka A. 2001. Poziom zarażenia psów i stopień zanieczyszczenia piaskownic jajami geohelmintów na terenie Warszawy i okolic. *Wiadomości Parazyologiczne* 47 (Supl. 2): 7.
- Glickman T., Schantz P.M., Cypress R.M. 1979. Canine and human toxocariasis: review of transmission, pathogenesis, and clinical disease. *Journal of the American Veterinary Medical Association* 175: 1265-1269.
- Głuszkowska A. 1955. Robaki pasożytnicze przewodu pokarmowego kotów w Łodzi. *Wiadomości Parazyologiczne* 11: 261-266.
- Gundlach J., Sadzikowski A., Tomczuk K., 1996. Zanieczyszczenie jajami *Toxocara* spp. wybranych środowisk miejskich i wiejskich. *Medycyna Weterynaryjna* 52: 395-396.
- Gundlach J., Tomczuk K., Sadzikowski A. 1991. Pasożyty wewnętrzne psów i kotów w aspekcie zagrożenia zdrowia człowieka. Materiały z XVI Zjazdu PTP, Poznań, 12-13.09.1991: 63.
- Jarosz W. 2001. Zanieczyszczenie gleby jajami *Toxocara* spp. na terenie Elbląga. *Wiadomości Parazyologiczne* 47: 143-149.
- Jarosz W., Kacprzak E., Rychlicki W., Mizgajska-Wiktor H. 2003. Soil contamination with *Toxocara* spp. eggs and toxocarosis in people from rural areas of Poznań region (Kołaczkowo, Lusowo). *Wiadomości Parazyologiczne* 49: 399.
- Juszko J., Marczyńska M., Żarnowska H. 1994. Diagnostyka i leczenie postaci ocznej infekcji larwą glisty psiej *Toxocara canis*. *Klinika Oczna* 96: 275-280.
- Kornaś S., Nowosad B., Skalska M. 2001. Występowanie glisty *Toxocara canis* u psów w krakowskim schronisku dla bezdomnych zwierząt. *Wiadomości Parazyologiczne* 47: 755-762.
- Kozakiewicz B. 1983. Ekstensywność inwazji *Toxocara canis* u psów i jej aspekty epidemiologiczne w aglomeracji miejskiej. *Medycyna Weterynaryjna* 39: 660-664.
- Marczyńska M. 1999. Przebieg kliniczny i leczenie toksokarozy u dzieci. *Polski Merkuriusz Lekarski* 6: 377-379.
- Maśnik E. 2000. Zależności między występowaniem jaj *Toxocara* w kale psów i w glebie. *Wiadomości Parazyologiczne* 46: 239-244.
- Mizgajska, H. 1997. The role of some environmental factors in the contamination of soil with *Toxocara* spp. and other geohelminth eggs. *Parasitology International* 46: 67-72.
- Mizgajska H. 1999. Biologiczne skażenie gleby na terenach popowodziowych we Wrocławiu. *Wiadomości Parazyologiczne* 45: 89-93.
- Mizgajska H. 2000. Zanieczyszczenie gleby jajami *Toxocara* spp. na terenie Krakowa i pobliskich wsi. *Wiadomości Parazyologiczne* 46: 105-110.
- Mizgajska H., Luty T. 1998. Toksokaroza u psów i zanieczyszczenie gleby jajami *Toxocara* spp. w aglomeracji poznańskiej. *Przegląd Epidemiologiczny* 54: 441-446.
- Mizgajska H., Jarosz W., Remejciak A. 2001. Rozmieszczenie źródeł inwazji *Toxocara* spp. w środowisku miejskim i wiejskim w Polsce. *Wiadomości Parazyologiczne* 47: 399-404.
- Okulewicz A. 1996. Toxocarosis problem in large city (Wrocław). *Parassitologia* 38: 374.
- Ołdakowska A., Marczyńska M., Popielska J. 1999. Hiperleukocytoza i eozynofilia w przebiegu toksokarozy – opis przypadków klinicznych. *Pediatria Polska* 5: 515-19.
- Petryszak A. 2001. Zanieczyszczenie jajami *Toxocara* spp. gleby zieleńców miejskich w Krakowie (dzielnica Krowodrza). *Zeszyty Naukowe Akademii Rolniczej w Krakowie* 386: 129-135.
- Petryszak A., Nosal P. 2003. Zanieczyszczenie jajami glist *Toxocara* spp. gleby zieleńców miejskich Bytomia. *Roczniki Naukowe Zootechniki* (Supl. 17): 779-782.
- Petryszak A., Nosal P., Nowosad B. 2002. Zanieczyszczenie jajami *Toxocara* spp. gleby zieleńców miejskich Krakowa. Materiały Konferencyjne Konferencji naukowej pt. „Zoonozy: problem nadal

- aktualny", Warszawa 6.12.2002: 65.
- Ramisz A., Martynowicz T. 1963. Parazytofauna przewodu pokarmowego psów i kotów m. Wrocławia i okolic, ze szczególnym uwzględnieniem nicieni z rodziny *Ancylostomidae*. *Wiadomości Parazytologiczne* 9: 125-127.
- Wolfe A., Wright I.P. 2003. Human toxocariasis and direct contact with dogs. *Veterinary Record* 152: 419-422.
- Żebrowska D. 1961. Helmintofauna kotów (*Felis catus domesticus*) Warszawy i jej okolic. *Wiadomości Parazytologiczne* 7 (Supl.): 279-282.

Zaakceptowano do druku 14 czerwca 2004