



Metlaouia Noubhani and Cappetta, 1997 (Chondrichthyes: Orectolobiformes) preoccupied by *Metlaouia* Dumont, 1928 (Insecta: Lepidoptera)

ABDELMAJID NOUBHANI and HENRI CAPPETTA

Several years ago, we published a series of new genera and species of elasmobranchs from the Maastrichtian–Paleogene of Morocco (Noubhani and Cappetta 1997). Among these is *Metlaouia* Noubhani and Cappetta, 1997 (Orectolobiformes, Orectolobidae), based on the type species *Squatirhina numidica* Arambourg, 1952.

Recently, Dr. John K. Page informed us of the existence of the homonym *Metlaouia* Dumont, 1928 Lepidoptera, Noctuidae). The lepidopteran genus is available and has priority, so that *Metlaouia* Noubhani and Cappetta, 1997 must be regarded as a junior homonym. In accordance with the International Code of Zoological Nomenclature (ICZN 1999, articles 52, 60), we herein propose a new replacement name, *Eometlaouia*, for the fossil elasmobranch in question.

Genus *Eometlaouia* nov.

(= *Metlaouia* Noubhani and Cappetta, 1997)

Type species: *Squatirhina numidica* Arambourg, 1952.

Etymology: Metlaoui, southern Tunisia, the type locality; in combination with the prefix eos (Greek, dawn), in reference to the Eocene epoch.

Distribution.—Metlaoui, southern Tunisia; and Ouled Abdoun Basin, Morocco; Ypresian (early Eocene).

Acknowledgements.—We thank John K. Page, Production Section Chief of the Zoological Record, Great Britain, who brought this problem of homonymy to our attention. This is Contribution de l’Institut des Sciences de l’Évolution, Université de Montpellier II, number 2002-004.

References

- Arambourg, C. 1952. Les vertébrés fossiles des gisements de phosphates (Maroc–Algérie–Tunisie). *Notes et Mémoires du Service géologique du Maroc* 92 : 1–372.
Dumont, C. 1928. Notes sur les Lépidoptères de Barbarie. In: P. Lechevalier (ed.), *Encyclopédie entomologique*, sér. B, 3, 17–37, 89–93. Paris.
International Commission on Zoological Nomenclature (cited as ICZN). 1999. *International Code of Zoological Nomenclature*, 4th edition, xxix + 306 pp. The International Trust for Zoological Nomenclature, c/o The Natural History Museum, London.
Noubhani, A. and Cappetta, H. 1997. Les Orectolobiformes, Carcharhiniformes et Myliobatiformes (Elasmobranchii, Neoselachii) des bassins à phosphate du Maroc (Maastrichtien–Lutétien basal). Systématique, biostratigraphie, évolution et dynamique des faunes. *Palaeo Ichthyology* 8: 1–327.

Abdelmajid Noubhani [noubhani@ucd.ac.ma], Université Chaouïda Doukkali, Faculté des Sciences, B.P. 20, El Jadida, Morocco;
Henri Cappetta [cappetta@isem.univ-montp2.fr], Laboratoire de Paléontologie, UMR 5554 “Institut des Sciences de l’Evolution”, Université de Montpellier II – Sciences et Techniques du Languedoc, CC 064, Place Eugène Bataillon, 34095 Montpellier Cedex 5, France.