



Choonsig Kim, Jaeyeob Jeong\*

## Comparision of soil CO<sub>2</sub> efflux rates in *Larix leptolepis*, *Pinus densiflora* and *P. rigitaeda* plantations in southern Korea

Received: 10 September 2015; Accepted: 18 May 2016

**Abstract:** This study compared soil CO<sub>2</sub> efflux rates in three adjacent coniferous plantations consisting of larch (*Larix leptolepis* Gordon), red pine (*Pinus densiflora* S. et Z.) and rigitaeda pine (*P. rigitaeda*: *P. rigida* Mill × *P. taeda* L.) species planted in the same year (1963). Soil CO<sub>2</sub> efflux, litter fall carbon (C) and root decomposition rates were measured with soil environmental factors for two years.

The mean annual soil CO<sub>2</sub> efflux rates (g CO<sub>2</sub> m<sup>-2</sup> hour<sup>-1</sup>) were the highest in rigitaeda pine (0.30), followed by red pine (0.27) and larch (0.24) plantations. An exponential regression of the CO<sub>2</sub> efflux rates against their corresponding soil temperatures showed a significant ( $P < 0.05$ ) relationship (red pine  $R^2 = 0.69$ ; rigitaeda pine  $R^2 = 0.67$ ; larch  $R^2 = 0.63$ ). The soil CO<sub>2</sub> efflux rates were negatively correlated with soil pH, but the soil water and soil organic C content were not significantly correlated with the CO<sub>2</sub> efflux rates. Soil CO<sub>2</sub>-C efflux rates were correlated by litter fall C inputs in the larch and red pine plantations, and the decomposition rates of 5–10 mm diameter roots in the larch plantation.

This result is useful in understanding potential responses of soil CO<sub>2</sub> efflux rates with changes in stand and soil environmental factors induced by different tree species.

**Keywords:** carbon cycling, carbon dynamics, litter fall, root decomposition, soil respiration

**Addresses:** C. Kim, Department of Forest Resources, Gyeongnam National University of Science and Technology, Korea, e-mail: [ckim@gntech.ac.kr](mailto:ckim@gntech.ac.kr)

J. Jeong, Centre for Environmental Risk Assessment and Remediation, University of South Australia, Adelaide, SA 5095, Australia, e-mail: [jeojy003@mymail.unisa.edu.au](mailto:jeojy003@mymail.unisa.edu.au)

\*Corresponding author

## Introduction

Quantitative evaluation of soil CO<sub>2</sub> efflux rates among different tree species is key to understanding the C cycling in forest ecosystems because tree species can have a significant influence on soil CO<sub>2</sub> efflux rates (Pangle & Seiler 2002; Lee & Jose 2003; Lee et al., 2010) due to their different nutrient re-

quirements (Dijkstra et al., 2009), litter fall fluxes (Vesterdal et al., 2012), litter decomposition mechanisms (Davidson & Janssens, 2006), fine root production (Valverde-Barrantes, 2007) and decomposition (Jeong & Kim, 2014). However, it is difficult to compare the soil CO<sub>2</sub> efflux rates of different tree species due to the potentially confounding effects of site macroclimatic and forest management fac-

tors (Kim et al., 2009). In addition, conflicting results have been reported regarding the influences of different tree species on soil CO<sub>2</sub> efflux rates. For example, tree species affected soil CO<sub>2</sub> efflux rates (Lee & Jose, 2003; Bréchet et al., 2009; Metcalfe et al., 2011), while others did not affect by different tree species (Valverde-Barrantes, 2007; Olsson et al., 2012). This disparity could be due to differences in soil environmental factors, such as nutrient availability, soil water content and soil temperature (Raich & Tufekcioglu, 2000), as well as stand environmental factors such as tree density, litter fall production and litter decomposition (Søe & Buchmann, 2005; Davidson & Janssens, 2006; Noh et al., 2010) induced by different tree species.

Although many studies generally describe the relationship between soil CO<sub>2</sub> efflux rates and biotic and abiotic factors, including soil temperature, nutrient supply, soil water, litter fall, litter decomposition and root decomposition, at regional and global scales (Davidson et al., 2000; Davidson & Janssens, 2006; Metcalfe et al., 2011), few studies describe the relationship between soil CO<sub>2</sub> efflux rates and soil and stand attributes (Søe & Buchmann, 2005; Noh et al., 2010), which are more important criteria that govern the soil CO<sub>2</sub> efflux rates of different tree species at a small, plot-based scale given similar site conditions.

Red pine (*Pinus densiflora* S. et Z.), rigitaeda pine (*P. rigitaeda*: *P. rigida* Mill × *P. taeda* L.) and larch

(*Larix leptolepis* Gordon) have been the three most important coniferous tree species planted throughout Korea during the last forty years. Red pine is the most important coniferous tree species occupying more than 23.5% (about 1.5 million ha) of Korean forest lands. Larch forests were also planted on about 600,000 hectares and rigitaeda pine showed good growth characteristics compared with rigida pine (*P. rigida* Mill.) in Korea. Information is needed to enable evaluation of the direction and rates of change in soil CO<sub>2</sub> efflux rates by various tree species because of C storage potential in plantations (Fox, 2000). The objectives of this study were as follows: 1) to quantify the impacts of established forest stands on soil CO<sub>2</sub> efflux rates in adjacent, even-aged larch and pine plantations under similar site conditions, and 2) to correlate soil CO<sub>2</sub> efflux rates with stand (litter fall C and root decomposition) and soil environmental factors (soil temperature, soil water content, soil pH, soil organic C concentration).

## Methods

The study was conducted in the Sambong National Exhibition Forests, administered by Korea Forest Service. The mean annual precipitation in this area is 1,322 mm and the annual mean temperature is 12.8°C. Experimental plots consisting of one decid-




Fig. 1. Location of the study site in Gyeongnam province, southern Korea. Box indicates three replicated plots in each plantation

Table 1. General characteristics in *Larix leptolepis*, *Pinus densiflora* and *P. rigitaeda* plantations

Plantation	Location	Elevation (m)	Stand density (trees ha <sup>-1</sup> )	DBH (cm)		Basal area (m <sup>2</sup> ha <sup>-1</sup> )	
				2006	2007	2006	2007
<i>L. leptolepis</i>	35°27'27" N	669	350 (17)	31.07	31.55	27.57	28.43
	127°38'30" E			(2.58)	(2.57)	(5.13)	(5.14)
<i>P. densiflora</i>	35°27'27" N	665	216 (58)	34.80	35.27	20.70	21.27
	127°38'28" E			(1.76)	(1.69)	(0.51)	(0.41)
<i>P. rigitaeda</i>	35°27'27" N	671	550 (29)	29.40	29.88	35.78	37.18
	127°38'33" E			(0.54)	(0.49)	(3.29)	(3.83)

N=3 plots per plantation. Values in parenthesis indicate standard error. DBH: diameter at breast height.

uous (larch) and two evergreen (red pine; rigitaeda pine) coniferous plantations were located adjacent to each other (Fig. 1). All three plantations were established in 1963 on northeast facing slopes (5–15°). The study sites were characterized by identical macroclimatic conditions and plantation age. The data for this study were collected from three replicated plots (20 m × 10 m) as an experimental unit of each tree plantation, with a small distance (<20 m) among plantation types because sampling bias due to differing site components can be reduced by using a sampling scheme with identical designs for adjacent sites (Oksanen, 2001). Sampling within each plantation was repeated monthly enough to interpret the magnitude and consistency of treatment effects because plantations of each tree species were not replicated (Hurlbert, 1984). Thus, the results from a single replicate need to be viewed and interpreted cautiously due to pseudoreplication (Hurlbert, 1984; Binkley, 2008).

Mean stand densities, diameters at breast height (DBH), and basal areas of the three plantations are shown in Table 1. More information about the soil conditions, understory vegetation, and nutrient dynamics in relation to litter fall and root decomposition of this study site can be found elsewhere (Kim et al., 2010; Kim et al., 2013; Jeong & Kim, 2014).

Soil CO<sub>2</sub> efflux rates were measured monthly *in situ* using an infrared gas analyzer system (Model EGM-4, PP systems, Hitchin, UK) equipped with a flow-through closed chamber (Model SRC-2, same manufacturer) from three randomly selected locations of each plot across each plantation, at time between 13:00 and 17:00, monthly from July 2006 through June 2008. The soil CO<sub>2</sub> efflux rates from each date were considered the midpoint of a sampling month and were used to calculate the monthly CO<sub>2</sub> efflux rates (Bowden et al., 2004).

To compare soil factors among the three plantations, soil temperature was measured monthly at 20 cm depth using a soil temperature probe (Model STP-1) attached to the EGM-4. Soil samples to measure soil water content were collected monthly at 20 cm depth using an Oakfield soil sampler. The core samples were placed in plastic bags, transported to a laboratory and dried in an oven for 48 hours at 105°C to quantify the soil gravimetric water con-

tent. The organic C concentration of the collected soil samples was determined by loss on ignition at 550°C for 4 hours, and soil pH was measured by glass electrode.

To measure the litter fall, three circular litter traps made of 1.5 mm nylon net with individual surface area of 0.25 m<sup>2</sup> were installed at randomly chosen locations in each plot (total 27 traps). Litter was collected at monthly intervals between July 2006 and June 2007. The collected litter from each trap was transported to a laboratory and then oven-dried at 65°C for 48 hours. The dried samples were then separated into needles and other components, and the mass of each portion was recorded. The C concentrations of litter samples were determined by elemental analyzer (Thermo Scientific, Flash 2000, Italy).

Root decomposition rates were estimated using the *in situ* buried root decay bag technique, which has been the most common approach for comparisons of root decay rates (Silver & Miya, 2001; Jeong & Kim, 2014), employing 15 cm × 15 cm nylon bags with < 0.5 mm mesh size. Fresh roots from each plantation were collected from a sampling point located randomly in each plot at approximately 15 cm mineral soil depth on July 15, 2006. For this study, the root system was defined as woody and non-woody roots < 2 mm, 2–5 mm and 5–10 mm in diameter together with their associated root tips. After collection, the roots were gently rinsed with tap water, sorted into these size classes, and air dried to constant mass at room temperature for 20 days. Root samples with an air-dried mass of 1 g were weighed to the nearest 0.01 g and placed in numbered bags. Sub-samples from each root type were also taken to determine oven-dried mass at 65°C for 48 hours. A total of 81 root bags (3 tree species types × 3 replication plots × 3 root diameter size classes × 3 replication samples) were incubated in the mineral soil layer of each plantation (installed 29 July 2006). The bags were inserted vertically into the mineral soil to a depth of 15 cm with a straight-blade shovel and collected after 378 days (11 August 2007). Collected bags were oven-dried at 65°C for 48 hours, cleaned by gentle brushing with a soft paintbrush to remove mineral soil and weighed to




Fig. 2. Monthly fluctuations of soil CO<sub>2</sub> efflux, soil temperature, soil pH, soil water content and organic carbon content in *Larix leptolepis*, *Pinus densiflora* and *P. rigitaeda* plantations. Mean and standard error are shown (n = 9). Different letters among plantations of each month represent a significant difference at P < 0.05



determine root mass loss rates. Information about the C and nitrogen dynamics in fine root decomposition processes of this study site can be found elsewhere (Jeong & Kim, 2014).

The exponential regressions between the soil CO<sub>2</sub> efflux rates and soil temperature were calculated as follows:

$$\text{Soil CO}_2 \text{ efflux rates} = B_0 e^{b_1 ST}$$

where  $B_0$  and  $b_1$  are the coefficients estimated through the regression analysis and  $ST$  is the soil temperature. The  $Q_{10}$  values were calculated by the equation  $Q_{10} = e^{10 \times b_1}$ .

The effect of tree species on soil CO<sub>2</sub> efflux rates and soil or stand environmental factors was evaluated by a nested analysis of variance (Kroodsma et al., 2011) with the PROC MIXED procedure in SAS (SAS Institute Inc., 2003). Three experimental units of each plantation were considered to be a possible covariate (a random effect) in the analysis. When significant differences at  $P < 0.05$  occurred, a comparison of the treatment means was compared using Tukey adjustment of least squares means. Pearson correlation coefficients were also calculated for the soil CO<sub>2</sub> efflux rates and the stand or soil environmental factors.

## Results

### Factors affecting the fluctuation of monthly soil CO<sub>2</sub> efflux rates

Temporal fluctuations in soil CO<sub>2</sub> efflux rates for the three plantations were closely related to soil temperature fluctuations. Soil CO<sub>2</sub> efflux rates reached their maximum values in July and August (Fig. 2) but declined during the fall (September and October) and spring (April and May). An exponential regression of monthly CO<sub>2</sub> efflux rates against the corresponding soil temperature at a depth of 20 cm (Fig. 3) was significant in all three plantations (larch:  $R^2 = 0.63$ ,  $P < 0.05$ ; red pine:  $R^2 = 0.69$ ,  $P < 0.05$ ; rigitaeda pine:  $R^2 = 0.67$ ,  $P < 0.05$ ; all plantations:  $R^2 = 0.66$ ,  $P < 0.05$ ).  $Q_{10}$  values in the three plantations were 4.76 in red pine, followed by 4.61 in rigitaeda pine and 3.96 in larch plantations.

Monthly fluctuations of CO<sub>2</sub> efflux rates were negatively correlated (Fig. 4) with the fluctuations of soil pH in all three plantations (larch:  $r = -0.71$ ,  $P < 0.01$ ; red pine:  $r = -0.79$ ,  $P < 0.01$ ; rigitaeda pine:  $r = -0.66$ ,  $P < 0.01$ ). In contrast to soil pH, the correlation coefficient of soil CO<sub>2</sub> efflux rates against soil water content was not significant ( $P > 0.05$ ) because no monthly fluctuation in soil water contents was ob-




Fig. 3. An exponential regression of CO<sub>2</sub> efflux rates against the soil temperature in *Larix leptolepis* (a), *Pinus densiflora* (b), *P. rigitaeda* (c) and all plantations (d)




Fig. 4. Correlation relationships between CO<sub>2</sub> efflux rates and soil pH, soil water content and soil organic carbon content in *Larix leptolepis*, *Pinus densiflora* and *P. rigitaeda* plantations (\*\*\*:  $P < 0.001$ )

served in the study stands. Soil organic C availability was not the major controlling factor ( $r = 0.20$ – $0.35$ ) of monthly fluctuations of soil CO<sub>2</sub> efflux in the three plantations.

### Factors affecting mean annual soil CO<sub>2</sub> efflux rates

The mean annual soil CO<sub>2</sub> efflux rates for the 2-year study period were significantly higher in rigitaeda pine ( $0.30 \text{ g CO}_2 \text{ m}^{-2} \text{ hour}^{-1}$ ) than in larch ( $0.24 \text{ g CO}_2 \text{ m}^{-2} \text{ hour}^{-1}$ ), but the rates were not significantly different between the red pine ( $0.27 \text{ g CO}_2 \text{ m}^{-2} \text{ hour}^{-1}$ ) and rigitaeda pine plantations (Fig. 5). In this study, abiotic factors such as the mean annual soil temperature, mean annual soil water content, mean annual soil pH and mean annual soil organic C content, which are regarded as the main factors influencing soil respiration, were significantly different among the three plantations (Fig. 5).

The soil CO<sub>2</sub>-C efflux rates were the highest in rigitaeda pine ( $8.48 \text{ Mg C ha}^{-1} \text{ year}^{-1}$ ), followed by red pine ( $7.23 \text{ Mg C ha}^{-1} \text{ year}^{-1}$ ) and larch ( $6.45 \text{ Mg C ha}^{-1} \text{ year}^{-1}$ ) plantations. There was a significant negative correlation between annual soil CO<sub>2</sub>-C efflux rates and total litter fall C fluxes during the study period in the larch and pine plantations (Fig. 6), although the efflux rates were not significantly correlated with the needle litter C fluxes in the rigitaeda pine plantation. There was no correlation between soil CO<sub>2</sub>-C efflux rates and decomposition rates from various sizes of roots in three plantations, except for the decomposition rates of 5–10 mm roots of larch (Fig. 7).




Fig. 5. Mean ( $n = 9$ ) annual soil CO<sub>2</sub> efflux rates, soil temperature, soil pH, soil water content and organic carbon content in *Larix leptolepis*, *Pinus densiflora* and *P. rigitaeda* plantations. Vertical bars indicate standard error. Different letters on the bars represent a significant difference at  $P < 0.05$




Fig. 6. Correlation relationships between soil CO<sub>2</sub>-C efflux rates and needle or total litterfall carbon fluxes in *Larix leptolepis*, *Pinus densiflora* and *P. rigitaeda* plantations (\*:  $P < 0.05$ ). Each data point is the mean value of 3 observations within each replication plot




Fig. 7. Correlation relationships between soil CO<sub>2</sub>-C efflux rates and decomposition rates of various root sizes in *Larix leptolepis*, *Pinus densiflora* and *P. rigitaeda* plantations (\*:  $P < 0.05$ ). Each data point is the mean value of 3 observations within each replication plot

## Discussion

### Factors affecting the fluctuation of monthly soil CO<sub>2</sub> efflux rates

Temporal fluctuations in monthly soil CO<sub>2</sub> efflux rates were affected by different tree species. Relatively low soil CO<sub>2</sub> efflux in the larch plantation could be attributed to the difference in soil temperature compared to the two pine plantations (Fig. 2). For example, soil temperatures in July and August 2006 were significantly lower for the larch (17.2°C) than for the rigitaeda pine (18.0°C) plantations. Several studies have reported significant effects of soil temperature on monthly soil CO<sub>2</sub> efflux rates because microbial activity increases with rising soil temperature (Buchmann, 2000; Davidson & Janssens, 2006).

An exponential increase in monthly soil CO<sub>2</sub> efflux rates with respect to soil temperature has been observed in other different forest types in Korea (Kim et al., 2009; Noh et al., 2010) because of the temperature dependency of microbial decay and root growth activity. Also, sensitivity of soil CO<sub>2</sub> efflux rates to soil temperature is commonly expressed by the coefficient  $Q_{10}$  values (Bond-Lamberty, 2011). However,  $Q_{10}$  values were not related to the difference of soil temperature among the three plantations because mean annual soil temperature was significantly lower for red pine than for rigitaeda pine plantations. The highest  $Q_{10}$  value at 20 cm soil depth in red pine could have resulted mainly from high C availability for microbial decay with adequate soil water content (Fig. 5). The  $Q_{10}$  values in this study are comparable to those observed in other pine forests in Korea,

which have been reported as 3.45–3.77 at 12 cm soil depth in red pine stands (Noh et al., 2010). The difference in  $Q_{10}$  values between this previous study and the current one may be due to the difference in soil depth at which temperature was measured. For example, in another study,  $Q_{10}$  values increased from 1.9 to 3.5 with increasing soil depth in European beech, Norway spruce, and Scots pine forests (Borken et al., 2002).

The strong correlation between monthly  $\text{CO}_2$  efflux rates and soil pH has been observed due to a seasonal trend in soil pH, which was depressed in the summer season when soil  $\text{CO}_2$  efflux reached its maximum value and was elevated in the winter season when soil  $\text{CO}_2$  efflux reached its minimum value. In contrast to soil pH, no monthly fluctuation in soil water contents was observed during the study period. Generally, the effect of soil water content on soil  $\text{CO}_2$  efflux is limited to either extremely high or low water levels, such as those observed during heavy rainfall events or prolonged droughts (Pangle et al., 2002; Tang et al., 2005), whereas seasonal soil temperature variation is minor (Davidson & Janssens, 2006). Soil organic C availability was not the major controlling factor ( $r = 0.20\text{--}0.35$ ) of monthly fluctuations of soil  $\text{CO}_2$  efflux in the three plantations because of large temporal and spatial variations of its component (Reth et al., 2005).

### Factors affecting mean annual soil $\text{CO}_2$ efflux rates

Mean annual soil  $\text{CO}_2$  efflux rates in forest ecosystems can be affected (Lee & Jose, 2003; Metcalfe et al., 2011) or relatively be unaffected (Valverde-Barrantes, 2007; Olsson, 2012) by different tree species. Many studies reported that differences in  $\text{CO}_2$  efflux between tree species could be attributed to differences in the conditions for decomposition of organic matter and root respiration, which are the two main sources of soil  $\text{CO}_2$  efflux rates (Metcalfe et al., 2007; Lee et al., 2010). For example, rigitaeda pine plantations showed favorable environmental conditions for microbial or root growth activity and organic matter decomposition, such as higher soil temperature (approximately  $0.61\text{--}0.89^\circ\text{C}$ ) and soil pH (approximately 0.16 units), compared to red pine or larch plantations (Fig. 5). Soil pH is an important variable that affects soil  $\text{CO}_2$  efflux rates because it is closely related to microbial activity and nutrient availability (Lee & Jose, 2003). In addition, the soil organic C content was significantly lower in rigitaeda pine plantations than in red pine plantations (Fig. 5), indicating an accelerated turnover for soil C pools due to relatively high soil temperature compared to the other two plantations. Low soil water content in rigitaeda pine

has also been attributed to enhanced diffusivity for  $\text{CO}_2$  due to increased air-filled pore space (Tang et al., 2005).

The soil  $\text{CO}_2\text{-C}$  efflux rates observed in these larch and pine plantations were comparable to those reported for other pine forests ( $6.21\text{--}7.45 \text{ Mg C ha}^{-1} \text{ year}^{-1}$ ) in central Korea (Noh et al., 2010) and for temperate evergreen coniferous forests ( $6.81 \text{ Mg C ha}^{-1} \text{ year}^{-1}$ ) and deciduous needle-leaved forests ( $6.41 \text{ Mg C ha}^{-1} \text{ year}^{-1}$ ) in the world's terrestrial vegetation (Raich & Schlesinger, 1992). There was a significant correlation between annual soil  $\text{CO}_2\text{-C}$  efflux rates and total litter fall C fluxes during the study period in the larch and red pine plantations. This result supports the previously reported finding that annual soil  $\text{CO}_2$  efflux rates were correlated to annual litter fall production across forest ecosystems (Yan et al., 2006). In this study, annual soil  $\text{CO}_2\text{-C}$  efflux rates were 3.9 times more than the total litter fall C fluxes in rigitaeda pine, 3.3 times more in larch and 3.1 times more in red pine plantations. The annual soil  $\text{CO}_2\text{-C}$  efflux rates observed in these three plantations were slightly higher than those reported for a global compilation of litter fall and soil  $\text{CO}_2$  efflux data, which showed an average value of 3.0 (Raich & Tufekcioglu, 2000).

There was no correlation between annual soil  $\text{CO}_2\text{-C}$  efflux rates and annual rates of decomposition from various sizes of roots in three plantations except for the decomposition rates of the larch roots, although heterotrophic soil respiration contributed approximately 66% of temperate coniferous forests in Korea (Lee et al., 2010). Other study reported that root decomposition in forest soils was related to soil  $\text{CO}_2$  efflux rates for approximately 3 months after initial incubation (Raich & Schlesinger, 1992) due to species-specific differences in fine root productivity, tissue chemistry, and turnover rates of organic matter (Søe & Buchmann, 2005).

### Conclusions

Three coniferous plantations in this study are capable of altering the soil  $\text{CO}_2$  efflux and various soil resource and environmental variables such as soil temperature, soil pH, soil water and soil organic C content. The increased soil  $\text{CO}_2$  efflux in the rigitaeda compared with the larch plantations was attributed to changes in soil temperature and soil pH that may have enhanced the decomposition rates of stored organic matter and microbial activity and root growth in the mineral soil layer. Soil  $\text{CO}_2\text{-C}$  efflux rates were influenced by total litter fall C fluxes of larch or red pine plantations, while these efflux rates were not affected by needle litter fall C fluxes and decomposing roots. This study indicates that soil  $\text{CO}_2$  efflux rates



in one deciduous and two coniferous plantations could be determined by changes in stand and soil resource factors caused by the different tree species when established at sites with similar macroclimatic and management conditions.

## Acknowledgments

This work was partially supported by the National Research Foundation Grant funded by the Korean Government (KRF-2006-331-F00024, 2010-0022193) and by Forest Science & Technology Projects (Project No. S211212L030320) provided by Korea Forest Service.

## References

- Binkley D (2008) Three key points in the design of forest experiments. *Forest Ecology and Management* 255: 2022–2023.
- Bond-Lamberty B, Bronson D, Bladyka E & Gower ST (2011) A comparison of trenched plot techniques for partitioning soil respiration. *Soil Biology and Biochemistry* 43: 2108–2114.
- Borken W, Xu YJ, Davidson EA & Beese F (2002) Site and temporal variation of soil respiration in European beech, Norway spruce, and Scots pine forests. *Global Change Biology* 8: 1205–1216.
- Bowden RD, Davidson E, Savage K, Arabia C & Steudler P (2004) Chronic nitrogen additions reduce total soil respiration and microbial respiration in temperate forest soils at the Harvard Forest. *Forest Ecology and Management* 196: 43–56.
- Bréchet L, Ponton S, Roy J, Freycon V, Coûteaux MM, Bonal D & Epron D (2009) Do tree species characteristics influence soil respiration in tropical forests; A test based on 16 tree species planted in monospecific plots. *Plant and Soil* 319: 235–246.
- Buchmann N (2000) Biotic and abiotic factors controlling soil respiration rates in *Picea abies* stands. *Soil Biology & Biochemistry* 32: 1625–1635.
- Davidson EA, Verchot LV, Cattânio JH, Ackermani IL & Carvalho JEM (2000) Effects of soil water content on soil respiration in forests and cattle pastures of eastern Amazonia. *Biogeochemistry* 48: 53–69.
- Davidson EA & Janssens IA (2006) Temperature sensitivity of soil carbon decomposition and feedbacks to climate change. *Nature* 440: 165–173.
- Dijkstra FA, West JB, Hobbie SE & Reich PB (2009) Antagonistic effects of species on C respiration and net N mineralization in soils from mixed coniferous plantations. *Forest Ecology and Management* 257: 1112–1118.
- Fox TR (2000) Sustained productivity in intensively managed forest plantations. *Forest Ecology and Management* 138: 187–202.
- Hurlbert SH (1984) Pseudoreplication and the design of ecological field experiments. *Ecological Monographs* 54: 187–211.
- Jeong J & Kim C (2014) Carbon and nitrogen status of decomposing roots in three adjacent coniferous plantations. *Annual of Forest Research* 57: 109–117.
- Kim C, Son Y, Lee WK, Jeong J & Noh NJ (2009) Influences of forest tending works on carbon distribution and cycling in a *Pinus densiflora* S. et Z. stand in Korea. *Forest Ecology and Management* 257: 1420–1426.
- Kim C, Jeong J, Cho HS & Son Y (2010) Carbon and nitrogen status of litterfall, litter decomposition and soil in even-aged larch, red pine and rigitaeda pine plantations. *Journal of Plant Research* 123: 403–409.
- Kim C, An HC, Cho HS & Choo GC (2013) Base cation fluxes and release by needle litter in three adjacent coniferous plantations. *Forest Science and Technology* 9: 225–228.
- Kroodsma DE, Byers BE, Goodale E, Johnson S & Liu WC (2011) Pseudoreplication in playback experiments, revisited a decade later. *Animal Behaviour* 61: 1029–1033.
- Lee KH & Jose S (2003) Soil respiration, fine root respiration, fine root production, and microbial biomass in cottonwood and loblolly plantations along a nitrogen fertilization gradient. *Forest Ecology and Management* 185: 263–273.
- Lee N, Koo JW, Noh NJ, Kim J & Son Y (2010) Autotrophic and heterotrophic respiration in needle fir and *Quercus*-dominated stands in a cool-temperate forest, central Korea. *Journal of Plant Research* 123: 485–495.
- Metcalf DB, Meir P, Aragão LEOC, Malhi Y, da Costa ACL, Braga A, Gonçalves PHL, de Athaydes J, de Almeida SS & Williams M (2007) Factors controlling spatio-temporal variation in carbon dioxide efflux from surface litter, roots, and soil organic matter at four rain forest sites in the eastern Amazon. *Journal of Geophysical Research* 112: G04001.
- Metcalf DB, Fisher RA & Wardle DA (2011) Plant communities as drivers of soil respiration: pathways, mechanisms, and significance for global change. *Biogeosciences* 8: 2047–2061.
- Noh NJ, Son Y, Lee SK, Yoon TK, Seo KW, Kim C, Lee WK, Bae SW & Hwang J (2010) Influence of stand density on soil CO<sub>2</sub> efflux for a *Pinus densiflora* forest in Korea. *Journal of Plant Research* 123: 411–419.

- Oksanen L (2001) Logic of experiments in ecology: is pseudoreplication a pseudoissue? *Oikos* 94: 27–38.
- Olsson BA, Hansson K, Persson T, Beuker E & Helmissaari HS (2012) Heterotrophic respiration and nitrogen mineralisation in soils of Norway spruce, Scots pine and silver birch stands in contrasting climates. *Forest Ecology and Management* 269: 197–205.
- Pangle RE & Seiler J (2002) Influence of seedling roots, environmental factors and soil characteristics on soil CO<sub>2</sub> efflux rates in a 2-year-old loblolly pine (*Pinus taeda* L.) plantation in the Virginia Piedmont. *Environmental Pollution* 116: 85–96.
- Raich JW & Schlesinger WH (1992) The global carbon dioxide flux in soil respiration and its relationship to vegetation and climate. *Tellus* 44B: 81–99.
- Raich JW & Tufekcioglu A (2000) Vegetation and soil respiration: correlations and controls. *Biogeochemistry* 48: 71–90.
- Reth S, Reichstein M & Falge E (2005) The effect of soil water content, soil temperature, soil pH-value and the root mass on soil CO<sub>2</sub> efflux- A modified model. *Plant and Soil* 268: 21–33.
- SAS Institute Inc (2003) SAS/STAT statistical software. version 9.1. SAS publishing, Cary, NC. USA.
- Silver WL & Miya RK (2001) Global patterns in root decomposition: comparisons of climate and litter quality effects. *Oecologia* 129: 407–419.
- Søe ARB & Buchmann N (2005) Spatial and temporal variations in soil respiration in relation to stand structure and soil parameters in an unmanaged beech forest. *Tree Physiology* 25: 1427–1436.
- Tang J, Qi Y, Xu M, Misson L & Goldstein AH (2005) Forest thinning and soil respiration in a ponderosa plantation in the Sierra Nevada. *Tree Physiology* 25: 57–66.
- Valverde-Barrantes OJ (2007) Relationships among litterfall, fine root growth, and soil respiration for five tropical tree species. *Canadian Journal of Forest Research* 37: 1954–1965.
- Vesterdal L, Elberling B, Christiansen JR, Callesen I & Schmidt IK (2012) Soil respiration and rates of soil carbon turnover differ among six common European tree species. *Forest Ecology and Management* 264: 185–196.
- Yan J, Wang Y, Zhou G & Zhang D (2006) Estimates of soil respiration and net primary production of three forests at different succession stages in South China. *Global Change Biology* 12: 810–821.