

Determination of Bispyribac Sodium 10 % SC (Herbicide) Residue Level in Straw, Grain and Soil Using HPLC Method

C. Tamilselvan*, S. John Joseph, V. Angayarkanni

Bioscience Research Foundation, Porur, Chennai - 600 116, Tamil Nadu, India

E-mail address: brfchennai@gmail.com

ABSTRACT

A field trial was conducted to evaluate the residues of Bispyribac sodium 10 % SC on rice crop during Kharif season 2013 at Kandikai in Thiruvallur District, Tamilnadu, India. Randomized block design was followed with three treatments for three replicates. Bispyribac sodium 10 % SC 200 g a.i./ha, 500 g a.i./ha and control (water spray) was sprayed using hand operated Maax battery sprayer with a spray volume of 300 litres per hectare at 15 days after transplantation of rice crop (*ADT 45*). At harvest, samples of grain, straw and soil were collected replicate wise from each treatment along with the control. These samples were stored in icebox and transfer to the laboratory under cooled condition for analysis. All the residues samples were analyzed for Bispyribac sodium content by a validated HPLC method at the minimum detectable concentration of 0.01 ppm. The result revealed that no detectable level of Bispyribac sodium in straw, grain and soil at harvest. The post treatment and pre-harvest interval was 60 days after transplantation.

Keywords: Bispyribac Sodium; Herbicide; Residue; Straw; Grain; Soil and High performance liquid chromatography

1. INTRODUCTION

Pesticides acted as the most widely used form of chemical in agriculture. They are used to kill any undesired organism interfering with agricultural production. A pesticide is a chemical or biological agent to control organisms that are considered to be harmful [1].

Pesticides are categorized into four main categories which are herbicides, fungicides, insecticides and rodenticides. Target pests can include insects, plants pathogen, weeds, mollusks, birds, mammals, and fish nematodes (roundworm) that destroy disease spreading properties or vectors for diseases. Although the use of pesticides is for human benefits, some have weaknesses such as potential toxicity to human and other animals. Over dose of pesticide cause soil contamination and undegraded pesticides remain as a residue in the soil [2].

Herbicide is the type of pesticide used to kill unwanted plants and weeds. Selective herbicides kill specific targets, while leaving the non-targeted planted crop without giving any harm. Herbicides are commonly used in agricultural production system throughout the world and Indian plantations are no exception.

The efficacy of herbicide in controlling weeds is important, its residual impact should also be considered for environmental safety for all processes which include soil adsorption, breakdown and degradation. The most of the soil-bound pesticides are due to adsorption of soil particles [3]. Transfer in soil is the movement of pesticide downward and spreading away from the target plants [4].

The objectives of the research are to determine residue of Bispyribac sodium which is newer herbicide widely used in agriculture for control major weeds in the rice crop. This research also aims to standardize and develop HPLC method for determination of Bispyribac sodium residues and their extraction, isolation and purification method in rice straw, grain and soil samples.

2. MATERIALS AND PREPARATION

Solvents (acetonitrile, silver nitrate, phosphoric acid and reference standard Bispyribac sodium) and HPLC water used for this study were Sigma Aldrich and Merck respectively.

2. 1. Instrumentation and analytical conditions

HPLC analyses were performed using the LC-10AT VP and SPD-10A UV-VIS Detector of Shimadzu with PC integrator. A reversed-phase column (Qualisil BDS 5u C18, Size-250 x 4.6 mm (i.d); particle size 5 μ m) was used. The room temperature was maintained at 32 °C. The mobile phase was a mixture of 0.1 % phosphoric acid: Acetonitrile (1:1) with a flow rate of 1.0 mL/min. at 248 nm.

2. 2. Sample Preparation Procedure

2. 2. 1. Sample Preparation Straw:

About 2 kg of straw was collected from each replication and cut into small pieces, mixed thoroughly by the method of cone and quartering and rejecting the opposite ends. The sample of 500 g was taken to the laboratory in icebox and stored in deep freezer until analysis.

2. 2. 2. Sample Preparation Grain

At harvest 2 kg of grain was collected after threshing from each replication and mixed thoroughly by the method of cone and quartering and rejecting the opposite ends. The sample 500 g was taken to the laboratory in icebox and stored in deep freezer until analysis.

2. 2. 3. Sample Preparation Soil

About 1 kg of soil was collected during harvest (0-15 cm) in 10 locations using soil auger and removed unwanted roots and debris from soil. After thorough mixing 500 g of sample was collected with quartering technique. The samples were packed in polyethylene bags with clear identification mark and taken to the laboratory in icebox and stored in deep freezer until analysis.

2. 2. 4. Reference Standard Preparation

A known weight of standard accurately was taken in 100 ml standard flask. Added sufficient volume of Acetonitrile, mixed well and made to the mark on the standard flask using the same solvent after dissolved the content completely. Allowed the standard solution to stand for 30 minutes at room temperature for equilibration. Stored in deep freezer and necessary dilutions were made from the stock whenever required.

2. 2. 5. Sample Storage

Stored the samples as such or if the bulk is too much, after the preparation of the sample. Depending upon the nature of the sample, keep the samples or their extracts either in deep freezer at 15 °C or in refrigerator until taken up for analysis. Ensure that the samples do not absorb or lose moisture during storage. Avoid undue long storage periods.

3. METHOD OF ANALYSIS

3. 1. Extraction of Straw and Grain

Exactly 50 g of representative sample was taken in a high speed blender; add 2 ml of 0.1 N silver nitrate solution and 25 ml mixture of 0.1 % phosphoric acid: Acetonitrile (1:1) solutions were added. Then it was homogenized for 2 minutes. The homogenized sample was extracted with 100 ml of methanol. Extraction process was done on an end-over-end mechanical shaker for a period of 15 minutes. Extract was filtered through whattman filter paper. The extraction process was repeated twice with the 75 ml of solvent after adding add 2 ml of 0.1 N silver nitrate solution and 25 ml mixture of 0.1 % phosphoric acid: Acetonitrile (1:1) solutions were added. The combined filtrate was concentrated to approximately 5ml of volume using rotary vacuum evaporator at temperature of 40 °C.

3. 2. Extraction of Soil

Exactly 50 g of representative sample was taken in a high speed blender; add 2 ml of 0.1 N silver nitrate solution and 25 ml mixture of 0.1 % phosphoric acid: Acetonitrile (1:1) solutions were added. Then it was homogenized for 2 minutes. The homogenized sample was extracted with 100 ml of methanol. Extraction process was done on an end-over-end mechanical shaker for a period of 15 minutes. Extract was filtered through whattman filter paper. The extraction process was repeated twice with the 75 ml of solvent after adding add 2 ml of 0.1 N silver nitrate solution and 25 ml mixture of 0.1 % phosphoric acid: Acetonitrile (1:1) solutions were added. The combined filtrate was concentrated to smaller volume (5 ml) using rotary vacuum evaporator at temperature of 40 °C.

3. 3. Calibration details

Different known concentrations of Bispyribac sodium (0.01, 0.1, 0.5, 1.0, 1.5, 2.0 & 2.5 ppm) were prepared in mobile phase by diluting the stock solution. Injected 20 µl of standard solution and the peak area resulting from the elution of compound strictly adhering to absorbance of peak retention was measured. A calibration curve has been plotted for concentration of the standards injected versus area observed and the curve was found linear up to the lowest concentration range 0.01 ppm. Thecalibration chart details are presented in the Figure 1 given below.




Figure 1. Calibration chart of Bispyribac sodium concentration Vs peak area.

3. 4. Recovery

Preparation of fortified standard solutions:

Bispyribac sodium was prepared taken in to three separate 100 ml standard flasks and added sufficient volume of mobile phase to dissolve the contents. To this fortified three levels of standards 0.5, 1.0, 1.5 ppm respectively. Made the standard flasks up to the mark-using Mobile phase and mixed well. The standard solution of 20 μ l was injected in the HPLC system continuously and recorded the respective peak areas (Table 1, 2 and 3). The typical HPLC chromatogram of fortified and blank samples of straw, grain and soil were presented in Figures 2 to 7.




Figure 2. Typical HPLC chromatogram of Rice straw sample-Blank.




Figure 3. Typical HPLC chromatogram of Rice straw sample-Fortified.

Table 1. Recovery of Straw.

Fortified Concentration (ppm)	Fortified Area (mV·Sec)	Recovery (%)	Mean ± SD
0.5	49.7352	95.84	96.06±0.27
1.0	97.4805	96.04	
1.5	149.4826	96.38	

Table 2. Recovery of Grain.

Fortified Concentration (ppm)	Fortified Area (mV·Sec)	Recovery (%)	Mean ± SD
0.5	49.1489	96.64	97.01± 0.39
1.0	96.7483	96.97	
1.5	148.3946	97.43	

Table 3. Recovery of Soil.

Fortified Concentration (ppm)	Fortified Area (mV·Sec)	Recovery (%)	Mean ± SD
0.5	50.2639	97.76	98.10 ± 0.33
1.0	97.6953	98.13	
1.5	147.1898	98.42	




Figure 4. Typical HPLC chromatogram of Rice grain sample-Blank.




Figure 5. Typical HPLC chromatogram of Rice grain sample-Fortified.




Figure 6. Typical HPLC chromatogram of Soil sample-Blank.




Figure 7. Typical HPLC chromatogram of Soil sample-Fortified.

4. RESULTS AND DISCUSSION

A field trial was conducted at Kandikai in Thiruvallur District, Tamilnadu, India to evaluate the residues of Bispyribac sodium 10 % SC on rice crop during Kharif season 2013. A randomized block design was followed with three replicates for three treatments. Bispyribac sodium 10 % SC @ 200 g a.i./ha, 500 g a.i./ha and water spray served as a control. After 15 days from the transplantation spraying was carried out by hand operated Maax battery sprayer with a spray volume of 300 litres per hectare. Rice variety 'ADT 45'

was grown in a net area of 25 sq·m per replication. The local agriculture practice of using as fertilizer and other plant production chemical was applied twice as per recommended dosage. The water was irrigated daily wise up to harvest and it was stopped before 5 days of harvest. At harvest samples of grain, straw and soil was collected replication wise from each treatment along with the control. The analyses of result revealed that no Bispyribac sodium residues could be deducted in straw, grain and soil at harvest in all treatment. The post treatment and pre-harvest interval was 60 days (Table 5, 6 and 7).

The validated HPLC method clearly indicated that the present method of the investigation is recommended for reduce analysis of any commodities. The validation is reflected the linearity, precision and accuracy which are very acceptable limit and high resolution without interference of matrix up to <0.01 ppm level (Figure 1).

The method validation ensures analyses creditability. In this study the parameters precision, linearity, accuracy, limit of detection (LOD) and limit of quantification (LOQ) were adopted.

4. 1. Linearity

For all method calibration curve were prepared on single day. The result obtained was plotted in the linear regression curve shown Figure 1. The curves are linear and gradually slope. The intercept and correlation coefficient is found to be 0.99.

The standard solution of Bispyribac sodium injected using HPLC model LC-10AT VP and SPD-10A UV-VIS Detector of Shimadzu. A reversed-phase column (Qualisil BDS 5u C18, Size-250 x 4.6 mm (i.d); particle size 5 µm) was used. The mobile phase was a mixture of 0.1 % phosphoric acid: Acetonitrile (1:1) with a flow rate of 1.0 mL/min and the elute was monitored at the wavelength of 248 nm. The column was maintained at a temperature of 25 °C. The retention time and peak area of the test substance were recorded at 10.14 min. The calibration curve for the HPLC analyses was conducted by plotting the peak area of normalization of Bispyribac sodium on Y-axis against concentration X-axis the obtained was shown in Figure 1. The peak a curve is straight line and linear.

4. 2. Precision




Figure 8. The peak area for Bispyribac sodium standards solution.

The intra-day precision method was repeated for analyses at three concentration levels. The RSD values clearly showed that variation was very minimal and accurate value of the respective concentration was obtained. The accuracy of an analytical method is the absence of test result true value which clearly reflected the recovery result. The mean recovery of straw, grain and soil were 96.06, 97.01 and 98.10 respectively (Table 1, 2 and 3).

Table 4. Summarization of Bispyribac sodium Standard Solution Data of HPLC.

Standard Solution	Retention time	Peak Area
Bispyribac sodium Standard Solution	10.140	52.8199

The chromatogram in Figure 8 showed the peak area for Bispyribac sodium standard solution containing Bispyribac sodium as the active ingredient. The peak on retention time was 10.140, which matched with the retention time of Bispyribac sodium compound; hence the standard solution was confirmed to contain certain level of Bispyribac sodium concentration. Table 4 summarized the peak area and retention time of Bispyribac sodium standard solution. The residue determination of Bispyribac sodium 10 % SC content in rice straw, grain and soil are presented in Table 5, 6 and 7.

Table 5. Residue Determination of Bispyribac sodium 10% SC Content in Rice Straw.

Treatment	Volume of the sample (ml)	Peak area of Bispyribac sodium in sample	Bispyribac sodium content (ppm)	Average of Bispyribac sodium content (ppm)
T1R1	5	ND	<0.01	<0.01
T1R2	5	ND	<0.01	
T1R3	5	ND	<0.01	
T2R1	5	ND	<0.01	<0.01
T2R2	5	ND	<0.01	
T2R3	5	ND	<0.01	
T3R1	5	ND	<0.01	<0.01
T3R2	5	ND	<0.01	
T3R3	5	ND	<0.01	

ND = Non-detectable (<0.01 ppm)

Table 6. Residue Determination of Bispyribac sodium 10 % SC Content in Rice Grain.

Treatment	Volume of the sample (ml)	Peak area of Bispyribac sodium in sample	Bispyribac sodium content (ppm)	Average of Bispyribac sodium content (ppm)
T1R1	5	ND	<0.01	<0.01
T1R2	5	ND	<0.01	
T1R3	5	ND	<0.01	
T2R1	5	ND	<0.01	<0.01
T2R2	5	ND	<0.01	
T2R3	5	ND	<0.01	
T3R1	5	ND	<0.01	<0.01
T3R2	5	ND	<0.01	
T3R3	5	ND	<0.01	

ND = Non-detectable (<0.01 ppm)

Table 7. Residue Determination of Bispyribac sodium 10 % SC Content in Soil Residue.

Treatment	Volume of the sample (ml)	Peak area of Bispyribac sodium in sample	Bispyribac sodium content (ppm)	Average of Bispyribac sodium content (ppm)
T1R1	5	ND	<0.01	<0.01
T1R2	5	ND	<0.01	
T1R3	5	ND	<0.01	
T2R1	5	ND	<0.01	<0.01
T2R2	5	ND	<0.01	
T2R3	5	ND	<0.01	
T3R1	5	ND	<0.01	<0.01
T3R2	5	ND	<0.01	
T3R3	5	ND	<0.01	

ND = Non-detectable (<0.01 ppm)

5. CONCLUSION

The residue analysis from the HPLC result shows that Bispyribac sodium could not be deducted in straw, grain and soil at harvest (<0.01 ppm). The post treatment and pre-harvest interval was 60 days after transplantation. Hence no residues found in rice crop and well below the allowable limit (0.01 ppm). The Bispyribac sodium standard and fortified sample chromatogram clearly indicate that the absence of matrix interference in the retention time of the substance and uniform retention time. From the result it is concluded, the developed method was clearly indicates that it is simple, fast and reliable. This method can be adopted by any researcher for their routine estimation. Further it is concluded that the absence of Bispyribac sodium in rice crop and soil substrate not pose any threat to the environment and any other commodities as it used in the recommended concentration.

References

- [1] Tomar RAS. Encyclopedia of Agricultural Chemistry, New Delhi, Anmol Publications PVT. LTD. 2010; Vol. 1: pp. 1-20.
- [2] M. Watts, Paraquat at Pesticide Action Network Asia & the Pacific 2011. Retrieved from <http://wssroc.agron.ntu.edu.tw/note/Paraquat.pdf>.
- [3] H. Muhamad, BS. Ismail, M. Sameni, N. Mat, *Environ Monitor Assessment* 49 (2010) 466-69.
- [4] USDA Natural Resources Conservation Services. Understanding Soil Risks and Hazards: Using Soil Survey to Identify the Areas with Risks and Hazards to Human Life and Property 1998, Retrieved from: <http://www.nature.nps.gov/geology/soils/Understanding%20Soil%20Risks%20and%20Hazards.pdf>.

(Received 27 May 2014; accepted 03 June 2014)