



EFFECT OF UV-B RADIATION ON ANTIOXIDATIVE ENZYME ACTIVITY IN CUCUMBER COTYLEDONS

MAGDALENA RYBUS-ZAJĄC* AND JAN KUBIŚ

*Department of Plant Physiology, University of Life Sciences in Poznań,
Wołyńska 35, 60-637 Poznań, Poland*

Received July 5, 2010; revision accepted October 22, 2010

Ultraviolet-B (UV-B) radiation negatively affects plant cells, causing reactive oxygen species (ROS) to be generated. To study the effects of increased UV-B exposure on antioxidant processes, we exposed germination-stage cucumber seedlings to increased ultraviolet radiation and analyzed hydrogen peroxide content and the activity of superoxide dismutase (SOD), catalase (CAT), guaiacol peroxidase (GPX) and syringaldazine peroxidase (SPX). Enzymatic antioxidant system activity generally increased after UV-B supplementation. Activation of CAT, GPX and SPX in cucumber cotyledons correlated positively with increases in SOD activity and hydrogen peroxide content. The results indicate that ROS accumulated despite higher engagement of the enzymatic antioxidant system, and that elevated UV-B radiation triggered oxidative stress in the cucumber cotyledons.

Key words: Catalase, guaiacol peroxidase, H₂O₂, superoxide dismutase, syringaldazine peroxidase.

INTRODUCTION

Ozone depletion in the stratosphere, caused by trace gases such as chlorofluorocarbons (CFCs) and NO_x, results in increased levels of ultraviolet-B radiation (UV-B, 280–315 nm) reaching the Earth's surface. Stratospheric ozone recovery was not expected before 2010, and will be slowed due to the effect of greenhouse gas emissions (Kakani et al., 2003). The decreased ozone levels are expected to recover to 1970 levels by 2050 (UNEP 2002). Current levels of UV-B during the cropping season are anywhere between 2 and 12 kJ m⁻² per day on the Earth's surface, an increase of 6–14% UV-B radiation over pre-1980 levels, and the effects of enhanced radiation may increase in future years (Madronich et al., 1998, UNEP 2002). This range of radiation has a multifaceted, most often negative, effect on crops. Plants must adapt to the deleterious effects of UV-B radiation because they depend on sunlight for photosynthesis and therefore cannot avoid exposure to UV-B radiation. The intensity of damage caused by stress depends on the plant species and its development phase, anatomical and morphological characteristics, environmental conditions, as well as the dose and duration of UV-B. In addition to directly

damaging DNA, UV-B radiation frequently triggers oxidative stress through the formation of reactive oxygen species (ROS), which in turn cause enhanced lipid and protein oxidation (Yanarelli et al., 2006a). Thus, oxygen toxicity is observed to increase in plant cells (Langebartels et al., 2002, Mittler 2002; Jain et al., 2004), accompanied by the formation of elevated amounts of ROS (Lamb and Dixon, 1997; Foyer and Noctor, 2003, 2005), which include superoxide anions (O₂^{•-}), hydrogen peroxide (H₂O₂), hydroxyl radicals (OH[•]) and singlet oxygen (¹O₂) (Scandalios 2002). The formation of H₂O₂ occurs in such physiological processes as photosynthesis and respiration. This metabolite participates indirectly in generation of structural barriers or serves signaling functions (Mittler, 2002; Neil et al., 1999, 2002; Veljavic-Jovanovic et al., 2002). The specific signaling pathway that controls the expression of the set of genes engaged in defense against UV-B is rapidly up-regulated (Agrawal et al., 2009). H₂O₂ is rather stable, but other ROS are relatively short-lived in cells (Asada, 1999). In response to ROS generation the antioxidant system is triggered; it includes superoxide dismutase (SOD), catalase (CAT) and peroxidases (POXs), as well as low-molecular antioxidants (Scandalios 2002). Superoxide dismutase generates

e-mail: magrybus@jay.up.poznan.pl

H_2O_2 , and catalase is one of the main enzymes that scavenge hydrogen peroxide. Peroxidases, present in almost every organism, catalyze oxidation of a wide range of substrates, mainly phenols, dependent on hydrogen peroxide decomposition. In plants they serve numerous physiological functions including hydrogen peroxide detoxification, lignin biosynthesis, hormonal signaling and stress response. Almost every organism contains POXs (Passardi et al., 2004).

Cucumber is a plant highly sensitive to stress factors. It is also one of the more common vegetable species. When grown in the field its cotyledons are at real risk of exposure at early stages of germination to stresses such as elevated UV radiation. We analyzed hydrogen peroxide levels and the activity of superoxide dismutase (SOD), guaiacol peroxidase (GPX), catalase (CAT), syringaldazine peroxidase (SPX) in cucumber cotyledons under exposure to elevated ultraviolet-B radiation. We assessed the sensitivity of the species at germination stage.

MATERIAL AND METHODS

The experimental material consisted of seedlings of cucumber cv. Dar grown on perlite under controlled growth conditions: 25°C, 14 h photoperiod, and 120 $\mu\text{mol m}^{-2} \text{s}^{-1}$ photon flux density (400–700 nm). Seven-day-old seedlings were subjected to UV-B irradiation supplied by Philips TL 20 W/01 RS lamps, max. 315 nm, 16 $\text{kJ m}^{-2} \text{day}^{-1}$ intensity, for 8 h per day (3.25 $\mu\text{mol m}^{-2} \text{s}^{-1}$ photon flux density) for 3–10 days.

Hydrogen peroxide concentration was assayed using the method of Messner and Boll (1994). Cotyledon samples were ground in 3 ml cold K phosphate buffer (pH 7.0) containing 0.02 g Polyclar AT. The homogenate was centrifuged for 25 min at 15,000 g at 4°C. The reaction mixture contained 1.5 ml extract, 0.15 ml K phosphate buffer (pH 7.0), 50 ml horseradish peroxidase (1 mg/1ml 100 mM K phosphate buffer, 60 units/mg) and 0.05 ml 0.05 M ABTS (2,2'-Azino-bis(3-ethylbenzothiazoline-6-sulfonic acid) diammonium salt). Absorbance increase according to the H_2O_2 level at 415 nm was measured after 3 min and compared to the standard curve of freshly prepared 0–30 nM H_2O_2 solutions in 100 mM K-phosphate buffer (pH 7.0). The results are given in nmol hydrogen peroxide (H_2O_2) per mg protein.

Superoxide dismutase activity was assayed using the method of Beauchamp and Fridovich (1971), which measures inhibition in photochemical reduction of nitroblue tetrazolium (NBT) at 560 nm. Cotyledons were ground in 3 ml cold solution containing 0.05 M Na phosphate buffer (pH 7.0), 1% PVP (polyvinylpyrrolidone), 1 mM EDTA (ethylenediaminetetraacetic acid) and 0.5 M

NaCl, and centrifuged for 25 min at 15,000 g at 4°C. The supernatant was used as enzyme extract. The 3 ml reaction mixture contained 0.05 M Na phosphate buffer (pH 7.8), 0.1 mM EDTA, 97 mM L-methionine, 120 mM riboflavin, 2 mM NBT and 30 μl enzyme extract. The reaction was carried out for 10 min under a fluorescent lamp. One unit of activity was estimated as the quantity of enzyme reducing absorbance to 50% of that of tubes lacking the enzyme. Total enzyme activity was expressed in units per mg protein.

For catalase and peroxidase activity the cotyledons were ground in 0.1 M phosphate buffer (pH 7.0) containing 0.5% polyethylene glycol (PEG 6.000) and 40 mg Polyclar AT, and centrifuged at 10,000 g for 15 min at 4°C, and the supernatant was used as enzyme extract.

Catalase activity was determined by measuring H_2O_2 consumption (Dhindsa et al., 1981) in reaction mixture containing 50 mM sodium phosphate buffer (pH 7.0), 15 mM H_2O_2 and extract, at 240 nm against a calibration curve. Activity was expressed as μM decomposed H_2O_2 per mg protein.

Guaiacol peroxidase was determined according to the method of Hammerschmit et al., (1982). The reaction mixture contained 0.2 mM guaiacol, 0.09 mM H_2O_2 and enzyme extract. H_2O_2 -dependent oxidation of guaiacol is followed by an increase of absorbance at 470 nm. Enzyme activity was calculated as the increase of absorbance (ΔA) per mg protein.

Syringaldazine peroxidase was determined according to Imberty et al. (1985). The reaction mixture contained 0.1 M phosphate buffer (pH 6.0), 0.5 mM H_2O_2 , syringaldazine (3.1 mg dissolved in 1 ml methanol and mixed 1:2 with dioxane) and enzyme extract. Enzyme activity was calculated as the absorbance increase at 530 nm (ΔA) per mg protein.

Protein content was determined by Bradford's (1976) method.

Photosynthesis rate was analyzed for whole cucumber seedlings by measuring CO_2 loss in a closed system using an AirTECH 2500-P CO_2 analyzer. The results were expressed as mg CO_2 consumed per g fresh weight.

Statistical analysis. The experimental data were subjected to one-way ANOVA, and significant differences between means were determined by Tukey's multiple range test ($p < 0.05$, $p < 0.01$).

RESULTS

Elevated UV-B radiation applied at the germination stage led to a time-dependent increase of hydrogen peroxide level in cucumber cotyledons, by 93%, 70% and 73% respectively at 5, 7 and 10 days of stress versus the controls (Fig. 1). Thus the




Fig. 1. Effect of progressive UV-B on hydrogen peroxide level in cucumber cotyledons. Significant difference between stressed plants and control: * $p < 0.05$, ** $p < 0.01$




Fig. 2. Effect of progressive UV-B on superoxide dismutase (SOD) activity in cucumber cotyledons. Significant difference between stressed plants and control: * $p < 0.05$, ** $p < 0.01$




Fig. 3. Effect of progressive UV-B on catalase (CAT) activity in cucumber cotyledons. Significant difference between stressed plants and control: * $p < 0.05$, ** $p < 0.01$




Fig. 4. Effect of progressive UV-B on guaiacol peroxidase (GPX) activity in cucumber cotyledons. Significant difference between stressed plants and control: * $p < 0.05$, ** $p < 0.01$

changes were already considerable at day 5 of stress. Superoxide dismutase activity also increased, by 14% (day 3 of stress), 70% (day 5), 56% (day 7) and 19% (day 10) versus the controls (Fig. 2). Except for the last day of the stress these changes were correlated with higher levels of hydrogen peroxide. In control plants the hydrogen peroxide level remained stable.

UV-B irradiation stress induced increased catalase activity in cucumber cotyledons as compared with the stable levels in the controls (Fig. 3). Under UV-B exposure it reached 16.1 units (day 3, vs 9.8 units in control), 16.5 units (day 5, vs 10.5), 12.7 units (day 7, vs 9.9) and 13.5 units (day 10, vs 10.1).

Guaiacol peroxidase activity at successive days of elevated UV-B irradiation amounted to 6.73 units

(day 3, vs 5.12 units in control), 15.31 units (day 5, vs 11.13), 14.58 units (day 7, vs 10.75) and 19.90 units (day 10, vs 9.28) (Fig. 4). Those levels in UV-B treated plants are 31%, 37%, 36% and 114% higher than the values from the respective controls.

Syringaldazine peroxidase activity in seedlings exposed to UV-B irradiation reached 18.62 units (day 3, vs 19.90 units in control), 32.56 units (day 5, vs 21.56), 36.62 units (day 7, vs 22.00), and 53.55 units (day 10, vs 38.78) (Fig. 5). Those levels in UV-B-treated plants are 51% (day 5), 66.5% (day 7) and 38% (day 10) higher than in the controls.

Under increased UV-B irradiation the photosynthesis rate decreased in a time-dependent manner versus the stable levels in the controls (Fig. 6); in UV-B treated cucumber seedlings, CO₂ assimilation




Fig. 5. Effect of progressive UV-B on syringaldazine peroxidase (SPX) activity in cucumber cotyledons. Significant difference between stressed plants and control: $p < 0.05$, ** $p < 0.01$

declined to 2.85 units (day 3, vs 3.08), 2.18 units (day 5, vs 2.85), 2.09 units (day 7, vs 3.02) and 2.82 units (day 10, vs 2.89).

DISCUSSION

In plants the most frequent effect of exposure to elevated UV radiation is oxidative stress and ROS overproduction, in which a significant role is played by hydrogen peroxide. H_2O_2 is an important signalling molecule (Foyer and Noctor, 2003, 2005) which induces further responses of plants to environmental stresses, both biotic and abiotic (Dat et al., 2000; van Breusegem et al., 2001; Neil et al., 2002; Bolkhina et al., 2003). It is generated via superoxide, presumably in a non-controlled manner, during electron transport processes such as photosynthesis and mitochondrial respiration (Alscher et al., 2002; Neill et al., 2002). Potential sources of hydrogen peroxide synthesis include NADPH oxidase, amine oxidase, oxalate oxidase and flavin-containing oxidases (Bolwell and Wojtaszek, 1997; Bolwell et al., 2002).

We noted enhanced hydrogen peroxide content after enhanced UV-B irradiation, with correlated increases of SOD activity. This suggests that under UV-B stress H_2O_2 was formed intensively as a result of enzymatic superoxide anion dismutation. Similar changes have been reported by Baumbusch et al. (1998), Dai et al. (1997), Kubo et al. (1999), Rybus-Zajac (2005), and Kubis and Rybus-Zajac (2008), but those investigations used plants during intensive vegetative growth. Rarely analyzed is the plant response at the seedling stage, especially at a very early stage which is connected with high oxidative activity and the involvement of cotyledons (Jain et al., 2004). There are reports on changes in SOD activity




Fig. 6. Effect of progressive UV-B on photosynthesis rate in cucumber cotyledons. Significant difference between stressed plants and control: $p < 0.05$, ** $p < 0.01$

and overproduction of H_2O_2 as a result of exposure to UV-B radiation in sunflower cotyledons (Yanarelli et al., 2006b) and *Picea asperata* seedlings (Han et al., 2009). In cucumber cotyledons irradiated with ultraviolet UV-C (254 nm) or UV-A+UV-B (280-360 nm), Watanabe et al. (2006) detected H_2O_2 accumulation and suggested that UV irradiation causes oxidative damage to DNA in plant cells.

Oxidative stress arises from an imbalance in the generation and decomposition of ROS, with more ROS (such as H_2O_2) being produced than are metabolized (Neill et al., 2002). Recent work has confirmed that hydrogen peroxide is a signaling molecule which mediates responses to abiotic stresses in plants; enhanced UV-B increased the efficiency of the antioxidant defense system, including UV-B-absorbing compounds and antioxidant enzyme activity (Han et al., 2009). The H_2O_2 accumulation we observed in cucumber cotyledons was accompanied by an increase in the activity of antioxidant system enzymes catalyzing reactions of hydrogen peroxide removal: CAT, GPX and SPX. Balakumar et al. (1997) reported increased catalase activity in tomato plants after UV-B irradiation. Increased peroxidase activity under the influence of UV-B irradiation has been reported frequently in, for example, barley (Mazza et al., 1999), sugar beet (Panagopoulos et al., 1990), sunflower cotyledons (Yanarelli et al., 2006b), long green cucumber cultivar (Kataria et al., 2007), wheat and mung bean (Agrawal and Ratore, 2007), spinach (Lei et al., 2008), *Picea asperata* seedlings (Han et al., 2009) and peanut (Tang et al., 2010). Sometimes, however, the activity of this enzyme is found to decline when the irradiation dose is too high (Rao et al., 1996). UV-B radiation induces increased activity of other antioxidant system enzymes of cucumber cotyledons as well; Kataria et

al. (2007) reported increased glutathione reductase and ascorbic acid peroxidase activity.

UV-B stress influenced antioxidant activity in the cell wall region in cucumber cotyledons, as seen in increased syringaldazine peroxidase levels. Intensification of cell wall component synthesis can increase cell wall rigidity, which in turn can boost tolerance to drought, UV-B and co-stresses (Clifford et al., 1998; Garcíá et al., 2000; Kubiś and Rybus-Zajac, 2008).

Enhanced UV-B reduced photosynthesis in our tested cucumber cotyledons. Previous work also showed negative effects of UV-B on photosynthetic processes, resulting in decreased biomass production (Correia et al., 2005; Agrawal and Rathore, 2007; Han et al., 2009).

The disturbances recorded in seedlings under elevated UV-B radiation indicate that accumulation of the signaling molecule hydrogen peroxide triggered further defense mechanisms, altering the antioxidant defense capacity and thus minimizing the negative impact of UV-B on the plants. The higher concentration of H₂O₂ induced oxidative stress and boosted the antioxidant defense system.

REFERENCES

- AGRAWAL SB, and RATHORE D. 2007. Changes in oxidative stress defense system in wheat (*Triticum aestivum* L.) and mung bean (*Vigna radiata* L.) cultivars grown with and without mineral nutrients and irradiated by supplemental ultraviolet-B. *Environmental and Experimental Botany* 59: 21–33.
- AGRAWAL SB, SINGH S, and AGRAWAL M. 2009. Ultraviolet-B induced changes in gene expression and antioxidants in plants. *Advances in Botanical Research* 52: 47–86.
- ALSCHER RG, ERTURK N, and HEATH LS. 2002. Role of superoxide dismutases (SODs) in controlling oxidative stress in plant. *Journal of Experimental Botany* 53: 1331–1341.
- ASADA K. 1999. The water-water cycle in chloroplasts: scavenging of active oxygen and dissipation of excess photons. *Annual Review of Plant Physiology and Plant Molecular Biology* 50: 601–639.
- BALAKUMAR T, GAYATHRI B, and ANBUDURAI PR. 1997. Oxidative stress injury in tomato plants induced by supplemental UV-B radiation. *Biologia Plantarum* 39: 215–221.
- BAUMBUSCH LO, EIBLMEIER M, SCHNITZLER JP, HELLER W, SANDERMANN H, and POLLE JA. 1998. Interactive effects of ozone low UV-B radiation on antioxidants in spruce (*Picea abies*) and pine (*Pinus silvestris*) needles. *Physiologia Plantarum* 104: 248–254.
- BEAUCHAMP C, and FRIDOVICH I. 1971. Superoxide dismutase: improved assays and an assay applicable to acrylamide gels. *Analytical Biochemistry* 44: 276–278.
- BOLKHINA O, VIROLAINEN E, and FAGERSTEDT KV. 2003. Antioxidants, oxidative damage and oxygen deprivation stress: a review. *Annals of Botany* 91: 179–194.
- BOLWELL GP, and WOJTASZEK P. 1997. Mechanisms for the generation of reactive oxygen species in plant defence – a broad perspective. *Physiological and Molecular Plant Pathology* 51: 347–366.
- BOLWELL GP, BINDSCHEDLER LV, BLEE KA, BUTT VS, DAVIES DR, GARDNER SL, GERRISH C, and MINIBAYEVA F. 2002. The apoplastic oxidative burst in response to biotic stress in plants: a three-component system. *Journal of Experimental Botany* 53: 1367–1376.
- BRADFORD MM. 1976. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Analytical Biochemistry* 72: 248–254.
- CLIFFORD SC, ARNDT SK, CORLETT JE, JOSHI S, SANKHLA N, POPP M, JONES HG. 1998. The role of solute accumulation, osmotic adjustment and changes in cell wall elasticity in drought tolerance in *Ziziphus mauritiana* (Lamk.) *Journal of Experimental Botany* 49: 967–977.
- CORREIA CM, MOUTINHO PEREIRA JM, COUTINHO JF, BJÖRN LO, and TORRES-PEREIRA JMG. 2005. Ultraviolet-B radiation and nitrogen affect the photosynthesis of maize: a Mediterranean field study. *European Journal of Agronomy* 22:337–347.
- DAI Q, YAN B, HUANG S, LIU X, PENG S, MIRANDA ML, CHAVEZ AQ, VERGARA BS, and OLSZYK DM. 1997. Response of oxidative stress defense systems in rice (*Oryza sativa* L.) leaves with supplemental UV-B radiation. *Physiologia Plantarum* 101: 301–308.
- DAT J, VANDENABEELE S, VRANOVA E, VAN MONTAGU M, INZE D, and VAN BREUSEGEM F. 2000. Dual action of the active oxygen species during plant stress responses. *Cellular and Molecular Life Sciences* 57: 779–795.
- DHINDSA RS, PLUMB-DHINDSA P, and THORPE TA. 1981. Leaf senescence: correlated with increased levels of membrane permeability and lipid peroxidation and decrease levels of superoxide dismutase and catalase. *Journal of Experimental Botany* 32: 93–101.
- FOYER CH, and NOCTOR G. 2003. Redox sensing and signaling associated with reactive oxygen in chloroplasts, peroxisomes and mitochondria. *Physiologia Plantarum* 119: 355–364.
- FOYER CH, and NOCTOR G. 2005. Oxidant and antioxidant signaling in plants: re-evaluation of the concept of oxidative stress in physiological context. *Plant, Cell and Environment* 29: 1056–1071.
- GARCÍA AL, FUENTES V, and NICOLÁS N. 2000. Interactive effect of nitrogen and long-term moderate water stress on water relations in tomato (*Lycopersicon esculentum* Mill.) plants. *Journal of Plant Physiology* 156: 563–566.
- HAMMERSCHMIDT R, NUCKLES EM, and KUĆ J. 1982. Association of enhanced peroxidase activity with induced systemic resistance of cucumber to *Colletotrichum lagenarium*. *Physiological Plant Pathology* 20: 73–82.
- HAN C, LIU Q, and YANG Y. 2009. Short-term effects of experimental warming and enhanced ultraviolet-B radiation on photosynthesis and antioxidant defense of *Picea asperata* seedlings. *Plant Growth Regulation* 58: 153–162.
- IMBERTY A, GOLDBERG R, and CATESSON AM. 1985. Isolation and characterization of *Populus* isoperoxidases involved in the last step of lignin formation. *Planta* 164: 221–226.
- JAIN K, KATARIA S, and GURUPRASAD KN. 2004. Oxyradicals under UV-B stress and their quenching by antioxidants.

- Indian Journal of Biochemistry and Biophysics* 42: 884–892.
- KAKANI VG, REDDY KR, ZHAO D, and SAILAJA K. 2003. Field crops responses to ultraviolet-B radiation: a review. *Agricultural and Forest Meteorology* 120: 191–218.
- KATARIA S, JAIN K, and GURUPRASAD KN. 2007. UV-B induced changes in antioxidant enzymes and their isoforms in cucumber (*Cucumis sativus* L.) cotyledons. *Indian Journal of Biochemistry and Biophysics* 44: 31–37.
- KUBIS J, and RYBUS-ZAJAC M. 2008. Drought and excess UV-B irradiation differentially alter the antioxidant system in cucumber leaves. *Acta Biologica Cracoviensia Series Botanica* 50: 35–41.
- KUBO A, AONO M, NAKALIMA N, SAJI H, TANAKA K, and KONDO N. 1999. Differential responses in activity of antioxidant enzymes to different environmental stresses in *Arabidopsis thaliana*. *Journal of Plant Research* 112: 279–290.
- LAMB C, and DIXON RA. 1997. The oxidative burst in plant disease resistance. *Annual Review Plant Physiology and Plant Molecular Biology* 48: 251–275.
- LANGEBARTELS CH, WOHLGEMUTH H, KSCHIESCHAN S, GRÜN S, and SANDERMANN H. 2002. Oxidative burst and cell death in ozone-exposed plants. *Plant Physiology and Biochemistry* 40: 567–575.
- LEI Z, MINGYU S, XIAO W, CHAO L, CHUNXIANG Q, LIANG C, HAO H, XIAOQING L, and FASHUI H. 2008. Antioxidant stress is promoted by nano-anatase in spinach chloroplasts under UV-B radiation. *Biological Trace Element Research* 121: 69–79.
- MADRONICH S, MCKENZIE LO, BJÖRN LO, and CALDWELL MM. 1998. Changes in biologically active ultraviolet radiation reaching the earth's surface. *Journal of Phytochemistry and Photobiology, B, Biology* 46: 5–19.
- MAZZA CA, BATTISTA D, ZIMA AM, SZWARCBERG-BRACCHITTA M, GIORDANO CV, ACEVEDO A, and SCOPEL AL. 1999. The effects of solar ultraviolet-B radiation on the growth and yield of barley are accompanied by increased DNA damage and antioxidant response. *Plant Cell Environmental* 22: 61–70.
- MESSNER B, and BOLL. 1994. Cell suspension cultures of spruce (*P. abies*): inactivation of extracellular enzymes by fungal elicitor-induced transient release of hydrogen peroxide (oxidative burst). *Plant Cell, Tissue and Organ Culture* 39: 69–78.
- MITTLER R. 2002. Oxidative stress, antioxidants and stress tolerance. *Trends in Plant Science* 7: 405–410.
- NEILL SJ, DESIKAN R, CLARKE A, and HANCOCK J. 1999. H₂O₂ signaling in plant cells. In: *Plant Responses to Environmental Stress*. BIOS Scientific Publishers, Oxford 8: 59–64.
- NEILL SJ, DESIKAN R, and HANCOCK J. 2002. Hydrogen peroxide signaling. *Current Opinion in Plant Biology* 5: 388–395.
- PANAGOPOULOS I, BORNMAN F, and BJÖRN LO. 1990. Effect of ultraviolet radiation and visible light on growth, fluorescence induction, ultra weak luminescence and peroxidase activity in sugar beat plants. *Journal of Photochemistry and Photobiology, B: Biology* 8: 73–87.
- PASSARDI F, LONGET D, PENEL C, and DUNAND C. 2004. The class III peroxidase multigenic family in rice and its evolution in land plants. *Phytochemistry* 65: 1879–1893.
- RYBUS-ZAJAC M. 2005. Oxidative stress generation in *Taxus baccata* leaves affected by *Pestalotiopsis funerea* Desm. under different light conditions. *Dendrobiology* 54: 51–56.
- RAO MV, POLIYATH G, and ORMROD D. 1996. Ultraviolet and ozone-induced biochemical changes in antioxidant enzymes of *Arabidopsis thaliana*. *Plant Physiology* 110: 125–136.
- SCANDALIOS JG. 2002. The rise of ROS. *Trends in Biochemical Sciences* 27: 483–486.
- TANG K, ZHAN JC, YANG HR, and HUANG WD. 2010. Changes of resveratrol and antioxidant enzymes during UV-induced plant defense response in peanut seedlings. *Journal of Plant Physiology* 167: 95–102.
- UNEP. 2002. Executive summary. Final of UNEP/WMO Scientific Assessment of Ozone Depletion: 2002. Prepared by the Scientific Assessment Panel of the Montreal Protocol on Substances that Deplete the Ozone Layer. UNEP, Nairobi (released 23 August 2002).
- VAN BREUSEGEM F, VRANOVA E, DAT JF, and INZE D. 2001. The role of active oxygen species in plant signal transduction. *Plant Science* 161: 404–414.
- VELJOVIC-JOVANOVIC S, NOCTOR G, and FOYER CH. 2002. Are leaf hydrogen peroxide concentrations commonly overestimated? The potential influence of artefactual interference by tissue phenolics and ascorbate. *Plant Physiology and Biochemistry* 40: 501–507.
- WATANABE K, YAMADA N, and TAKEUCHI Y. 2006. Oxidative DNA damage in cucumber cotyledons irradiated with ultraviolet light. *Journal of Plant Research* 119: 239–246.
- YANNARELLI GG, NORIEGA GO, BATLLE A, TOMARO ML. 2006a. Heme oxygenase up-regulation in ultraviolet-B irradiated soybean plants involves reactive oxygen species. *Planta* 224: 1154–1162.
- YANNARELLI GG, GALLEGOS SM, and TOMARO ML. 2006b. Effect of UV-B radiation on the activity and isoforms of enzymes with peroxidase activity in sunflower cotyledons. *Environmental and Experimental Botany* 56: 174–181.