



RELATIONSHIP BETWEEN APHID INFESTATION AND CHLOROPHYLL CONTENT IN FABACEAE SPECIES

SYLWIA GOŁAWSKA*, ROBERT KRZYŻANOWSKI, AND IWONA ŁUKASIK

Department of Biochemistry and Molecular Biology,
University of Natural Sciences and Humanities in Siedlce,
Prusa 12, 08-110 Siedlce, Poland

Received October 1, 2009; revision accepted November 10, 2010

We determined the chlorophyll *a* and *b* levels (SPAD readings) in uninfested leaves and in leaves after 7 and 17 days of aphid infestation in four Fabaceae species (*Pisum sativum* L., *Vicia faba* L., *Trifolium pratense* L., *Medicago sativa* L.). Feeding by pea aphids *Acyrtosiphon pisum* Harris (Hemiptera: Aphididae) caused significant loss of chlorophyll *a* and *b* in the infested plants. Uninfested leaves on both short- and long-infestation plants had significantly higher chlorophyll *a* and *b* than infested leaves.

Key words: chlorophyll *a+b*, Fabaceae, *Acyrtosiphon pisum*, SPAD values.

INTRODUCTION

One of the most serious pests of commercial Fabaceae crops is the pea aphid *Acyrtosiphon pisum* Harris (Homoptera: Aphididae) (Farag et al., 2007). The pea aphid damages crops directly and is a vector of more than thirty viral diseases, including bean yellow mosaic virus, red clover vein mosaic virus and pea streak virus (Barnett and Diachun, 1986; Jones and Proudlove, 1991). All viral diseases reduce yield of Fabaceae (Cuperus et al., 1982; Garlinge and Robertson, 1998).

Chlorophyll content is one of the most important parameters in the relationships between plants and herbivores. Chlorophyll levels change during plant development (Costa et al., 2001), and can alter in response to a wide variety of stresses (Fanizza et al., 1991; Samdur et al., 2000; Lawson et al., 2001). Chlorophyll catabolism is equal to global chlorophyll synthesis, and can be reduced by insect feeding, nutritional deficiencies and pathogen infections (Ni et al., 2002). Chlorophyll loss caused by herbivore feeding is not fully understood, although herbivory-caused chlorophyll loss has been described (Carbera et al., 1994; Ni et al., 2002; Heng-Moss et al., 2003). Two well-known aphid species causing chlorophyll loss are the Russian wheat aphid *Diuraphis noxia* (Mordvilko) (Hemiptera: Aphididae) on wheat *Triticum aestivum* L. (Burd

and Elliott, 1996) and the greenbug *Schizaphis graminum* (Rondani) on sorghum *Sorghum bicolor* (L.) Moench (Girma et al., 1998) and wheat. Leaf feeding by sap-feeding insects causes chlorosis and necrosis, leading to significant crop loss worldwide (Ni et al., 2001). Herbivore-caused leaf chlorosis in growing plants should be studied in detail, as chlorophyll fluorescence might prove useful as an indicator of plant responses to stressors including insect damage (Haile et al., 1999).

In this work we determined the concentrations of photosynthetic pigment (chlorophyll *a* and *b*) in uninfested and aphid-infested legumes. No similar studies have been conducted to assess the effect of feeding by this species on chlorophyll levels in such a wide range of hosts; this research represents an initial effort to characterize the effect *A. pisum* feeding has on chlorophyll *a+b* loss in legumes.

MATERIALS AND METHODS

PLANT MATERIAL

The experiments used four legume species: pea *P. sativum* L. var. Tulipan, vetch *V. faba* L. var. Jaga, clover *T. pratense* L. var. Bona, and alfalfa *M. sativa* L. var. Radius. Seed samples of alfalfa were obtained from the Plant Breeding and Acclimatization Institute (IHAR) in Radzików/Błonie (near Warsaw, Poland),

e-mail: sylwia@ap.siedlce.pl

TABLE 1. Total chlorophyll concentration (SPAD units) (means \pm SD) in uninfested (control) and infested legumes 7 days after pea aphid infestation

Plant species	Total chlorophyll (SPAD units)		t	d.f.	P
	Uninfested plants	Infested plants			
<i>Pisum sativum</i>	39.82 \pm 2.39	36.57 \pm 6.96	3.81	98	<0.001
<i>Vicia faba</i>	38.55 \pm 6.97	31.10 \pm 7.14	5.27	98	<0.001
<i>Trifolium pratense</i>	38.84 \pm 4.67	28.88 \pm 6.17	9.10	98	<0.001
<i>Medicago sativa</i>	30.89 \pm 5.15	26.10 \pm 6.09	3.57	73	<0.001

Student's *t*-test, comparing means between SPAD units in the legume plants.

and the others were bought from Horticultural Plant Breeding, Seed Production and Nursery in Ożarów Mazowiecki (Warsaw, Poland). Seed samples were germinated in a climate chamber and kept at 21 \pm 1°C and 70% relative humidity (RH) under a 16 h photoperiod. The plants were grown in plastic pots (7 \times 7 \times 9 cm) with fine garden soil commonly used for greenhouse experiments, one plant per pot. The plants were watered regularly and not additionally fertilized.

APHIDS

The pea aphids *Acyrtosiphon pisum* Harris used in the experiments came from stock culture kept at the University of Natural Sciences and Humanities in Siedlce, Poland. The aphids were reared on pea seedlings *P. sativum* L. var. Tulipan (Fabaceae) in an environmental chamber (21 \pm 1°C, 16 h photoperiod, 70% RH). They were transferred to the studied legumes for one generation (Apablaza and Robinson, 1967). Then the adult apterous females were used in the experiments.

ENTOMOLOGICAL OBSERVATIONS

The entomological observations were made on isolated plants in plastic cylinders (50 \times 50 \times 50 cm) in an environmental chamber (21 \pm 1°C, 16 h photoperiod, 70% RH). The experiment was initiated when the legume plants were 3 days old. Then 25 adult apterous females were placed on each plant of each legume species. At two intervals of infestation (7 and 17 days) the pea aphids (adult apterae, larvae and adult alatae) were counted on 10 plants of each legume.

SPAD METER READINGS

The chlorophyll content in tissues of single leaves of the legume plants (infested, and uninfested as control) was determined with a SPAD-502 meter (Minolta Corp., Ramsey, NJ). This instrument has a self-contained light source for uniform lighting over the sampled leaf surface, and two detectors, one sensitive to red light (645 nm) and the other sensitive to infrared radiation (790 nm). The sensors convert the light into

electrical currents for calculation of the SPAD value: $SPAD = A(\log(I_{or}/I_r) - \log(I_{of}/I_f)) + B$, where A and B are constant, and I_r and I_f are respectively the currents from red and infrared detectors with sample in place and with no sample in place (I_{or} and I_{of}) (Fanizza et al., 1991). Five SPAD readings were averaged for each leaf to represent one observation. The results represent average measurements of chlorosis for five leaves on ten plants of each legume.

STATISTICAL ANALYSIS

Comparisons of total chlorophyll concentration (SPAD units) between infested and control legume plants were subjected to two-tailed and unpaired Student's *t*-test. One-way ANOVA was carried out for number of aphids on the studied plants, followed by Duncan's test. Correlations between chlorophyll *a* and *b* content and number of aphids were calculated. All statistical analyses used Statistica for Windows v. 6.0 (StatSoft, 2003).

RESULTS

Average values of SPAD readings decreased under the stress of *A. pisum* feeding. For uninfested plants at the first measurement they ranged from 30.89 for *M. sativa* to 39.82 for *P. sativum*, and at the second measurement from 29.65 for *M. sativa* to 36.80 for *V. faba* (Tabs. 1, 2). Aphid infestation significantly reduced the level of chlorophyll *a* and *b* irrespective of the duration of infestation. At 7 days of infestation, *A. pisum* caused significant loss of chlorophyll *a* and *b* versus the values for uninfested plants of all studied legumes (Tab. 1). The decrease during the shorter period of infestation was greatest in vetch and clover (Tab. 1). At 17 days of infestation, *P. sativum* plants still showed a slight, nonsignificant decrease of chlorophyll *a+b*; at that interval the differences in chlorophyll *a+b* content between control and infested plants were significant for *V. faba*, *T. pratense* and *M. sativa* (Tab. 2). At 17 days the prolonged infestation produced stronger stress reactions in tissues of vetch and clover (Tab. 2).

TABLE 2. Total chlorophyll concentration (SPAD units) (means \pm SD) in uninfested (control) and infested legumes 17 days after pea aphid infestation

Plant species	Total chlorophyll (SPAD units)		t	d.f.	P
	Uninfested plants	Infested plants			
<i>Pisum sativum</i>	34.87 \pm 5.64	33.64 \pm 3.70	0.66	98	NS
<i>Vicia faba</i>	36.80 \pm 4.69	30.68 \pm 7.29	4.99	98	<0.001
<i>Trifolium pratense</i>	35.39 \pm 3.94	27.19 \pm 3.33	11.25	98	<0.001
<i>Medicago sativa</i>	29.65 \pm 3.88	23.80 \pm 2.70	6.75	73	<0.001

Student's *t*-test, comparing means between SPAD units in the legume plants.

TABLE 3. Abundance (means \pm SD) of pea aphids on the studied legume species

Plant species	Days after infestation*	
	7 days	17 days
<i>Pisum sativum</i>	134.70 \pm 15.05 a	213.60 \pm 38.84 a
<i>Vicia faba</i>	119.70 \pm 14.46 b	150.60 \pm 24.53 b
<i>Trifolium pratense</i>	10.90 \pm 2.33 c	2.60 \pm 0.70 c
<i>Medicago sativa</i>	6.50 \pm 1.35 c	1.70 \pm 0.95 c

*Values with different letters within columns differ significantly by Duncan's test at $p < 0.01$. Values are means from 10 plants on the sampling date for each legume.

There were clear differences in pea aphid population development between the studied legume plants. The number of pea aphids was highest on pea plants and lowest on alfalfa (Tab. 3).

The number of aphids on the studied plants was significantly correlated with SPAD readings only for alfalfa after seven days of infestation ($R = 0.96$; $p < 0.05$; Pearson correlations) (Fig. 1).

DISCUSSION

In this work, *A. pisum* infestation was shown to reduce chlorophyll levels in several species of the *Fabaceae* family. This indicates symptoms of chlorosis in the infested plants and adds important new data for *A. pisum*, an aphid species whose genome has been sequenced (Ollivier et al., 2010; IAGC, 2010) and which has been identified as a serious pest of legume crops. In *Vitis vitifera*, Fanizza et al. (1991) reported a drop in chlorophyll content under a different stress treatment. Leaves of stressed plants apparently synthesized less chlorophyll pigment. In our work the studied *Fabaceae* cultivars differed in the effect of pea aphid feeding on chlorophyll *a+b* concentrations. Other studies also show that responses differ. Rafi et al. (1996) found that *D. noxia* reduced chlorophyll levels in resistant cereals, while Heng-Moss et al. (2003) reported that total chlorophyll and




Fig. 1. Relationship between SPAD values and abundance of pea aphid on alfalfa 7 days after infestation ($y = -1.282 + 0.32166 * A\text{-SPAD}$, $R = 0.96$).

carotenoid concentrations differed among Beta wheat isolines in response to *D. noxia* feeding: infested Beta-*Dn2* plants had higher levels of chlorophylls and carotenoids than uninfested plants, but infested Beta-*Dn1* plants had the same chlorophyll and carotenoid levels as uninfested plants. Burd and Eliot (1996) found a significant decline in chlorophyll concentration in infested leaf tissue of *D. noxia*-susceptible wheat and barley, whereas total chlorophyll concentration was not significantly affected by *D. noxia* in resistant wheat or barley. In our study the amount of chlorophyll (as SPAD units) differed between infested and uninfested plants. The chlorophyll *a+b* concentration in uninfested *Fabaceae* plants was significantly higher than in aphid-infested *Fabaceae* plants. Aphid feeding adversely affected the plants and directly affected chlorophyll content. Interestingly, the chlorophyll concentration in pea plant tissues at 17 days of infestation was similar to the level in the respective uninfested plants. This indicates that aphid feeding may have less effect on chlorophyll loss in this species in the long term.

The exact mechanism by which aphids affect plant metabolism is not fully understood, but Heng-Moss et al. (2003) speculated that by feeding mainly on phloem tissue the aphids change the pH either on the luminal side of the thylakoid membrane, preventing the formation of zeaxanthin, or on the stromal side where regeneration of violaxanthin takes place. Burd and Elliott (1996) showed that aphid feeding could reduce protein synthesis, making photoinhibition irreversible as well as blocking electron transport on the acceptor site of the photosystem II reaction center, causing over-reduction in the system. One of two pathways of natural degradation of chlorophyll *a* is the oxidative bleaching pathway (Janave, 1997). Ni et al. (2001) showed that feeding by chlorosis-eliciting *D. noxia* or the non-chlorosis-eliciting bird cherry-oat aphid *R. padi* did not cause any changes in the oxidative bleaching pathway or chlorophyllase activity as compared with uninfested plants. However, Ni et al. (2002) showed that *D. noxia* feeding caused significant loss of chlorophyll *a* and *b* in the damaged regions: on two different sampling dates, undamaged regions of *D. noxia*-infested leaves showed significantly higher chlorophyll *a* and *b* concentrations than in uninfested leaves. *D. noxia*-infested wheat leaves showed significantly greater Mg-dechelataase activity than *R. padi*-infested and uninfested wheat leaves. Those assays of chlorophyll degradation enzymes indicated that *D. noxia* feeding significantly increased Mg-dechelataase activity in damaged and non-damaged leaf regions as compared with uninfested leaves, on both sampling dates. Mg-dechelataase activity in aphid-damaged regions was significantly higher than in undamaged leaf regions. This suggested that undamaged regions of *D. noxia*-infested leaves compensated for the pigment loss in the damaged regions, and that Mg-dechelataase activity changed dynamically from a localized response to a systemic response as the infestation period prolonged. Those assays clearly demonstrate the dynamic nature of plant responses to aphid feeding, which were initially localized and limited to the site of feeding, and then transformed to a whole-leaf response, indicating that *D. noxia* may elicit signaling chemical transduction between damaged and undamaged regions of infested leaves (Ni et al., 2002).

Haile et al. (1999) and Heng-Moss et al. (2003) found a significant decline of the photosynthetic rate in aphid-injured leaves and speculated that it may have resulted from increased synthesis of chemical defense compounds in response to herbivory. The decline in chlorophyll concentration found in our experiment may also be due to increased production of defensive compounds. Among the studied Fabaceae species the number of aphids was lowest on *M. sativa* plants, indicating that they are less attractive to *A. pisum*. Earlier work demonstrated that

alfalfa plants are resistant to *A. pisum*. Alfalfa contains numerous secondary plant metabolites, including carotenoids (Livingstone et al., 1980) and saponins (Oleszek, 1999, 2000; Oleszek et al., 1992; Stochmal et al. 2001a, b). They have been suggested as possible chemical defensive agents of alfalfa plants against generalist herbivores (Nozzolillo et al., 1997; Oleszek, 1999; Osbourn, 2003). Goławska et al. (2008) found that saponins play an important role in alfalfa defense against the pea aphid. Szykarczyk et al. (2001) showed that high-saponin alfalfa lines reduced pea aphid performance and phloem sap ingestion. Goławska et al. (2006) found big differences in pea aphid feeding behavior on alfalfa lines with low and high saponin content. Three alfalfa saponins (zanhic acid tridesmoside, 3-GlcA, 28AraRhaXyl medicagenic acid glycoside, 3-GlcA, 28AraRha medicagenic acid glycoside) were found to be inhibitors of *A. pisum* feeding (Goławska, 2007).

The present data on the effect of *A. pisum* feeding on chlorophyll *a+b* concentrations in Fabaceae tissues show changes in chlorophyll *a+b* concentrations in response to *A. pisum* feeding, and suggest the presence of a feeding-induced stress response in the studied legume species. SPAD measurements comprise a reliable, quick and nondestructive method useful not only for study of chlorophyll accumulation rates in different parts of the same leaf but also for determination of insect-plant interactions. Further research should investigate the mechanisms of acceptance/resistance for the Fabaceae species and explore the potential use of photosynthetic pigments as markers for identifying germplasm resistant to *A. pisum* and other chlorosis-causing insects.

REFERENCES

- APABLAZA HJV, and ROBINSON AG. 1967. Effect of three species of grain aphids (Homoptera: Aphididae) reared on wheat, oats or barley and transferred as adult to wheat, oats and barley. *Entomologia Experimentalis et Applicata* 10: 358–362.
- BARNETT OW, and DIACHUN S. 1986. Virus diseases of clovers: etiology and crop losses. In: Edwardson JR, and Christie RG [eds.], *Viruses Infecting Forage Legumes*, 625–675. Florida Agricultural Experiment Station, Gainesville.
- BURD JD, and ELLIOTT NC. 1996. Changes in chlorophyll *a* fluorescence induction kinetics in cereals infested with Russian wheat aphid (Homoptera: Aphididae). *Journal of Economic Entomology* 89: 1332–1337.
- CABRERA HM, ARGANDONA VH, and CORCUERA LJ. 1994. Metabolic changes in barley seedlings at different aphid infestation levels. *Phytochemistry* 35: 317–319.
- COSTA C, DWYER LM, DUTILLEUL P, STEWART DW, MA LB, and SMITH DL. 2001. Inter-relationships of applied nitrogen, SPAD, and yield of leafy and non-leafy maize genotypes. *Journal of Plant Nutrition* 24: 1173–1194.

- CUPERUS CW, RADCLIFFE EB, BARNES DK, and MARTEN GC. 1982. Economic injury levels and economic thresholds for pea aphid, *Acyrtosiphon pisum* (Harris) on alfalfa. *Crop Science* 1: 453–463.
- FANIZZA G, RICCIARDI L, and BAGNULO C. 1991. Leaf greenness measurements to evaluate water stressed genotypes in *Vitis vitifera*. *Euphytica* 55: 27–32.
- FARAG MA, HUHMANN DV, LEI Z, and SUMNER LW. 2007. Metabolic profiling and systematic identification of flavonoids and isoflavonoids in roots and cell suspension cultures of *Medicago truncatula* using HPLC-UV-ESI-MS and GC-MS. *Phytochemistry* 68: 342–354.
- GARLINGE J, and ROBERTSON D. 1998. Crop variety sowing guide for Western Australia. *Bulletin of Agriculture for Western Australia* 4341: 214–250.
- GIRMA M, KOFOID KD, and REESE JC. 1998. *Sorghum* germplasm tolerant to greenbug (Homoptera: Aphididae) feeding damages measured by reduced chlorophyll loss. *Journal of the Kansas Entomological Society* 71: 108–115.
- GOLAWSKA S. 2007. Deterrence and toxicity of plant saponins for the pea aphid *Acyrtosiphon pisum* Harris. *Journal of Chemical Ecology* 33: 1598–1606.
- GOLAWSKA S, LUKASIK I, and LESZCZYŃSKI B. 2008. Effect of alfalfa saponins and flavonoids on pea aphid. *Entomologia Experimentalis et Applicata* 128: 147–153.
- GOLAWSKA S, OLESZEK W, and LESZCZYŃSKI B. 2006. Effect of low and high-saponin of alfalfa on pea aphid. *Journal of Insect Physiology* 52: 737–743.
- HAILE FJ, HIGLEY LG, NI X, and QUISENBERRY SS. 1999. Physiological and growth tolerance in wheat to Russian wheat aphid (Homoptera: Aphididae) injury. *Environmental Entomology* 28: 787–794.
- HENG-MOSS TM, NI X, MACEDO T, MARKWELL JP, BAXENDALE FP, QUISENBERRY SS, and TOLMAY V. 2003. Comparison of chlorophyll and carotenoid concentrations among Russian wheat aphid (Homoptera: Aphididae)-infested wheat isolines. *Plant Resistance* 96: 475–481.
- IAGC (THE INTERNATIONAL APHID GENOMICS CONSORTIUM). 2010. Genome sequence of the pea aphid *Acyrtosiphon pisum*. *Plos Biology* 8(2): e1000313. doi:10.1371/journal.pbio.1000313
- JANAVE MT. 1997. Enzymatic degradation of chlorophyll in cavendish bananas: In vitro evidence for two independent degradative pathways. *Plant Physiology and Biochemistry* 35: 837–846.
- JONES RAC, and PROUDLOVE W. 1991. Further studies on cucumber mosaic virus infection of narrow-leaved lupin (*Lupinus angustifolius*): seed-borne infection, aphid transmission, spread and effects on grain yield. *Annals of Applied Biology* 118: 319–329.
- LAWSON T, CRAIGON J, TULLOCH AM, BLACK CR, COLLS JJ, and LANDON G. 2001. Photosynthetic responses to elevated CO₂ and ozone in field-grown potato (*Solanum tuberosum*). *Journal of Plant Physiology* 158: 309–323.
- LIVINGSTON AL, KOHLER GD, and KUZMICKY DD. 1980. Comparison of carotenoid storage stability in alfalfa leaf protein (Pro-Xan) and dehydrated meals. *Journal of Agriculture and Food Chemistry* 28: 652–656.
- NI X, QUISENBERRY SS, MARKWELL J, HENG-MOSS T, HIGLEY L, BAXENDALE F, SARATH G, and KLUCAS R. 2001. In vitro enzymatic chlorophyll catabolism in wheat elicited by cereal aphid feeding. *Entomologia Experimentalis et Applicata* 101: 159–166.
- NI X, QUISENBERRY SS, HENG-MOSS T, MARKWELL J, HIGLEY L, BAXENDALE F, SARATH G, and KLUCAS R. 2002. Dynamic change in photosynthetic pigments and chlorophyll degradation elicited by cereal aphid feeding. *Entomologia Experimentalis et Applicata* 105: 43–53.
- NOZZOLILLO C, ARNASON JT, CAMPOS F, DONSKOV N, and JURZYSTA M. 1997. Alfalfa leaf saponins and insects resistance. *Journal of Chemical Ecology* 23: 995–1002.
- OLESZEK W. 1999. Allelopathic significance of plant saponins. In: Macias FA, Galindo JCG, Molinillo JMG, and Cutler HG [eds], *Recent Advances in Allelopathy. A science for the Future*, 167–178. Servicio de Publicaciones, Universidad de Cadiz, Cadiz, Spain.
- OLESZEK W. 2000. Alfalfa saponins: chemistry and application. In: Bidlack WR, Omaye ST, Meskin MS, and Topham DK [eds], *Phytochemicals as Bioactive Agents*, 167–188. Technomic Publishing Company, Switzerland, Basel.
- OLESZEK W, JURZYSTA M, PŁOSZYŃSKI M, COLQUHOUN IA, PRICE KR, and FENWICK GR. 1992. Zanic acid tridesmoside and other dominant saponins from alfalfa (*Medicago sativa* L.) aerial parts. *Journal of Agriculture and Food Chemistry* 40: 191–196.
- OLLIVIER M, LEGEAI F, and RISPE C. 2010. Comparative analysis of the *Acyrtosiphon pisum* genome and EST-based gene sets from other aphid species. *Insect Molecular Biology* 19: 33–45.
- OSBORN AE. 2003. Molecules of interest, saponins in cereals. *Phytochemistry* 62: 1–4.
- RAFI MM, ZEMETRA RS, and QUISENBERRY SS. 1996. Interaction between Russian wheat aphid (Homoptera: Aphididae) and resistant and susceptible genotypes of wheat. *Journal of Economic Entomology* 89: 239–246.
- RICHARDSON AD, DUGAN SP, and BERLYN GP. 2002. An evaluation of noninvasive methods to estimate foliar chlorophyll content. *New Phytologist* 153: 185–194.
- SAMDUR MY, SINGH AL, MATHUR RK, MANIVEL P, CHIKANI BM, GOR HK, and KHAN MA. 2000. Field evaluation of chlorophyll meter for screening groundnut (*Arachis hypogaea* L.) genotypes tolerant to iron-deficiency chlorosis. *Current Science Bangalore* 79: 211–214.
- STATSOFT INC. 2003. Statistica (Data Analysis Software System), version 06. www.statsoft.com.
- STOCHMAL A, PIACENTE S, PIZZA C, DE RICCARDIS F, LEITZ R, and OLESZEK W. 2001a. Alfalfa (*Medicago sativa* L.) flavonoids. 1. Apigenin and luteolin glycosides from aerial parts. *Journal of Agriculture and Food Chemistry* 49: 753–758.
- STOCHMAL A, SIMONET AM, MACIAS FA, OLIVEIRA MA, ABREU JM, NASH R, and OLESZEK W. 2001b. Acylated apigenin glycosides from alfalfa (*Medicago sativa* L.) var. Artal. *Phytochemistry* 57: 1223–1226.
- SZYŃKARCZYK S, LESZCZYŃSKI B, OLESZEK W, and STASZEWSKI Z. 2001. Development of pea aphid, *Acyrtosiphon pisum* (Harris) on alfalfa lines varied in saponin content. In: Cichocka E, Goszczyński W, Leszczyński B, Ruszkowska M, Wojciechowski W [eds.], *Aphids and Other Homopterous Insects*, 121–130. Polish Academy of Sciences, Poland, Siedlce.