

## KARYOLOGY OF SEVEN *TRIFOLIUM* L. TAXA GROWING IN TURKEY

YASAR KIRAN<sup>1</sup>, AHMET SAHIN<sup>2</sup>, ISMAIL TÜRKÖĞLU<sup>3</sup>, MURAT KURSAT<sup>4</sup>,  
AND IFRAN EMRE<sup>5\*</sup>

<sup>1</sup>Department of Biology, Fırat University, 23169, Elazığ, Turkey

<sup>2</sup>Department of Secondary Science and Mathematics Education,  
Erciyes University, 38039, Kayseri, Turkey

<sup>3</sup>Department of Secondary Science and Mathematics Education,  
Fırat University, 23169, Elazığ, Turkey

<sup>4</sup>Department of Biology, Bitlis Eren University, 13000, Bitlis, Turkey

<sup>5</sup>Department of Primary Education, Fırat University, 23169, Elazığ, Turkey

Received March 17, 2010; revision accepted December 18, 2010

The paper reports karyomorphometric features and somatic chromosome numbers for seven *Trifolium* L. taxa. Three basic chromosome numbers are identified ( $x=5, 7, 8$ ). *Trifolium speciosum* Willd. and *Trifolium physodes* Stev. ex Bieb. var. *physodes* have somatic chromosome number  $2n=16$ ; *Trifolium campestre* Schreb., *Trifolium pratense* L. var. *americanum* Harz and *Trifolium sylvaticum* Gerard ex Lois. have  $2n=14$ ; *Trifolium bocconei* Savi has  $2n=10$ ; and *Trifolium repens* L. var. *repens* is tetraploid  $2n=32$ . All chromosomes have the centromeres at the median point (M) or median region (m) except for one chromosome of *Trifolium sylvaticum* Gerard ex Lois, which is submetacentric (sm). The only satellite observed was in the genome of *Trifolium bocconei* Savi.

**Key words:** Chromosome numbers, karyotype, *Trifolium* L.

### INTRODUCTION

The Leguminosae are the third largest family of flowering plants, with 727 genera and 19,325 species given (Lewis et al., 2005; Khatoon and Ali, 2006). *Trifolium* L. is one of the largest genera of the Leguminosae, comprising more than 250 species belonging to the tribe *Trifolieae*, of which ten are of considerable agricultural importance (Vizintin et al., 2006; Abberton, 2007). The genus name refers to the distinctive leaves, usually consisting of three leaflets (Salimpour et al., 2008). *Trifolium* originates in the Mediterranean area, Europe and the mountainous regions of Africa and Central, South and North America (Lange and Schifino-Wittman, 2000). The species of *Trifolium* are divided into eight sections: *Lotoidea*, *Paramesus*, *Mystillus*, *Vesicaria*, *Chronosemium*, *Trifolium*, *Tricocephalum* and *Involucrarium* (Lange and Schifino-Wittmann, 2000; Vizintin et al., 2006). All members of *Trifolium* are herbaceous perennials or annuals, frequently prostrate and rarely more than 50 cm in height (Ellison et al., 2006). In the flora of Turkey, *Trifolium* is repre-

sented by 118 taxa including 96 species, 22 subspecies and varieties (Davis, 1970; Davis et al., 1988).

Eighty percent of *Trifolium* species are diploid  $2n=16$ , and the basic chromosome number in the majority of examined species is  $x=8$  (Goldblatt, 1981; Zohary and Heler, 1984; Ellison et al., 2006). However, the basic chromosome numbers in *Trifolium* form an aneuploid series of  $x=5, 6$  and  $7$  (Zohary and Heler, 1984; Vizintin et al., 2006). Several studies reported additional chromosome numbers 11, 12 and 13 (Vizintin et al., 2006). Less than 20% of *Trifolium* species are polyploids and show ploidy variation (Ansari et al., 1999; Majumdar et al., 2004). In this study we examined chromosome morphology and determined chromosome number for seven *Trifolium* species growing in Turkey.

### MATERIALS AND METHODS

*Trifolium* specimens and seeds were collected from natural habitats in 2005 and 2006 and the seeds were used for karyological analysis. Voucher

e-mail: iemre@firat.edu.tr

TABLE 1. Somatic chromosome number, ploidy level, karyotype formula, chromosome length range, total karyotype length (TKL), asymmetry indexes ( $A_1$ ,  $A_2$ ) of Romero Zarco (1986) for the investigated *Trifolium* species

| Taxon  | 2n | Ploidy level | Karyotype formula | Chromosome length range ( $\mu\text{m}$ ) | TKL ( $\mu\text{m}$ ) | $A_1$ | $A_2$ |
|--|----|--------------|-------------------|---|-----------------------|-------|-------|
| <i>T. repens</i> L.var. <i>repens</i>                  | 32 | 4x           | 2M+14m            | 1.17–1.72                                 | 22.88                 | 0.18  | 0.09  |
| <i>T. speciosum</i> Willd.                             | 16 | 2x           | 8m                | 0.99–1.64                                 | 9.77                  | 0.22  | 0.16  |
| <i>T. campestre</i> Schreb.                            | 14 | 2x           | 1M+6m             | 1.13–1.73                                 | 9.79                  | 0.15  | 0.14  |
| <i>T. physodes</i> Stev. ex Bieb. var. <i>physodes</i> | 16 | 2x           | 8m                | 1.48–2.10                                 | 14.19                 | 0.20  | 0.13  |
| <i>T. pratense</i> L.var. <i>americanum</i> Harz       | 14 | 2x           | 1M+6m             | 1.20–1.79                                 | 9.97                  | 0.22  | 0.14  |
| <i>T. sylvaticum</i> Gerard ex Lois.                   | 14 | 2x           | 6m+1sm            | 1.13–1.81                                 | 9.44                  | 0.32  | 0.17  |
| <i>T. bocconeii</i> Savi                               | 10 | 2x           | 5m                | 1.53–2.31                                 | 972                   | 0.20  | 0.17  |

specimens (voucher numbers 5001–5007) and seeds are stored in the Firat University Herbarium (FUH).

Seeds were germinated at 25°C on moist filter paper in petri dishes. Actively growing root tips 1.5–2 cm in length were excised from the germinating seeds and pretreated with aqueous colchicine (0.05%) for 3 h at room temperature, then fixed with acetic alcohol (1:3) for 24 h and stored in 70% alcohol at 2–4°C. Then the root tips were rinsed in water and hydrolyzed with 1 N HCl for 10–18 min at 60°C and rinsed in running water for a minimum of 3–5 min. Feulgen staining of root tips was carried out for 1–2 h and root tips were put in 1% aceto-orcein for 5–10 min to improve staining. Finally, squash preparations were made in 1% aceto-orcein (Elci, 1982; Bairiganjan and Patnaik, 1989; Sahin, 1993).

Micrographs of good quality metaphase plates were taken using an Olympus BX51 microscope and an Olympus Camedia C-4000 digital camera. The long arm, short arm and total length of each chromosome were measured and the relative lengths, arm ratios, and centromeric indices were determined. Chromosome nomenclature follows Levan et al. (1964). The intra-chromosomal asymmetry index ( $A_1$ ) was calculated according to the formula given by Romero Zarco (1986), and the inter-chromosomal asymmetry index ( $A_2$ ) was calculated as the ratio of chromosome length to mean chromosome length.

## RESULTS AND DISCUSSION

Somatic chromosome numbers, ploidy level, karyotype formula, chromosome length ranges, total karyotype length (TKL), asymmetry index ( $A_1$ ,  $A_2$ ; Romero Zarco, 1986), karyomorphological parameters, metaphase chromosomes and haploid idiograms of the seven *Trifolium* species studied are presented in Tables 1 and 2 and Figures 1–8).

*Trifolium repens* L. var. *repens* (Tabs. 1, 2; Figs. 1, 8a) B7, Elaziğ, Baskil-Bolucuk village, 1580 m M. Kursat, 2006.

We found that the tetraploid number of the chromosomes for *T. repens* var. *repens* is  $2n=4x=32$  ( $x=8$ ). The karyotype of this species consists of 2 chromosomes with the centromere at the median point (M) and 14 chromosomes with centromeres in the median region (m). No satellite was observed. Our results are compatible with the findings of many other studies (Zohary and Heller, 1984; Taylor, 1987; Schifino-Wittmann and Moraes-Fernandes, 1988; Khatoon and Ali, 1993; Ansari et al., 1999). Some previous studies reported  $2n=16$ , 28 and 30 for *T. repens* (Williams and Verry, 1981; Pandey et al., 1987; Lavrenko et al., 1991).

*Trifolium speciosum* Willd. (Tabs. 1, 2; Figs. 2, 8b) B7 Elaziğ, Kamislik Mountain, 1260 m, I. Türkoğlu, 2006.

Its chromosome number was found to be  $2n=16$  ( $x=8$ ). All chromosomes have the centromere in median regions (m). No satellite was observed. The same chromosome number was noted by El Baba (1980) and Petrova and Kozuharov (1982).

*Trifolium campestre* Schreb. (Tabs. 1, 2; Figs. 3, 8c) B7 Elaziğ, Kamislik Mountain, 1760 m, I. Türkoğlu, 2006.

The chromosome number was determined as  $2n=14$  ( $x=7$ ). The karyotype of this species has 6 chromosomes with the centromere in the median region (m) and one with it at the median point (M). Satellites were not observed in the karyotype of this species. The same chromosome count was given by Lövkvist and Hultgård (1999), Vizintin et al. (2006) and Salimpour et al. (2008).

*Trifolium physodes* Stev. ex Bieb. var. *physodes* (Tabs. 1, 2; Figs. 4, 8d)

B7 Elaziğ, Baskil-Bolucuk village, 1480 m, M. Kursat, 2006.

The diploid number of the chromosomes was found to be  $2n=16$  ( $x=8$ ). All chromosomes are metacentric

TABLE 2. Karyomorphological parameters of *Trifolium* species

| Pair No.                                  | RL   | AR   | CI    | Type | Pair No.             | RL   | AR   | CI    | Type |
|---|------|------|-------|------|----------------------|------|------|-------|------|
| <i>T. repens</i> var. <i>repens</i>       |      |      |       |      | <i>T. speciosum</i>  |      |      |       |      |
| I   | 1.72 | 1.32 | 43.02 | m    | I                    | 1.64 | 1.27 | 43.90 | m    |
| II  | 1.64 | 1.56 | 39.02 | m    | II                   | 1.36 | 1.26 | 44.11 | m    |
| III                                       | 1.56 | 1.16 | 46.15 | m    | III                  | 1.24 | 1.38 | 41.93 | m    |
| IV  | 1.52 | 1.37 | 42.10 | m    | IV                   | 1.20 | 1.22 | 45.00 | m    |
| V   | 1.51 | 1.32 | 43.04 | m    | V                    | 1.15 | 1.30 | 43.47 | m    |
| VI  | 1.49 | 1.29 | 43.62 | m    | VI                   | 1.10 | 1.29 | 43.63 | m    |
| VII                                       | 1.44 | 1.25 | 44.44 | m    | VII                  | 1.09 | 1.18 | 45.87 | m    |
| VIII                                      | 1.42 | 1.15 | 46.47 | m    | VIII                 | 0.99 | 1.30 | 43.43 | m    |
| IX  | 1.41 | 1.31 | 43.26 | m    | <i>T. campestre</i>  |      |      |       |      |
| X   | 1.40 | 1.00 | 50.00 | M    | I                    | 1.73 | 1.30 | 43.35 | m    |
| XI  | 1.37 | 1.24 | 44.52 | m    | II                   | 1.57 | 1.18 | 45.85 | m    |
| XII                                       | 1.35 | 1.36 | 42.22 | m    | III                  | 1.46 | 1.11 | 47.26 | m    |
| XIII                                      | 1.33 | 1.18 | 45.86 | m    | IV                   | 1.34 | 1.18 | 49.25 | m    |
| XIV                                       | 1.31 | 1.14 | 46.56 | m    | V                    | 1.30 | 1.00 | 50.00 | M    |
| XV  | 1.24 | 1.00 | 50.00 | M    | VI                   | 1.26 | 1.29 | 43.65 | m    |
| XVI                                       | 1.17 | 1.48 | 40.17 | m    | VII                  | 1.13 | 1.26 | 44.24 | m    |
| <i>T. physodes</i> var. <i>physodes</i>   |      |      |       |      | <i>T. sylvaticum</i> |      |      |       |      |
| I   | 2.10 | 1.33 | 42.85 | m    | I                    | 1.81 | 1.58 | 38.67 | m    |
| II  | 1.98 | 1.23 | 44.94 | m    | II                   | 1.47 | 1.67 | 37.41 | m    |
| III                                       | 1.85 | 1.25 | 44.32 | m    | III                  | 1.37 | 1.24 | 44.52 | m    |
| IV  | 1.80 | 1.09 | 47.77 | m    | IV                   | 1.26 | 1.86 | 34.92 | sm   |
| V   | 1.79 | 1.20 | 45.25 | m    | V                    | 1.21 | 1.46 | 40.49 | m    |
| VI  | 1.67 | 1.19 | 45.50 | m    | VI                   | 1.19 | 1.33 | 42.85 | m    |
| VII                                       | 1.52 | 1.41 | 41.44 | m    | VII                  | 1.13 | 1.30 | 43.36 | m    |
| VIII                                      | 1.48 | 1.34 | 42.56 | m    | <i>T. bocconei</i>   |      |      |       |      |
| <i>T. pratense</i> var. <i>americanum</i> |      |      |       |      | I                    | 2.31 | 1.26 | 44.15 | m    |
| I   | 1.79 | 1.35 | 42.45 | m    | II                   | 2.27 | 1.45 | 29.07 | m*   |
| II  | 1.55 | 1.50 | 40.00 | m    | III                  | 1.91 | 1.17 | 46.07 | m    |
| III                                       | 1.47 | 1.40 | 41.49 | m    | IV                   | 1.70 | 1.32 | 42.94 | m    |
| IV  | 1.40 | 1.00 | 50.00 | M    | V                    | 1.53 | 1.15 | 46.40 | m    |
| V   | 1.33 | 1.29 | 43.60 | m    |                      |      |      |       |      |
| VI  | 1.23 | 1.32 | 43.08 | m    |                      |      |      |       |      |
| VII                                       | 1.20 | 1.22 | 45.00 | m    |                      |      |      |       |      |

\*Chromosome with satellite. Abbreviations: RL – relative length; AR – arm ration (L/S); CI – centromeric index (100S/TL); m – median region; sm – submedian region; M – median point. Chromosome pairs are Roman-numbered.

(m) and no satellites were observed in the karyotype. Giri et al. (1981) and Munoz-Rodriguez (1993a) also gave  $2n=16$  for this taxon.

*Trifolium pratense* L. var. *americanum* Harz (Tabs. 1, 2; Figs. 5, 8e)

B7 Elaziğ, Kamislik Mountain, 1430 m, I. Türkoğlu, 2006.


The somatic chromosome count was found to be  $2n=14$  ( $x=7$ ). The karyotype of this species consists of 6 chromosomes with the centromere in the median region (m) and one with it at the median point (M). Satellites were not observed in the karyotype of this taxon. The present results are in accord those of previous studies (Taylor, 1987; Nikolov, 1991;

Muñoz-Rodriguez, 1993b; Vizintin et al., 2006; Abberton, 2007), except for Salimpour et al. (2008) who gave  $2n=16$  and Armstrong and Cleveland (1970) who gave  $2n=28$ .

*Trifolium sylvaticum* Gerard ex Lois. (Tabs. 1, 2; Figs. 6, 8f)

B7 Elaziğ, Baskil-Bolucuk village, 1580 m, M. Kursat, 2006.

We found chromosome number  $2n=14$  ( $x=7$ ) for this species. Its karyotype consists of 6 chromosomes with the centromere in median regions (m) and one sub-metacentric chromosome (sm). No satellite was observed. Muñoz-Rodriguez (1993b) noted the same chromosome number for this species.




**Figs. 1–7.** Metaphase chromosomes of *Trifolium* species. **Fig. 1.** *T. repens* var. *repens* ( $2n=32$ ). **Fig. 2.** *T. speciosum* ( $2n=16$ ). **Fig. 3.** *T. campestre* ( $2n=14$ ). **Fig. 4.** *T. physodes* var. *physodes* ( $2n=16$ ). **Fig. 5.** *T. pratense* var. *americanum* ( $2n=14$ ). **Fig. 6.** *T. sylvaticum* ( $2n=14$ ). **Fig. 7.** *T. bocconei* ( $2n=10$ ). Bar=10  $\mu\text{m}$ .

*Trifolium bocconei* Savi (Tabs. 1, 2; Figs. 7, 8h) B7, Elaziğ, Kayabeyli village, 1450 m, M. Kursat, 2006. We determined the chromosome number of this species as  $2n=10$  ( $x=5$ ). All chromosomes are metacentric (m) and chromosome II has a satellite. Previous reports give two other chromosome numbers for *T. bocconei*:  $2n=12$  (Petrova and Kozuharov, 1982; Muñoz-Rodriguez, 1993b) and  $2n=14$  (Issolah and Abdelguerfi, 1999).

## CONCLUSION

The chromosome numbers we found are generally in accord with previous results, except for *T. repens* var.



**Fig. 8.** Haploid idiograms of *Trifolium* taxa. **(a)** *T. repens* var. *repens* ( $2n=32$ ), **(b)** *T. speciosum* ( $2n=16$ ), **(c)** *T. campestre* ( $2n=14$ ), **(d)** *T. physodes* var. *physodes* ( $2n=16$ ), **(e)** *T. pratense* var. *americanum* ( $2n=14$ ), **(f)** *T. sylvaticum* ( $2n=14$ ), **(g)** *T. bocconei* ( $2n=10$ ). Bar=10  $\mu\text{m}$ .

*repens*, *T. pratense* var. *americanum* and *T. bocconei*. Some previous studies reported chromosome counts differing from our findings (Armstrong and Cleveland, 1970; Williams and Verry, 1981; Pandey et al., 1987; Lavrenko et al., 1991; Salimpour et al., 2008). We determined three different basic chromosome numbers ( $x=5, 7$  and  $8$ ). *T. speciosum* and *T. physodes* have  $2n=16$  chromosomes. *T. campestre*, *T. pratense* var. *americanum* and *T. sylvaticum* have  $2n=14$ . *T. bocconei* has  $2n=10$ , and *T. repens* var. *repens* (tetraploid) has  $2n=32$ . Previous studies suggest that two diploid species, *T. occidentale* ( $2n=16$ ) and *T. nigrescens* ( $2n=16$ ) and the tetraploid species *T. uniflorum* ( $2n=4x=32$ ) are closely related to *T. repens* (Ansari et al., 1999; Jones et al., 2003). All studied chromosomes have the centromeres at the median point (M) and in the median region (m), except for the submetacentric (sm) chromosome of *T. sylvatica*. Only *T. bocconei* was found to have a satellite.

## ACKNOWLEDGEMENT

This work was financially supported by the Firat University Scientific Research Projects Unit (FÜBAP), project no. 1221.

## REFERENCES

- ABBERTON MT. 2007. Interspecific hybridization in the genus *Trifolium*. *Plant Breeding* 126: 337–342.
- ANSARI HA, ELLISON NW, READER SM, BADAEVAS ED, FRIEBES B, MILLER TE, and WILLIAMS WM. 1999. Molecular cytogenetic organisation of 5S and 18S-26S rDNA loci in white clover (*Trifolium repens* L.) and related species. *Annals of Botany* 83: 199–206.
- ARMSTRONG KC, and CLEVELAND RW. 1970. Hybrids of *Trifolium pratense* × *Trifolium pallidum*. *Crop Science* 10: 354–357.
- BAIRANGANJAN GC, and PATNAIK SN. 1989. Chromosomal evolution in Fabaceae. *Cytologia* 54: 51–64.
- DAVIS PH. 1970. *Trifolium* L. *Flora of Turkey and East Aegean Islands*, vol. 3. Edinburgh University Press, Edinburgh.
- DAVIS PH, MILL RR, and KIT T. 1988. *Flora of Turkey and East Aegean Islands*, vol.10. Edinburgh University Press, Edinburgh.
- EL BABA J. 1980. Contribution a l'etude cytotaxinomique des *Trifolium* du Liban. *Bulletin de la Société Botanique de France* 127: 53–58.
- ELCI S. 1982. Sitogenetikte gözlemler ve araştırma yöntemleri. Firat Üniversitesi Fen-Edebiyat Fakültesi, Biyoloji Elazığ.
- ELLISON NW, LISTON A, STEINER JJ, WILLIAMS WM, and TAYLOR NL. 2006. Molecular phylogenetics of the clover genus (*Trifolium* – Leguminosae). *Molecular Phylogenetics and Evolution* 39: 688–705.
- GIRI N, TAYLOR NL, and COLLINS GB. 1981. Chromosome numbers in some *Trifolium* species with a karyotype for *T. physodes*. *Canadian Journal of Genetics and Cytology* 23: 621–626.
- GOLDBLATT P. 1981. Cytology and the phylogeny of Leguminosae. In: Polhill RM and Raven PH [eds.], *Advances in Legume Systematics*, vol. 2, 427–464. Royal Botanic Gardens, Kew.
- ISSOLAH R, and ABDELGUERFI A. 1999. Chromosome numbers within some spontaneous populations of 10 *Trifolium* species in Algeria. *Caryologia* 52: 151–154.
- JONES ES, HUGHES LJ, DRAYTON MC, ABBERTON MT, MICHAELSON-YEATES TPT, BOWEN C and FORSTER JW. 2003. An SSR and AFLP molecular marker-based genetic map of white clover (*Trifolium repens* L.). *Plant Science* 165: 531–539.
- KHATOON S, and ALI AI. 1993. *Chromosome Atlas of the Angiosperms of Pakistan*, 232. Department of Botany, BCC&T press, University of Karachi,
- KHATOON S, and ALI SI. 2006. Chromosome numbers and polyploidy in the legumes of Pakistan. *Pakistan Journal of Botany* 38 (4): 935–945.
- LANGE O, and SCHIFINO-WITTMANN MT. 2000. Isozyme variation in wild and cultivated species of the genus *Trifolium* L. (Leguminosae). *Annals of Botany* 86: 339–345.
- LAVRENKO AN, SERDITOV NP, and ULLE ZG. 1991. Chromosome numbers in some species of vascular plants from the Pechoro-Ilychsky Reservation (Komi ASSR). *Botanicheskiy Zhurnal* 76: 473–476.
- LEVAN A, FREDGA K, and SANDBERG AA. 1964. Nomenclature for centromeric position on chromosomes. *Hereditas* 52: 201–220.
- LEWIS G, SCHRIRE B, MACKINDER B, and LOCK M. 2005. *Legumes of the World*. Royal Botanic Gardens, Kew.
- LÖVKVIST B, and HULTGÅRD UM. 1999. Chromosome numbers in south Swedish vascular plants. *Opera Botanica* 137: 1–42.
- MAJUMDAR S, BANERJEE S, and KUMAR DE K. 2004. Meiotic behaviour of chromosome in PMCs and karyotype of *Trifolium repens* L. From Darjeeling Himalaya. *Acta Biologica Cracoviensis Series Botanica* 46: 217–220.
- MUÑOZ-RODRIGUEZ AF. 1993a. Estudio biosistemático de *Trifolium* sect. *Vesicastrum* en la Península Ibérica. *Studia Botanica, Universidad de Salamanca* 11: 259–295.
- MUÑOZ-RODRIGUEZ A. 1993b. *Trifolium* sect. *Trifolium* II. Estudio cariológico. *Acta Botanica Malacitana* 18: 89–118.
- NIKOLOV NA. 1991. Chromosome numbers of Bulgarian angiosperms from North Pirin Mountain: Reserve "Bajuv Dupki-Dzindzirica". *Fitologija* 41: 70–75.
- PANDEY KK, GRANT JE, and WILLIAMS EG. 1987. Interspecific hybridization between *Trifolium repens* and *T. uniflorum*. *Australian Journal of Botany* 35: 171–182.
- PETROVA AV, and KOZUHAROV SI. 1982. Cytotaxonomic study of *Trifolium* L. in Bulgaria: I. *Fitologija* 19: 3–23. [In Bulgarian with English summary].
- ROMERO ZARCO C. 1986. A new method for estimating karyotype asymmetry. *Taxon* 35: 526–530.
- SAHIN A. 1993. Karyotype analysis of some *Lathyrus* L. species of Turkey I. *Turkish Journal of Botany* 17: 65–69. (In Turkish with an English abstract).
- SALIMPOUR F, SHARIFNIA F, HAJRASOLIHAI MS, and UKHNEH E. 2008. Chromosome counts and determination of ploidy levels in Iranian species of *Trifolium*. *Chromosome Botany* 3: 53–63.
- SCHIFINO-WITTMANN MT, and MORAES-FERNANDES B. 1988. Chromosome numbers, karyotypes and meiotic behavior of populations of some *Trifolium* (Leguminosae) species. *Revista Brasileira Genética* 11 (2): 379–390.
- STEBBINS GL. 1971. *Chromosomal Evolution in Higher Plants*. Edward Arnold, London.
- TAYLOR NL. 1987. Chromosome numbers of *Trifolium* species from Romania. *Notulae Botanicae Horti Agrobotanici* 17: 39–45.
- VIZINTIN L, JAVORNIK B, and BOHANEK B. 2006. Genetic characterization of selected *Trifolium* species as revealed by nuclear DNA content and ITS rDNA region analysis. *Plant Science* 170: 859–866.
- WILLIAMS EG, and VERRY IM. 1981. A partially fertile hybrid between *T. repens* and *T. ambiguum*, *New Zealand Journal of Botany* 19: 1–8.
- ZOHARY M, and HELLER D. 1984. *The Genus Trifolium*. Israel Academy of Sciences and Humanities, Jerusalem, Israel.