



Impact of *Enterococcus italicus* ONU547 on the growth and acclimatization of micropropagated *Rubus fruticosus* L. and *Paulownia tomentosa* Steud. plants to *ex vitro* conditions

NADIYA TYTARENKO^{1*}, NATALIA TESLIUK¹, ANDRII MERLICH¹, THOMAS HAERTLÉ², VOLODYMYR IVANYTSIA¹

¹Odesa I.I. Mechnikov National University, Odesa, Ukraine

²Institut National des la Recherche Agronomique, Nantes, France

Received: 10 January 2023; revised: 19 March 2023; accepted: 18 May 2023

Abstract

Clonal micropropagation is an effective method for plant reproduction, applicable in both scientific and industrial domains. However, a significant number of microclones are lost during the *ex vitro* acclimatization process. To address this, the introduction of beneficial microorganisms into the rhizosphere of micropropagated plants could have a positive effect on the survival rates and external characteristics of acclimatized plantlets. The aim of this study was to determine the protective and growth-promoting potential of *Enterococcus italicus* ONU547 and its effect on micropropagated plants during acclimatization. The antagonistic activity of the bacteria was determined using the agar block method. *Lepidium sativum* L. seeds were inoculated with bacterial suspensions at concentrations of 10⁶, 10⁷, and 10⁸ CFU/ml. Subsequently, the roots of the microclones were treated with suspensions of 10⁶ and 10⁷ CFU/ml, and biometric characteristics were measured. The results demonstrated antagonistic properties against various phytopathogenic fungi, including *Aspergillus niger*, *Cladosporium cladosporioides*, *Alternaria alternata*, *Alternaria tenuissima*, *Rhizoctonia cerealis*, *Penicillium expansum*, and *Paecilomyces variotii*. Inoculation of *L. sativum* L. seeds resulted in improved germination rates, increased root numbers, and enhanced root and shoot lengths. Similarly, the effects of the studied bacteria on *Rubus fruticosus* L. and *Paulownia tomentosa* Steud. during the acclimatization stage led to higher survival rates, increased shoot lengths, greater node numbers, and larger leaf areas. A concentration of 10⁷ CFU/ml was identified as optimal for inoculating the microclones. The findings indicate that *E. italicus* ONU547 holds promise for the inoculation of micropropagated plants during the acclimatization process. Further research is recommended to establish the specific interaction mechanisms between these bacteria and plants.

Key words: *Paulownia tomentosa*, *Rubus fruticosus*, *ex vitro* acclimatization, *Enterococcus italicus*, plant growth-promoting bacteria, antagonism

Introduction

Clonal micropropagation is a fast and convenient method of plant reproduction that enables the production of large amounts of planting material in a short period. This technique offers several advantages, including the ability to protect seedlings from viral and bacterial infections and the potential for year-round production of valuable plant material. Microclones occupy little space and, therefore, do not require a large area to grow like

ordinary seedlings. In addition, it is possible to propagate plants of certain species and varieties that are difficult to reproduce using traditional methods. However, the effectiveness of micropropagation varies among different plant species, requiring specific approaches and adjustments to standard procedures. The success of micropropagation largely depends on the quality of the resulting seedlings and their ability to adapt from environmental conditions to the natural environment du-

* Corresponding author: Odesa I.I. Mechnikov National University, Odesa, Ukraine; e-mail: tatti383@gmail.com

ring acclimatization (Tytarenko et al., 2020). This process often leads to the loss of a large amount of plant material due to reduced resistance to infections, physiological limitations (such as poorly developed cuticles, excessive transpiration, and low photosynthetic efficiency), and inadequate stress tolerance compared to conventional plants (Shekhawat et al., 2021). Alleviating acclimatization stress for micropropagated plants is currently a key priority in plant biotechnology.

Paulownia, also known as the Princess tree (*Paulownia tomentosa* (Thunb.) Steud.), is a rapidly growing tree crop that has gained popularity in horticulture worldwide. Its easy cultivation, decorative value, and rapid growth make it highly sought after for large batches of paulownia seedlings. Traditional methods of reproduction through seeds or layering are less effective and do not guarantee healthy plants. Conversely, clonal micropropagation offers a more efficient approach to rapidly obtaining a large number of healthy paulownia seedlings (Crisan and Petrus-Vancea, 2016).

Common blackberry (*Rubus fruticosus* L.) is a traditional fruit and berry crop that is becoming increasingly popular in Ukrainian gardens. Thornless varieties are particularly favored due to their convenience for harvesting and excellent fruit quality. Although blackberries can be easily propagated using conventional vegetative methods, clonal micropropagation generates larger numbers of planting materials with desired characteristics (Fathy et al., 2018).

For both paulownia and common blackberry, clonal micropropagation not only proves to be effective and convenient but also offers economic benefits as a method of reproduction. The profitability of this process can be further enhanced by optimizing certain stages of micropropagation, which are known to result in the loss of plant material. One such critical stage is the acclimatization to *ex vitro* conditions. Considering that micropropagated plants do not possess a well-established microbiota, and knowing the importance of the plant – microorganism interaction in nature, it is plausible to assume that inoculating microclones with beneficial bacteria during the acclimatization phase can increase the survival rate of plantlets in the soil and positively impact the biometric characteristics of the planting material.

Lactic acid bacteria have been recognized for their potential benefits to plants and are currently used as biocontrol agents against phytopathogens, plant growth

promoters, biopesticides, remediation agents, and biofertilizers (Raman et al., 2022). These bacteria can be isolated from various plant parts and the rhizosphere, with the greatest diversity typically observed near the roots (Fhoula et al., 2013). *Enterococcus* bacteria are occasionally found in the plant rhizosphere, but their role in plant associations remains poorly studied. Mostly, enterococci are considered as part of the transient microbiota (Svec et al., 2012; Kumar et al., 2016). They exist as members of rhizosphere consortia (Strafella et al., 2021) and can play a significant role in plant resistance against biotic and abiotic stresses (Panwar et al., 2016).

Enterococci have been observed to produce phytohormones (Lee et al., 2015) and bacteriocins with different activity spectra (Okkers et al., 1999; H-Kittikun et al., 2015; Almeida-Santos et al., 2021). They can also synthesize biosurfactants, volatile compounds, and other substances that protect plants against phytopathogens, making them potential biocontrol agents (Kumar et al., 2016; David and Onifade, 2018; Maany et al., 2019; Chaurasia et al., 2021; Diaz et al., 2021).

The search for and identification of potentially beneficial plant-associated enterococci is an important area of research aimed at supporting sustainable agricultural practices (Cesa-Luna et al., 2019). The use of growth-promoting and protective microorganisms in plant cultivation can reduce the reliance on chemical fertilizers and toxic plant protection agents while increasing the yield of important agricultural crops. Enterococci with growth-promoting properties can also find applications in various plant-related fields, including clonal micropropagation.

Enterococcus italicus ONU547 was isolated from plant material at Odesa I. I. Mechnikov National University (Merlich et al., 2019). This strain has been extensively studied in the field of plant–microbe interactions. Certain characteristics of *E. italicus* ONU547, such as bacteriocin and organic acid production, inhibition of phytopathogenic bacteria growth, and the formation of biofilms on plant surfaces, suggest the potential of these bacteria for enhancing the growth and protection of *R. fruticosus* and *P. tomentosa* plants (Merlich et al., 2017a, 2017b, 2019).

Therefore, the aim of this study was to evaluate the growth-promoting and protective potential of *E. italicus* ONU547 for *R. fruticosus* and *P. tomentosa* plants, parti-

cularly during the effect of micropropagated plants to *ex vitro* conditions. To achieve this purpose, the antagonistic activity of the bacteria against phytopathogenic fungi was determined, a seed germination assay was conducted using *Lepidium sativum* L. as a model plant, and acclimatization experiments were performed on blackberry and paulownia microclones by inoculating the roots with the bacterial strain.

Materials and methods

Plant material

Young, nonlignified shoots of paulownia (*P. tomentosa* (Thunb.) Steud.) and blackberry (*R. fruticosus* L. var. Thornfree) were chosen as the initial plant materials for establishing *in vitro* cultures. The donor material was collected between 2018 and 2021 from plants grown in the open ground at the Botanical Garden of Odesa I.I. Mechnikov National University. Nodal segments containing apical or axillary buds were used as the initial explants.

After surface sterilization and cutting the shoots, nodal segments ($n = 30$ for each plant species) were cultured on Murashige and Skoog (MS) medium (Murashige and Skoog, 1962) supplemented with 25 g/l sucrose and 1 mg/l of 6-benzylaminopurine (6-BAP). During the multiplication stage, the phytohormonal composition of the media was modified as follows: for blackberry plants, 1.5 mg/l 6-BAP and 0.5 mg/l naphthylacetic acid were used; for paulownia plants, 2 mg/l 6-BAP and 0.5 mg/l indolylacetic acid were used. The plant material ($n = 60$ for each plant species) was cultivated in an incubation room at 25°C, with a light intensity of 2500 lx, relative humidity of 56–70%, and an 8/16 h (night/day) photoperiod.

For the preacclimatization stage, 60 microclones (separately for blackberry and paulownia) were transferred to MS medium without phytohormones. During the acclimatization experiments, plants with a well-developed root system and similar heights ranging from 3.0 to 3.5 cm for *R. fruticosus* and 4.5 to 5.0 cm for *P. tomentosa* were used. At this stage, the microclones were divided into three groups, with each group consisting of 20 plants. Two groups were treated with different concentrations of bacteria, while the third group served as an uninoculated control. All experiments were repeated three times.

Microorganisms and incubation conditions

For this research, *E. italicus* ONU547, which is stored in the Ukrainian Collection of Microorganisms of D.K. Zabolotny Institute of Microbiology and Virology of the National Academy of Sciences of Ukraine (IMV NASU, Kyiv, Ukraine), was utilized. The bacteria were incubated in De Man, Rogosa, and Sharpe (MRS) liquid nutrient medium at 37°C for 24 h.

The tests strains of mycelial fungi, including *Aspergillus niger* F-16706, *Paecilomyces variotii* F-424, *Cladosporium cladosporioides* F-2235, *Penicillium expansum* F-575, *Alternaria alternata* F-16866, *Fusarium oxysporum* F-54201, were obtained from the collection of IMV NASU (Kyiv, Ukraine). The test strains of *Rhizoctonia cerealis* F-30 and *Alternaria tenuissima* F-24 were obtained from the collection of the Department of Microbiology, Virology, and Biotechnology of Odesa I.I. Mechnikov National University (Odesa, Ukraine). These selected species are commonly associated with plant diseases in field conditions and are also frequent contaminants in plant tissue culture (Omamor et al., 2007; El-Baky et al., 2021; Liu et al., 2021; Wang et al., 2021). The fungi were cultured on potato dextrose agar (PDA) at 28°C.

Antagonistic activity assay

The antagonistic activity of *E. italicus* ONU547 against mycelial fungi was determined using the agar block method. The bacteria were cultured overnight on MRS agar at 37°C. Subsequently, a block containing *E. italicus* ONU547 was cut from an agar plate and placed in a Petri dish on a PDA medium that had been freshly seeded with fungal spores. To obtain the fungal inoculum, *A. niger* F-16706, *P. expansum* F-575, and *F. oxysporum* F-54201 were incubated for 2 days until the sporulation stage, *P. variotii* F-424, *C. cladosporioides* F-2235, and *R. cerealis* F-30 were incubated for 3 days, and *A. alternata* F-16866 and *A. tenuissima* F-24 were incubated for 5 days. Fungal spore suspensions were prepared in a sterile physiological solution and inoculated on a PDA medium following the method described by Shobha and Kumudini (2012). After placing the bacterial blocks, the fungi were incubated at 28°C for 5 days, and the diameter of the growth inhibition zone was measured.

Study of the effect of *E. italicus* ONU547 on the growth of *L. sativum* L. plants

To determine the ability of *E. italicus* ONU547 to promote plant growth *in vitro*, a seed germination assay was conducted using watercress (*L. sativum* L.) as the model plant. The influence of the bacteria on the germination rate and biometric characteristics of watercress seedlings was assessed.

E. italicus ONU547 was cultured overnight in MRS liquid medium at 37°C with shaking at 180 rpm. The liquid culture was then centrifuged for 10 min at 3500 × g and washed twice with sterile water. Cell suspensions with concentrations of 10⁶, 10⁷, and 10⁸ CFU/ml were prepared for inoculation.

Watercress seeds ($n = 80$) were surface sterilized in a 3.5% sodium hypochlorite solution for 20 min and washed three times with distilled water. The seeds were then placed in the bacterial suspensions for an hour, with 20 seeds in each group. Afterward, the seeds were transferred to sterile moist chambers for germination. The control group seeds were exposed only to distilled water.

Observations were performed over a period of 10 days. The average length of the shoots and roots of the watercress seedlings was measured, along with the germination rate and the number of roots per plant.

Acclimatization of micropropagated plants to *ex vitro* conditions

The acclimatization process consisted of two stages: preacclimatization and subsequent acclimatization. During the preacclimatization stage, holes were made in the lids of containers housing the microclones. The purpose of these holes was to establish gas exchange with the atmosphere to facilitate transpiration processes. The diameter of the holes gradually increased over the 5 days, starting with 0.5 cm for the first 3 days and then enlarging to 1 cm for the last 2 days. This contributed to the establishment of transpiration processes in drier air compared with high humidity *in vitro* (Leite et al., 2021).

After the preacclimatization stage, the microclones were carefully removed from the MS medium and washed with sterile water. The roots of the microclones were then inoculated with *E. italicus* ONU547 by immersing them in a suspension of the bacteria at different concentrations (10⁶ and 10⁷ CFU/ml) for 1 h. In contrast, the roots of plants in the control group were

treated with distilled water. The microclones were subsequently planted in autoclaved soil mixed with 10% agroperlite and placed in an acclimatization room with controlled environmental conditions. The acclimatization room maintained a temperature range of 22–24°C, with a light intensity of 2500 lx, relative humidity of 56–70%, and an 8/16 h (night/day) photoperiod.

During the first 2 weeks, the microclones were irrigated with sterile tap water twice a week. Afterward, irrigation was continued using nonsterile tap water at the same frequency.

Observations were conducted over a period of 30 days after planting. On the 7th, 14th, and 30th days of cultivation, the survival rate of the microclones in the soil was measured, along with several biometric characteristics including average shoot length, number of nodes, and leaf area.

The survival rate was determined as the percentage of viable plants among all the plants in the experimental group. The average shoot length and number of nodes were manually measured. The ImageJ software, as described by Cosmulescu et al. (2020), was utilized to measure the average leaf area.

Statistical analysis

All experiments were performed in triplicate, and the data are presented as the mean values ± standard deviations. The plant material was randomly chosen and distributed. Statistical analysis was carried out using a one-way analysis of variance, followed by Duncan's multiple range test (DMRT) utilizing the Statistical Package for the Social Sciences software, version 29. Significance was determined at $P \leq 0.05$.

Results

Antagonistic activity of *E. italicus* ONU547 against phytopathogenic fungi

E. italicus ONU547 showed antagonistic activity against seven strains of mycelial fungi (Table 1). The largest zones of growth inhibition were observed when exposed to *A. niger* (inhibition zone diameter: 19.3 ± 1.2 mm), *C. cladosporioides* (18.7 ± 1.2 mm), and *A. alternata* (16.7 ± 1.2 mm). Antagonistic activities were also identified against *A. tenuissima*, *R. cerealis*, and *P. variotii*, with average growth inhibition zones ranging from 14.0 to 14.7 mm. Furthermore, *E. italicus* ONU547

Table 1. Antagonistic activity of *E. italicus* ONU547 against mycelial fungi

Fungi strains	Average growth inhibition zone diameter [mm]
<i>A. tenuissima</i> F-24	14.7 ± 1.2
<i>R. cerealis</i> F-30	14.0 ± 0.0
<i>F. oxysporum</i> F-54201	0.0 ± 0.0
<i>C. cladosporioides</i> F-2235	18.7 ± 1.2
<i>P. variotii</i> F-424	14.7 ± 1.2
<i>P. expansum</i> F-575	14.0 ± 0.0 *
<i>A. alternata</i> F-16866	16.7 ± 1.2
<i>A. niger</i> F-16706	19.3 ± 1.2

* not growth inhibition, but a delay in sporulation was observed

exhibited antagonism against *P. expansum*, characterized by delayed sporulation rather than the absence of mycelial growth. However, no antagonistic activity against *F. oxysporum* was observed.

Effects of *E. italicus* ONU547 on germination and seedlings growth of *L. sativum* L.

To determine the growth-promoting potential and potential phytotoxic effect of *E. italicus* ONU547, a screening assay was conducted using watercress as a model plant. The assay evaluated the effects on seed germination rate on the 2nd and 3rd days, as well as the average number of roots per seedling and the average length of shoots and roots on the 7th and 9th days of cultivation.

The results showed that the effects of different concentrations of *E. italicus* ONU547 on the mentioned characteristics varied compared to the control and between each other (Fig. 1).

At a concentration of 10⁷ CFU/ml, the bacteria had a positive impact on germination, increasing the germination rate by 7.8 and 9.3% on the 2nd and 3rd day, respectively. Bacteria at a concentration of 10⁶ CFU/ml did not cause statistically significant changes compared to the control on the 2nd day, but they increased the germination rate by 2.5% on the following day. On the other hand, at a concentration of 10⁸ CFU/ml, the bacteria exhibited a certain inhibitory effect on seed germination, with rates 5.0% lower than the control on the 2nd day and 6.3% lower on the 3rd day.

Overall, on the 3rd day of the experiment, a significant difference in germination rate was observed among all




Fig. 1. The effect of *E. italicus* ONU547 on the germination rate of watercress seeds on the 2nd and 3rd days of cultivation; control seeds ($n = 20$) received no treatment while treated seeds ($n = 20$ for each concentration) were inoculated with bacteria; means followed by a common letter are not significantly different by the DMRT at $P \leq 0.05$




Fig. 2. The average number of roots in watercress seedlings on the 7th and 9th days after inoculation with *E. italicus* ONU547; control seeds ($n = 20$) received no treatment while treated seeds ($n = 20$ for each concentration) were inoculated with bacteria; means followed by a common letter are not significantly different by the DMRT at $P \leq 0.05$

experimental groups, with the highest proportion of germinated seeds observed after inoculation with bacteria at a concentration of 10⁷ CFU/ml (Fig. 1).

Regarding the average number of roots on the 7th and 9th days of cultivation, a statistically significant difference was observed between all experimental groups and the control (Fig. 2).

On the 7th day, the seedlings exposed to 10⁷ CFU/ml of *E. italicus* displayed the highest average number of roots (13.4 ± 0.6 roots). Although the average number remained the same on the 9th day for these seedlings and increased by 0.9 roots in the control, the plants inoculated with 10⁷ CFU/ml of bacteria still had the highest number of roots among all experimental groups. Conversely, seedlings exposed to 10⁸ CFU/ml of *E. italicus*




Fig. 3. The average length of roots and shoots of watercress seedlings on the 7th day after inoculation with *E. italicus* ONU547; control seeds ($n = 20$) received no treatment while treated seeds ($n = 20$ for each concentration) were inoculated with bacteria; means followed by a common letter are not significantly different by the DMRT at $P \leq 0.05$




Fig. 4. The average length of roots and shoots of watercress sprouts on the 9th day after inoculation with *E. italicus* ONU547; control seeds ($n = 20$) received no treatment while treated seeds ($n = 20$ for each concentration) were inoculated with bacteria; means followed by a common letter are not significantly different by the DMRT at $P \leq 0.05$

showed inhibition of root formation on the 7th and 9th days, while those inoculated with 10⁶ CFU/ml showed a minimal level of positive influence (Fig. 2).

Significant differences were observed among all experimental groups in terms of the average length of roots and shoots on the 7th and 9th day (Fig. 3).

On the 7th day, seedlings inoculated with 10⁶ CFU/ml of *E. italicus* exhibited the longest average shoot length, reaching 24.4 ± 1.1 mm. Those treated with 10⁷ CFU/ml had an average shoot length of 19.7 ± 0.6 mm, while the group with 10⁸ CFU/ml showed shoot lengths close to the control values (11.7 ± 0.7 mm). The average length of roots was highest in seedlings exposed to 10⁷ CFU/ml of *E. italicus* (27.7 ± 1.0 mm), followed by those inocula-

ted with 10⁶ CFU/ml (25.3 ± 0.8 mm). The control group exhibited an average root length of 21.1 ± 1.1 mm, while the group treated with 10⁸ CFU/ml of *E. italicus* ONU547 had an average root length of 11.1 ± 0.7 mm (Fig. 3).

On the 9th day, the trends for most characteristics remained consistent. The highest average shoot length was observed in seedlings treated with 10⁶ CFU/ml and 10⁷ CFU/ml of *E. italicus* (21.8 ± 0.6 and 26.6 ± 1.0 mm, respectively). The group with 10⁸ CFU/ml showed shoot lengths 0.5 mm shorter than the control, while the root length in this group was almost two times shorter (14.4 ± 0.4 and 26.6 ± 1.0 mm). Seedlings treated with 10⁷ CFU/ml of *E. italicus* on the 9th day exhibited an average root length of 32.3 ± 0.7 mm, while the highest root length was observed in plants inoculated with 10⁶ CFU/ml of bacteria (35.9 ± 0.8 mm) (Fig. 4).

In conclusion, the impact of *E. italicus* ONU547 on plants was found to depend on the bacterial inoculum concentration. In this study, a concentration of 10⁸ CFU/ml exhibited phytotoxic effects, while concentrations of 10⁶ CFU/ml and 10⁷ CFU/ml demonstrated plant growth-promoting properties.

Effects of *E. italicus* on acclimatization of *P. tomentosa* and *R. fruticosus* microclones to ex vitro conditions

During the acclimatization process of micropropagated plants to nonsterile environmental conditions, *E. italicus* ONU547 demonstrated a favorable impact on the biometric characteristics of the seedlings, as well as on their survival rate in the soil, when compared to the control group. This effect was observed in both experimental plant species, namely blackberry and paulownia.

Survival rate

The survival calculations revealed that on the 30th day, the highest average number of viable *R. fruticosus* plants was observed in the group of microclones inoculated with 10⁷ CFU/ml of *E. italicus* ONU547 (67.4 ± 1.2%). This resulted in a 15.9% increase in survival rate compared to the control group. In the group of microclones exposed to 10⁶ CFU/ml of bacteria, the survival rate increased by 12.2% on the final day of observation (Table 2).

Similarly, the survival rate of paulownia plants showed the greatest improvement when inoculated with 10⁷ CFU/ml of bacteria. On the 30th day, it reached 73.7 ± 0.5%, which was 6.7% higher than the control group. With the inoculation of 10⁶ CFU/ml of *E. italicus*,

Table 2. The average survival rate of *Rubus fruticosus* and *Paulownia tomentosa* microclones in the soil after inoculation with *E. italicus* ONU547

Cultivation time [days]	Inoculum concentration [CFU/ml]	Survival rate of the microclones [%]	
		<i>Rubus fruticosus</i>	<i>Paulownia tomentosa</i>
7	control	73.0 ± 2.8 ^b	80.7 ± 0.5 ^b
	10 ⁶	79.6 ± 1.2 ^a	84.4 ± 2.7 ^{ab}
	10 ⁷	83.0 ± 1.2 ^a	88.9 ± 0.8 ^a
14	control	51.9 ± 2.4 ^c	71.9 ± 0.8 ^b
	10 ⁶	65.2 ± 0.9 ^b	75.6 ± 1.4 ^a
	10 ⁷	70.4 ± 0.9 ^a	77.4 ± 0.9 ^a
30	control	51.5 ± 1.6 ^b	67.0 ± 1.2 ^b
	10 ⁶	63.7 ± 1.2 ^a	72.6 ± 0.5 ^a
	10 ⁷	67.4 ± 1.2 ^a	73.7 ± 0.5 ^a

* means followed by a common letter are not significantly different by the DMRT at $P \leq 0.05$

the survival rate reached $72.6 \pm 0.5\%$, representing a 5.6% increase compared to the control (Table 2).

Furthermore, the study identified differences in the effects of different bacterial concentrations on the plants. For blackberry microclones, it was observed that on the 7th day of acclimatization, bacteria concentrations of 10^6 and 10^7 CFU/ml had a similar effect. On the 14th day, the concentration of 10^7 CFU/ml of *E. italicus* demonstrated greater effectiveness than 10^6 CFU/ml, but by the 30th day, both experimental concentrations showed a comparable impact. During the acclimatization of paulownia microclones, only *E. italicus* at a concentration of 10^7 CFU/ml exhibited a statistically significant effect on the survival rate on the 7th day. However, on the 14th and 30th days, the effects of bacteria at both concentrations were statistically similar (Table 2).

Therefore, the utilization of *E. italicus* ONU547 at concentrations of 10^6 and 10^7 CFU/ml led to an increase in the survival rate of *R. fruticosus* and *P. tomentosa* microclones compared to the control group.

Plant length

E. italicus ONU547 also had an impact on plant growth. The increase in the average length of blackberry and paulownia microclones during the acclimatization period from the 1st to the 30th day is illustrated in Figures 5 and 6.

Figure 5 demonstrates that the average length of blackberry plants naturally increased over time, with the

greatest growth observed in microclones treated with 10^7 CFU/ml of bacteria. On the 30th day, their length was, on average, 2.3 cm longer than that of the control plants. From the 7th day until the end of the observation period, the concentration of 10^7 CFU/ml exhibited a distinct effect on the average length of plants compared to both the control and the concentration of 10^6 CFU/ml. Although 10^6 CFU/ml of *E. italicus* also displayed an effect, the acceleration of growth was observed with a delay compared to the 10^7 CFU/ml concentration. A significant difference from the control was observed only on the 30th day when the plants in this group were longer than the control plants by an average of 0.9 cm. However, no discernible difference in the effect of this experimental group compared to the control was detected from the first to the 14th day of acclimatization.

During the acclimatization of paulownia plants, differences between the control group and inoculated microclones were also observed (Fig. 6). Plants exposed to *E. italicus* ONU547 at both concentrations exhibited greater shoot growth throughout the entire observation period. On the 30th day, the average length of paulownia plants inoculated with 10^6 and 10^7 CFU/ml of bacteria was 9.5 ± 0.5 and 8.7 ± 0.5 cm, respectively, while control plants had an average height of 7.6 ± 0.5 cm. Differences in the effect of the experimental concentrations of *E. italicus* compared to the control and among themselves were observed from the 7th day until the end of the experiment.




Fig. 5. The average length of the *Rubus fruticosus* microclones after inoculation with *E. italicus* ONU547; control plants ($n = 20$) received no treatment while treated plants ($n = 20$ for each concentration) were inoculated with bacteria; means followed by a common letter are not significantly different by the DMRT at $P \leq 0.05$




Fig. 6. The average length of the *Paulownia tomentosa* microclones after inoculation with *E. italicus* ONU547; control plants ($n = 20$) received no treatment while treated plants ($n = 20$ for each concentration) were inoculated with bacteria; means followed by a common letter are not significantly different by the DMRT at $P \leq 0.05$

Node number

Figure 7A depicts the average number of nodes in blackberry plants from the day of planting to the 30th day of acclimatization. The results obtained indicate a correlation between the number of nodes and the length of the plants, with a steady increase over time. Additionally, the difference between the inoculated plants and the control group also increased throughout the 30-day observation period. Notably, from the 7th day until the end of the experiment, a discernible difference in the effect of the experimental concentrations compared to the control group and among themselves was observed. The control plants consistently exhibited the lowest mean values over the course of the 30 days. The use of 10^6 CFU/ml of *E. italicus* resulted in intermediate results, with 7.5 ± 0.5 nodes observed on the 30th day. The

highest number of nodes (an average of 8.8 ± 0.4 nodes/plant) was found in the group of plants inoculated with 10^7 CFU/ml of bacteria.

Similarly, for paulownia microclones, *E. italicus* ONU547 also increased the number of nodes in plants starting from the 7th day of acclimatization (Fig. 7B). However, in this case, both concentrations used in the experiment exhibited varying effects. On the 7th day of observation, the highest average number of nodes was observed in plants inoculated with 10^6 CFU/ml of *E. italicus*. On the 14th day, the effects of both concentrations were determined to be the same, and on the last day, the highest average number of nodes was found in microclones inoculated with 10^7 CFU/ml of bacteria (7.7 ± 0.5 nodes). Therefore, *E. italicus* ONU547 had a positive effect on the number of nodes in paulownia plants at both experimental concentrations.

Leaf area

Differences between the experimental and control plants were also noticeable in terms of the average leaf area in both blackberry and paulownia plants (Fig. 8A and Fig. 8B).

Measurements of the average leaf area in blackberry plants during the acclimatization experiment differed from those of the control group. However, on the last day of observation, both experimental concentrations of bacteria showed similar results (Fig. 9). On the 7th day of acclimatization, a statistically significant difference in the average leaf area of blackberry plants compared to the control group was observed only in the microclones inoculated with 10^7 CFU/ml of *E. italicus*. On the 14th day, both experimental concentrations exhibited differences from the control group as well as from each other. On the 30th day, the average leaf area of the experimental plants was equal to the control and was 0.6 cm^2 larger.

In the case of paulownia plants, bacteria at both experimental concentrations contributed to an increase in the average leaf area, with the highest average leaf area recorded in microclones inoculated with 10^7 CFU/ml of *E. italicus* throughout the entire observation period (Fig. 10). The effects of the studied bacterial concentrations varied over the course of the experiment. Although the difference in absolute values between leaf areas in plants exposed to 10^7 and 10^6 CFU/ml of bacteria was small, the statistical analysis revealed a significant dif-




Fig. 7. The average number of nodes in the microclones after inoculation with *E. italicus* ONU547: A – *Rubus fruticosus* and B – *Paulownia tomentosa*; control plants ($n = 20$) received no treatment while treated plants ($n = 20$ for each concentration) were inoculated with bacteria; means followed by a common letter are not significantly different by the DMRT at $P \leq 0.05$




Fig. 8. Microclones on the 14th day of acclimatization with *E. italicus* ONU547: A – *Rubus fruticosus* (1–2: control, 3–4: inoculated with bacteria); B – *Paulownia tomentosa* (1–6: control, 7–12: inoculated with bacteria); control plants ($n = 20$) received no treatment while treated plants ($n = 20$) were inoculated with 10^7 CFU/ml of bacteria

ference between them. Therefore, the best effect on the average leaf area of paulownia was demonstrated by *E. italicus* at a concentration of 10^7 CFU/ml, which was 0.8 cm^2 larger than the control on the 30th day of acclimatization.

Hence, the inoculation of *R. fruticosus* and *P. tomentosa* microclones with *E. italicus* ONU547 at concentrations of 10^6 and 10^7 CFU/ml exhibited a beneficial impact on various plant parameters, including the survival rate, average plant length, node count, and leaf area, when compared to the control group. Notably, the application of bacteria at a concentration of 10^7 CFU/ml yielded the most favorable results in terms of plant acclimatization to *ex vitro* conditions.

Discussion

This study represents the first comprehensive exploration of the potential application of *E. italicus* ONU547 as a plant growth promoter and beneficial

microorganism for acclimatization. The antagonistic activity of the investigated bacteria against mycelial fungi, including *A. niger*, *C. cladosporioides*, *A. alternata*, *A. tenuissima*, *R. cerealis*, *P. expansum*, and *P. variotii*, was successfully established. These findings align with previous research conducted by other scholars, who have also detected antagonistic properties in various strains of enterococci (Kumar et al., 2016; Maany et al., 2019; Diaz et al., 2021). Notably, the authors did not find any prior reports on the antagonism of *Enterococcus* spp. against *P. variotii*, *R. cerealis*, *A. tenuissima*, and *C. cladosporioides*. However, the inhibitory effects of enterococci on other tested fungi species, such as *A. niger*, *A. alternata*, and *P. expansum*, were already documented in existing literature (Fhoula et al., 2013; Ben Braïek et al., 2018; David and Onifade, 2018; Zabouri et al., 2021), corroborating the findings of our experiments, which showcased similar antagonistic properties.




Fig. 9. The average leaf area of *Rubus fruticosus* microclones after inoculation with *E. italicus* ONU547; control plants ($n = 20$) received no treatment while treated plants ($n = 20$ for each concentration) were inoculated with bacteria; means followed by a common letter are not significantly different by the DMRT at $P \leq 0.05$




Fig. 10. The average leaf area of *Paulownia tomentosa* microclones after inoculation with *E. italicus* ONU547; control plants ($n = 20$) received no treatment while treated plants ($n = 20$ for each concentration) were inoculated with bacteria; means followed by a common letter are not significantly different by the DMRT at $P \leq 0.05$

It is worth noting that the studied bacteria did not exhibit any antagonistic activities against *F. oxysporum*. This sets *E. italicus* ONU547 apart from related strains that have demonstrated inhibitory effects on the growth of *Fusarium* spp. in other studies (Ben Braiek et al., 2018; Kalantari et al., 2019; Houicher et al., 2021). Additionally, although the bacteria did not impede the growth of *P. expansum* mycelia, they did delay sporulation during cocultivation. This suppression of sporulation has the potential to decelerate the progression of fungal diseases in plants by influencing the subsequent dissemination of spores—a critical factor in disease spread (Dahlberg and Van Etten, 1982).

In previous *in vivo* studies (Merlich et al., 2017a), it was observed that *E. italicus* ONU547 exhibited inhibitory effects against *Rhizobium radiobacter* C58 and *Erwinia carotovora* ZM1 on carrot explants, resulting in

growth reductions of 33.3 and 24.8%, respectively. These findings highlight the antagonistic potential of the microorganisms within the studied strain against both fungal and bacterial phytopathogens.

Furthermore, this study revealed the growth-promoting properties of *E. italicus* ONU547 for microclones during the transition from *in vitro* conditions to a nonsterile environment. Moreover, our experiments demonstrated the positive effects of inoculating with cell suspensions of the microorganisms rather than their purified metabolites, as often reported in the existing literature (Lee et al., 2015; Pellegrini et al., 2020; Chaurasia et al., 2022; Msimbira et al., 2022). This indicates that for the potential development of biological preparations based on *E. italicus* ONU547, there is an opportunity to save time by bypassing the isolation and purification of specific synthesized compounds and simply utilizing the cell mass. Such an approach holds promise for the creation of more cost-effective preparations.

The concentration of bacteria during inoculation has been determined to be a crucial factor in its impact on plants. For example, when watercress seeds were inoculated *in vitro*, a concentration of 10^8 CFU/ml was found to be phytotoxic, inhibiting germination and plant growth. On the other hand, concentrations of 10^6 CFU/ml and 10^7 CFU/ml exhibited growth-promoting properties. Based on measurements of root and shoot length on the 9th day of observation, a concentration of 10^6 CFU/ml was deemed more effective. However, the highest germination rate and the greatest number of roots were observed in seedlings inoculated with 10^7 CFU/ml of *E. italicus*.

In the acclimatization experiments, it was found that inoculating plants with bacteria at a concentration of 10^7 CFU/ml yielded the most significant outcomes in terms of all evaluated characteristics. A statistically significant difference from the control group was observed from the 7th day of cultivation and remained evident throughout the observation period. Bacteria at a concentration of 10^6 CFU/ml also demonstrated a positive effect, although it generally required more time to manifest and exhibited a lesser extent of the impact.

In other studies investigating the effects of enterococci on plant growth, different concentrations of bacteria, inoculation procedures, and plant species were employed, resulting in varying observed influences. For example, Chaurasia et al. (2022) inoculated soil with

Enterococcus spp. at a concentration of 8×10^5 CFU/kg, leading to maize seed germination occurring three days faster than in the control group and a 53.7% increase in plant height, while no significant difference in the number of leaves was observed. In Kumar et al.'s (2016) experiments, *Vigna mungo* seeds were inoculated with a bacterial suspension (10^8 CFU/ml) mixed with 1% carboxymethyl cellulose, resulting in a 16.7% increase in germination rate, 23.7% increase in shoot length, and 27.3% increase in root length. Lee et al. (2012) found that bacterial inoculation significantly enhanced the shoot length of rice plants but did not yield positive effects on any of the analyzed biometric characteristics of cucumber plants. Simultaneously, the cell-free extract did increase the shoot length of plants by 40%. The exact concentration of the bacterial inoculum and the specific inoculation procedure was not specified in that research.

In the present study, the inoculation of *L. sativum* L. seeds with *E. italicus* ONU547 at a concentration of 10^7 CFU/ml through a 1 h soaking process resulted in a 9.3% increase in seed germination, 50.3% increase in *in vitro* shoot length, and 21.4% increase in root length compared to the control group. Additionally, for blackberry microclones during *ex vitro* acclimatization, the shoot length, node number, and leaf area exhibited increases of 36.5, 41.9, and 37.5% respectively. Similarly, for paulownia plants, the shoot length, node number, and leaf area increased by 25.0, 45.2, and 22.2%, respectively. A comparison of the aforementioned results suggests that the interactions between enterococci and plants can be species-specific and may depend on the inoculation procedure and inoculum concentration. Therefore, further extensive research is warranted to explore this topic.

The observed effects of *E. italicus* ONU547 on seed germination and the growth of micropropagated plants during the acclimatization stage can be preliminarily explained by several factors. It has been established that *E. italicus* ONU547 is a producer of organic acids, predominantly acetic acid (Merlich et al., 2017a), which may contribute to its antagonistic activity against phytopathogens. Additionally, *E. italicus* ONU547 has been found to produce bacteriocin, which exhibits increased activity after heating and demonstrates antagonistic effects against phytopathogenic bacteria (Merlich et al., 2019).

Previous studies have indicated that *E. italicus* ONU547 can form biofilms on the surface of watercress

roots, particularly when in consortia with other lactic acid bacteria. This adhesive property plays a significant role in effective plant protection and facilitates the better penetration of bacterial metabolites into the intercellular space (Merlich et al., 2017b).

The growth-promoting properties observed in this study may be attributed to the potential synthesis of phytohormones, as similar traits have been reported for enterococci (Lee et al., 2015). Additionally, the studied strain could potentially have a positive impact on plants through nitrogen fixation and phosphate solubilization, as demonstrated by other research on enterococci (Mello et al., 2020). Further investigations will be necessary to verify these hypotheses specifically for *E. italicus* ONU547.

Conclusion

E. italicus ONU547 demonstrated antagonistic activity against various phytopathogenic fungi, including *C. cladosporioides*, *A. alternata*, *A. tenuissima*, *A. niger*, *R. cerealis*, *P. variotii*, and *P. expansum*. Furthermore, the bacteria exhibited plant growth-promoting properties in the germination assay of *L. sativum* L. seeds, leading to increased average root and shoot length, the average number of roots, and germination rate compared to the control group. Additionally, positive effects were observed in the *ex vitro* acclimatization of micropropagated *P. tomentosa* Steud. and *R. fruticosus* L. plants, including an increase in average plant length, number of nodes, leaf area, and survival rate in the soil. The optimal inoculum concentration for effective acclimatization of microclones was determined to be 10^7 CFU/ml.

Given the outcomes of these experiments and the various potential mechanisms by which *E. italicus* ONU547 impacts plant tissues, further research on these bacteria is recommended. They show promise for inoculating micropropagated plants during the acclimatization stage to *ex vitro* conditions.

Acknowledgments

The authors would like to acknowledge the assistance of the D.K. Zabolotny Institute of Microbiology and Virology of the National Academy of Sciences of Ukraine, which provided the strains of mycelial fungi for this research.

References

- Almeida-Santos A.C., Novais C., Peixe L., Freitas A.R. (2021) *Enterococcus* spp. as a producer and target of bacteriocins:

- a double-edged sword in the antimicrobial resistance crisis context. *Antibiotics* 10(10): 1215.
- Ben Braïek O., Cremonesi P., Morandi S., Smaoui S., Hani K., Ghraïri T. (2018) *Safety characterization and inhibition of fungi and bacteria by a novel multiple enterocin-producing Enterococcus lactis 4CP3 strain*. *Microb. Pathog.* 118: 32–38.
- Cesa-Luna C., Baez A., Quintero-Hernandez V., De la Cruz-Enriquez J., Castaneda-Antonio M.D., Munoz-Rojas J. (2020) *The importance of antimicrobial compounds produced by beneficial bacteria on the biocontrol of phytopathogens*. *Acta Biol. Colomb.* 25(1): 140–154.
- Chaurasia L.K., Tirwa R.K., Tamang B. (2022) *Potential of Enterococcus faecium LM5.2 for lipopeptide biosurfactant production and its effect on the growth of maize (Zea mays L.)*. *Arch. Microbiol.* 204(4): 223.
- Cosmulescu S., Scriciecu F., Manda M. (2020) *Determination of leaf characteristics in different medlar genotypes using the ImageJ program*. *Hort. Sci. (Prague)* 47: 117–121.
- Crisan L.R., Petrus-Vancea A. (2016) *Paulownia tomentosa L. in vitro propagation*. *Nat. Resour. Sust. Dev.* 6: 30–37.
- Dahlberg K.R., Van Etten J.L. (1982) *Physiology and biochemistry of fungal sporulation*. *Ann. Rev. Phytopathol.* 20: 281–301.
- David O.M., Onifade O.E. (2018) *Effect of partially purified enterocins from Enterococcus faecalis strains on the growth of some phytopathogenic fungi*. *Ruhuna J. Sci.* 9(2): 160–168.
- Diaz D., Pizzolitto R., Vazquez C., Usseglio V., Zunino M., Dambolena J., Zygado J., Merlo C. (2021) *Effects of the volatile organic compounds produced by Enterococcus spp. strains isolated from maize grain silos on Fusarium verticillioides growth and fumonisin B1 production*. *J. Stored Prod. Res.* 93: 101825.
- El-Baky N.A., Abdel Rahman R.A., Sharaf M.M., Amara A.A.A.F. (2021) *The development of a phytopathogenic fungi control trial: Aspergillus flavus and Aspergillus niger infection in jojoba tissue culture as a model*. *Sci. World J.* 2021: 6639850.
- Fathy H., El-Leel O., Amin M., Abuel-Leel O. (2018) *Micropropagation and biomass production of Rubus fruticosus L. (Blackberry) plant*. *Middle East J. Appl. Sci.* 8(4): 1215–1228.
- Fhoula I., Najjari A., Turki Y., Jaballah S., Boudabous A., Ouzari H. (2013) *Diversity and antimicrobial properties of lactic acid bacteria isolated from rhizosphere of olive trees and desert truffles of Tunisia*. *Biomed. Res. Int.* 2013: 405708.
- H-Kittikun A., Biscola V., El-Ghaish S. Jaffres E., Xavier D., Pillot G., Haertle T., Chobert J.M., Hwanhlem N. (2015) *Bacteriocin-producing Enterococcus faecalis KT2W2G isolated from mangrove forests in southern Thailand: Purification, characterization and safety evaluation*. *Food Control.* 54: 126–134.
- Houicher A., Alkay Z., Bourahla M., Dertli E. (2021) *Virulence factors, antibiotic resistance, and antimicrobial activity of Enterococcus spp. isolated from different sources in Algeria*. *J. Microb. Biotech. Food. Sci.* 11(2): e1907.
- Kalantari N., Nazarian-Firouzabadi F., Darvishnia M. (2019) *The potential of Enterococcus faecium strains isolated from Iranian Oak sap to control plant pathogens*. *Biol. Contr. Pests Plant Dis.* 8(2): 87–99.
- Kumar S., Dubey R.C., Maheshwari D. (2016) *Biosurfactant-mediated biocontrol of Macrophomina phaseolina causing charcoal rot in Vigna mungo by a plant growth promoting Enterococcus sp. BS13*. *J. Plant Pathol. Microbiol.* 7: 11.
- Lee K.E., Radhakrishnan R., Kang S.M., You Y.H., Joo G.J., Lee I.J., Kim J.H. (2015) *Enterococcus faecium LKE12 cell-free extract accelerates host plant growth via gibberellin and indole-3-acetic acid secretion*. *J. Microbiol. Biotechnol.* 25(9): 1467–1475.
- Leite M.S., Furtado-Pinto T.E., Rabelo-Centofante A., Rubio-Neto A., Guimaraes-Silva F., Goncalves-Selari P.J.R., Martins P.F. (2021) *Acclimatization of Pouteria gardeneriana Radlk micropropagated plantlets: role of in vitro rooting and plant growth-promoting bacteria*. *Curr. Plant Biol.* 27: 100209.
- Liu Z., Jiao R.L., Chen S.Y., Ren Y., Zhang L., Zhang D., Chen J.Y., Guoying L. (2022) *First report of fruit rot of grapes (Vitis vinifera) caused by Cladosporium cladosporioides in Xinjiang, China*. *Plant Dis.* 106: 315.
- Maany D., Atwa N., Ahmed I., El-Diwanly, El-Wasief A., Amer M.N. (2019) *Production of a polypeptide antimicrobial compound by lactic acid bacteria isolated from rhizospheric soil of Egyptian organic farms*. *Int. J. Appl. Pure Sci. Agric.* 5: 2.
- Mello L., Santos A., Santos R., Kandasamy S., Lazarovits G., Rigobelo E. (2020) *Application of the bacterial strains Ruminobacter amylophilus, Fibrobacter succinogenes and Enterococcus faecium for growth promotion in maize and soybean plants*. *Aust. J. Crop Sci.* 14(12): 2020–2027.
- Merlich A., Galkin M., Choiset Y., Limanska N., Vasylieva N., Ivanytsia V., Haertlé T. (2019) *Characterization of the bacteriocin produced by Enterococcus italicus ONU547 isolated from Thai fermented cabbage*. *Folia Microbiol. (Praha)* 64(4): 535–545.
- Merlich A.G., Zhunko I.D., Limanska N.V., Ivanytsia V.O. (2017) *Antagonistic activity against plant pathogenic bacteria and ability to biofilm formation of Enterococcus italicus ONU547 bacteria and their consortia with Lactobacillus plantarum*. *J. V.N. Karazin Kharkiv Nat. Univ. Ser. Biol.* 29: 71–76.
- Merlich A.G., Zhunko I.D., Limanska N.V., Ivanytsia V.O. (2017) *Antagonistic activity of metabolic products of bacteria Lactobacillus plantarum and Enterococcus italicus with joint action against phytopathogenic bacteria*. *Microbiol. Biotech.* 3: 45–54.
- Msimbira L.A., Naamala J., Antar M., Subramanian S., Smith D.L. (2022) *Effect of microbial cell-free supernatants extracted from a range of pH levels on corn (Zea mays L.) and tomato (Solanum lycopersicum L.) Seed germination and seedling growth*. *Front. Sustain. Food Syst.* 6: 789335.

- Murashige T., Skoog F. (1962) *A revised medium for rapid growth and bioassays with tobacco tissue culture*. *Physiol Plant*. 15: 473–497.
- Omamor I.B., Asemota A., Eke C.R., Eziashi, E. (2007) *Fungal contaminants of the oil palm tissue culture in Nigerian institute for oil palm research (NIFOR)*. *Afr. J. Agric. Res.* 2: 534–537.
- Panwar M., Tewari R., Nayyar H. (2016) *Native halo-tolerant plant growth promoting rhizobacteria Enterococcus and Pantoea sp. improve seed yield of Mungbean (Vigna radiata L.) under soil salinity by reducing sodium uptake and stress injury*. *Physiol. Mol. Biol. Plants*. 22(4): 445–459.
- Pellegrini M., Pagnani G., Bernardi M., Mattedi A., Spera D.M., Gallo M.D. (2020) *Cell-free supernatants of plant growth-promoting bacteria: a review of their use as bio-stimulant and microbial biocontrol agents in sustainable agriculture*. *Sustainability* 12(23): 9917.
- Raman J., Kim J.S., Choi K.R., Eun H., Yang D., Ko Y.J., Kim S.J. (2022) *Application of lactic acid bacteria (LAB) in sustainable agriculture: advantages and limitations*. *Int. J. Mol. Sci.* 23: 7784.
- Shekhawat M.S., Mehta S.R., Manokari M., Priyadarshini S., Badhepuri M.K., Jogam P., Dey A., Rajput B.S. (2021) *Morpho-anatomical and physiological changes of Indian sandalwood (Santalum album L.) plantlets in ex vitro conditions to support successful acclimatization for plant mass production*. *Plant Cell Tiss. Organ. Cult.* 147: 423–435.
- Shobha G., Kumudini B.S. (2012) *Antagonistic effect of the newly isolated PGPR Bacillus spp. on Fusarium oxysporum*. *Int. J. Appl. Sci. Eng. Res.* 1(3): 463–474.
- Strafella S., Simpson D.J., Yaghoubi Khanghahi M., De Angelis M. Ganzle, M., Minervini F., Crecchio C. (2021) *Comparative genomics and in vitro plant growth promotion and biocontrol traits of lactic acid bacteria from the wheat rhizosphere*. *Microorganisms* 9(1): 78.
- Svec P., Vandamme P., Bryndova H., Holochova P., Kosina M., Maslanova I., Sedlacek I. (2012) *Enterococcus plantarum sp. nov., isolated from plants*. *Int. J. Syst. Evol. Microbiol.* 62(7): 1499–1505.
- Tytarenko N., Tesliuk N., Ivanytsia V. (2020) *Perspectives of using bacteria for cell and tissue plant culture*. *Microbiol. Biotech.* 3: 6–31.
- Wang K., Ngea G.L.N., Godana E.A., Shi Y., Lanhuang B., Zhang X., Zhao L., Yang Q., Wang S., Zhang H. (2021) *Recent advances in Penicillium expansum infection mechanisms and current methods in controlling P. expansum in postharvest apples*. *Crit. Rev. Food. Sci. Nutr.* 20: 1–14.
- Zabouri Y., Cheriguene A., Chougrani F., Merzouk Y., Marchetta A., Urzi C., De Leo F. (2021) *Antifungal activity of lactic acid bacteria against phytopathogenic Alternaria alternata species and their molecular characterization*. *J. Food Nutr. Res.* 60(1): 18–28.